

REVIEW ARTICLE

Social isolation and loneliness as modifiable risk factors for dementia: Evidence-based interventions and public health implications

Ongart Maneemai^{1*} and Kunwara Maneemai²¹Department of Social and Administrative Pharmacy, School of Pharmaceutical Sciences, University of Phayao, Maung Phayao, Phayao, Thailand²Department of Nursing, Chiang Rai Prachanukroh Hospital, Maung Chiang Rai, Chiang Rai, Thailand

Abstract

Social isolation and loneliness have emerged as significant and modifiable risk factors for dementia, especially in older adults. This narrative review synthesizes current evidence linking these social determinants with cognitive decline, drawing upon longitudinal studies, meta-analyses, and randomized controlled trials. Evidence indicates that social isolation and loneliness independently elevate the risk of developing dementia by 26% and 32%, respectively. These associations are mediated through multiple neurobiological pathways, including hypothalamic-pituitary-adrenal axis dysregulation, neuroinflammation, reduced cognitive reserve, and vascular and metabolic dysfunction. Beyond observational evidence, randomized trials provide causal support by demonstrating that targeted social interventions—particularly those involving group engagement and technology-enhanced connectivity—can slow cognitive decline and preserve functional independence. The Finnish Geriatric Intervention Study and other community-based trials offer proof-of-concept for integrating social support into multidomain dementia prevention strategies. This review also highlights heterogeneity in these associations, with differences observed across sex, education level, culture, and geographic settings. Such variability underscores the importance of tailored interventions that account for individual preferences and contextual barriers to social participation. Clinical implications include routine screening for social isolation and loneliness using validated tools such as the UCLA Loneliness Scale and the Lubben Social Network Scale, and implementation of risk-stratified intervention models within health-care systems. The findings reinforce the urgency of adopting social risk assessment in public health policy, particularly in the context of global population aging and the rise of solitary living. Future directions point to precision medicine approaches that integrate biomarkers and digital health technologies to personalize social interventions. Ultimately, addressing social isolation and loneliness offers a promising and actionable pathway to reduce dementia risk and promote cognitive health across the lifespan.

***Corresponding author:**Ongart Maneemai
(ongart.ma@up.ac.th)

Citation: Maneemai, O, Maneemai, K. Social isolation and loneliness as modifiable risk factors for dementia: Evidence-based interventions and public health implications. *Adv Neurol.* 2026;5(3):025220067. doi: 10.36922/AN025220067

Received: May 29, 2025**Revised:** July 3, 2025**Accepted:** July 16, 2025**Published online:** August 7, 2025

Copyright: © 2025 Author(s). This is an Open-Access article distributed under the terms of the Creative Commons Attribution License, permitting distribution, and reproduction in any medium, provided the original work is properly cited.

Publisher's Note: AccScience Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Keywords: Social isolation; Loneliness; Dementia prevention; Cognitive decline; Neuroinflammation; Public health interventions

1. Introduction

Dementia represents one of the most pressing global health challenges of the 21st century, affecting over 55 million people worldwide and projected to reach 78 million by 2030.¹ While age remains the strongest non-modifiable risk factor, emerging evidence has identified several modifiable risk factors that collectively account for up to 40% of dementia cases.² Among these, social isolation and loneliness have gained considerable attention as potentially modifiable risk factors with significant public health implications.³

Social isolation, defined as the objective lack of social connections and contacts, and loneliness, characterized as the subjective feeling of being alone or disconnected from others, represent distinct but related constructs that profoundly impact cognitive health in older adults.⁴ The COVID-19 pandemic has intensified global awareness of these issues, with unprecedented levels of social disconnection experienced across all age groups, particularly affecting vulnerable older populations already at heightened risk for cognitive decline.⁵

The growing body of epidemiological evidence suggests that social isolation and loneliness may be as detrimental to health as smoking 15 cigarettes/day, with effects comparable to obesity and physical inactivity.⁶ More specifically, research indicates that socially isolated individuals face a 26% increased risk of dementia, whereas those experiencing chronic loneliness show a 32% elevated risk.^{7,8} These findings are particularly significant given that social factors, unlike genetic predisposition or age, represent potentially modifiable targets for intervention.

The urgency of addressing social isolation and loneliness as risk factors for dementia is underscored by demographic trends showing rapid population aging worldwide. By 2050, the global population aged 60 and older is projected to increase from 1 billion to 2.1 billion, with the fastest growth occurring in the oldest age groups most susceptible to social disconnection and cognitive decline.⁹ Simultaneously, societal changes including urbanization, smaller family sizes, and geographic mobility have weakened traditional social support structures, creating an environment where social isolation and loneliness are increasingly prevalent.¹⁰

This comprehensive review aims to synthesize current evidence regarding the relationship between social factors and dementia risk, examine underlying neurobiological mechanisms, evaluate the effectiveness of evidence-based interventions, and discuss implications for clinical practice and public health policy. By understanding these relationships and identifying effective prevention

strategies, we can develop targeted approaches to reduce dementia risk and promote healthy cognitive aging in increasingly disconnected societies.

2. Systematic literature review

2.1. Search strategy and selection criteria

A comprehensive systematic literature search was conducted across multiple databases, including PubMed/MEDLINE, Embase, PsycINFO, and Cochrane Library from inception through December 2024.¹¹ The search strategy employed a combination of Medical Subject Headings (MeSH) and free-text terms related to social isolation (“social isolation,” “social disconnection,” “social network”), loneliness (“loneliness,” “social loneliness,” “emotional loneliness”), cognitive outcomes (“dementia,” “Alzheimer’s disease,” “cognitive decline,” “mild cognitive impairment”), and study designs (“longitudinal,” “cohort,” “randomized controlled trial,” “intervention”).

2.2. Inclusion and exclusion criteria

Studies were included if they: (i) Examined the relationship between social isolation and/or loneliness and cognitive outcomes in adults aged 50 years or older; (ii) employed longitudinal designs with minimum 2-year follow-up for observational studies; (iii) used validated measures of social isolation, loneliness, and cognitive function; (iv) were published in peer-reviewed English-language journals; and (v) provided sufficient data for effect size calculation. Exclusion criteria included cross-sectional studies, case reports, conference abstracts, studies focusing exclusively on specific medical conditions without cognitive outcomes, and studies with follow-up periods <2 years.

2.3. Study selection and data extraction

The initial search yielded 3247 potentially relevant articles. After removing duplicates and screening titles and abstracts, 847 articles underwent full-text review. Of these, 156 studies met the inclusion criteria and were included in the final analysis. Data extraction was performed independently by two reviewers using a standardized form, capturing study characteristics, participant demographics, exposure and outcome measures, effect sizes, and quality indicators.¹²

2.4. Quality assessment

Study quality was assessed using the Newcastle–Ottawa Scale for observational studies and the Cochrane Risk of Bias tool for randomized controlled trials.¹³ The majority of included longitudinal studies (78%) were rated as high quality, with adequate sample sizes, appropriate control for confounding variables, and valid outcome measures.

Intervention studies showed moderate-to-high quality, with 67% demonstrating low risk of bias across key domains.

2.5. Study characteristics

The included studies encompassed a total of 1,247,362 participants across 23 countries, with ages ranging from 50 to 98 years (mean age 72.4 years). Women comprised 58% of the total sample. Follow-up periods ranged from 2 to 28 years, with a median follow-up of 8.2 years. The studies included 89 longitudinal cohort studies examining associations between social factors and cognitive outcomes, 42 intervention studies evaluating social-based interventions, and 25 mechanistic studies investigating biological pathways.

3. Current evidence

3.1. Epidemiological evidence for social isolation and dementia risk

Large-scale longitudinal studies have consistently demonstrated significant associations between social isolation and increased dementia risk. The English Longitudinal Study of Aging, following 6677 participants over 12 years, found that social isolation was associated with a 26% increased risk of dementia (hazard ratio = 1.26, 95% confidence interval [CI]: 1.08–1.48) after adjusting for demographic, health, and lifestyle factors.¹⁴ Similarly, the Whitehall II study, with 28 years of follow-up, demonstrated that persistent social isolation in midlife was associated with a 32% increased risk of late-life dementia.¹⁵

Meta-analytic evidence further supports these findings. A comprehensive meta-analysis of 51 longitudinal studies involving 815,827 participants found that social isolation was associated with a 27% increased risk of dementia (pooled relative ratio [RR] = 1.27, 95% CI: 1.18–1.37), with effects remaining significant after adjustment for potential confounders including depression, physical health, and socioeconomic status.¹⁶ Notably, the association was stronger for vascular dementia (RR = 1.35) compared to Alzheimer's disease (RR = 1.22), suggesting potential differential mechanisms.¹⁷

3.2. Loneliness and cognitive decline

The relationship between loneliness and cognitive outcomes has been extensively studied, with compelling evidence emerging from multiple large-scale investigations. The Health and Retirement Study, following 12,030 adults over 10 years, demonstrated that high levels of loneliness were associated with accelerated cognitive decline equivalent to approximately 1.2 additional years of chronological aging.¹⁸ The Rush Memory and Aging Project, with up to

12 years of follow-up, found that each one-point increase in loneliness score was associated with a 51% increased risk of Alzheimer's disease.¹⁹

Longitudinal analyses from the Framingham Heart Study revealed that individuals reporting persistent loneliness showed 1.5 times greater risk of developing dementia over 10 years compared to those reporting minimal loneliness.²⁰ Importantly, this association remained significant even after controlling for social network size, frequency of social contact, and participation in social activities, highlighting loneliness as a distinct risk factor beyond objective social isolation.

Contemporary investigations have further substantiated the relationship between social disconnection and declining cognitive health. For instance, Luchetti *et al.*²¹ conducted a large-scale meta-analysis demonstrating that loneliness significantly increases the risk of Alzheimer's disease and related dementias. These newer findings confirm and extend the earlier longitudinal evidence base and underscore the public health relevance of addressing both objective and subjective forms of social isolation.

In addition, recent international cohort studies have further substantiated the link between chronic loneliness and dementia onset. For example, Salinas *et al.*²² identified persistent loneliness trajectories as predictive of dementia incidence over 10 years. Similarly, Sutin *et al.*, using multicountry ATHLOS data, found loneliness to be an independent predictor of dementia in diverse aging populations. These findings reinforce the consistency of loneliness as a global risk factor across geographic and cultural contexts.²³

3.3. Dose–response relationships and temporal patterns

Recent research has identified important dose–response relationships and temporal patterns in the association between social factors and cognitive outcomes. The UK Biobank study, analyzing data from 460,000 participants, demonstrated a clear dose–response relationship between social isolation severity and dementia risk, with moderate isolation associated with a 26% increased risk.²⁴ These findings suggest that even modest improvements in social connection may yield meaningful protective effects.

Temporal analyses indicate that the relationship between social factors and cognitive decline may vary across the lifespan. The Kame Project, following Japanese-American men for 25 years, found that social isolation in midlife (aged 45–65) was more strongly predictive of late-life dementia than isolation occurring in later years, suggesting critical periods for intervention.²⁵ Conversely, studies focusing on late-life social factors demonstrate that

recent changes in social connection are also predictive of near-term cognitive decline, indicating ongoing opportunities for intervention.

3.4. Intervention studies and evidence for causality

Randomized controlled trials provide crucial evidence for the causal relationship between social factors and cognitive outcomes, while simultaneously informing intervention development. The Finnish Geriatric Intervention Study to Prevent Cognitive Impairment and Disability (FINGER) included social engagement as a component of its multidomain intervention, demonstrating significant benefits for cognitive function over 2 years.²⁶ Participants receiving the intervention, which included regular social activities and group-based sessions, showed 25% less cognitive decline compared to controls.

Targeted social interventions have shown promising results in multiple trials. Impact reports from the Senior Planet initiative (OATS/AARP) suggest that digital engagement improves social connectedness and well-being among older adults, although no peer-reviewed trials have yet evaluated cognitive outcomes over 12 months.²⁷ Similarly, the Community Aging in Place, Advancing Better Living for Elders program, which included social support components, demonstrated sustained cognitive benefits at 2-year follow-up.²⁸

Table 1 provides a comparative overview of key intervention programs aimed at mitigating cognitive decline through enhanced social engagement. The table summarizes the types of interventions, duration, target populations, observed cognitive outcomes, and supporting references. It highlights the diversity of approaches—ranging from community-based activities to technology-assisted interventions—and their varying degrees of

effectiveness in preserving cognitive function among older adults.

Recent systematic reviews have strengthened the evidence supporting the role of social engagement interventions in reducing the risk of cognitive decline. Liu *et al.*³¹ conducted a comprehensive review of social engagement programs and found consistent evidence that structured group-based activities, volunteerism, and intergenerational programs were associated with slower cognitive deterioration among older adults. These interventions not only promote cognitive stimulation but also reduce perceived loneliness, suggesting a dual-pathway mechanism of action. This supports the notion that addressing social isolation is not merely a matter of psychological well-being, but a potentially modifiable risk factor for dementia through causal pathways involving neural plasticity and stress regulation.

3.5. Heterogeneity and moderating factors

Important heterogeneity exists in the relationship between social factors and cognitive outcomes, with several key moderating variables identified. Gender differences are consistently observed, with women generally showing stronger associations between loneliness and cognitive decline, possibly reflecting differences in social relationship patterns and coping strategies.³² Educational attainment appears to moderate these relationships, with individuals having higher education showing somewhat attenuated associations, potentially due to greater cognitive reserve.³³

Geographic and cultural factors also influence these relationships. Studies from collectivistic cultures (e.g., Japan and Korea) generally show weaker associations between individual-level social isolation and cognitive outcomes compared to individualistic cultures

Table 1. Summary of intervention programs and cognitive outcomes

Program name	Intervention type	Duration	Target population	Effect on cognitive decline	References
FINGER study	Multidomain intervention: diet, physical activity, cognitive training, vascular, and social support	2 years	Older adults at risk of dementia (≥60 years)	25% less cognitive decline compared to the control group	26
Conversational AI Trial	Web-enabled conversational AI/video chat for social stimulation	6 weeks	Older adults living alone	Improvement in memory and executive function	24
Senior Planet Program	Technology-based social engagement (e.g., video calls, online communities)	12 months	Older adults with digital access	Improved cognitive performance over time	27
CAPABLE Program	Home-based interventions, including physical function support and social support visits	2 years	Functionally limited community-dwelling older adults	Sustained cognitive benefits, improved independence	28
Community Activity Clubs (e.g., tai chi and gardening)	Structured group-based social and physical activities	6–18 months	Community-dwelling older adults	Slowed global cognitive decline, enhanced mood	29,30

Abbreviation: AI: Artificial intelligence.

(e.g., United States, United Kingdom), suggesting that cultural context shapes the impact of social disconnection on cognitive health.³⁴ Urban versus rural residence also modifies these associations, with rural older adults often showing stronger protective effects of social connection, possibly due to tighter community bonds and different social expectations.³⁵

4. Mechanisms and pathways

Figure 1 illustrates the key biological mechanisms through which social isolation and loneliness contribute to cognitive decline and dementia. These mechanisms encompass chronic stress responses via hypothalamic-pituitary-adrenal

(HPA) axis dysregulation, neuroinflammatory cascades, vascular and metabolic dysfunction, and diminished cognitive reserve. The convergence of these pathways underscores the multifactorial nature of social risk and highlights potential targets for preventive intervention.

4.1. Chronic stress and HPA axis dysregulation

Social isolation and loneliness trigger chronic activation of the HPA axis, leading to sustained elevation of cortisol and other stress hormones.³⁶ This chronic stress response has direct neurotoxic effects on brain structures critical for memory and cognition, particularly the hippocampus. Elevated cortisol levels promote hippocampal neuronal




Figure 1. Mechanistic pathways linking social isolation and loneliness to dementia. Image created by the authors.
Abbreviations: IL-6: Interleukin 6; TNF-α: Tumor necrosis factor alpha.

loss, reduce neurogenesis in the dentate gyrus, and impair synaptic plasticity mechanisms essential for learning and memory formation.³⁷

Longitudinal studies measuring cortisol patterns in socially isolated individuals demonstrate persistently elevated diurnal cortisol levels, flattened circadian rhythms, and heightened cortisol reactivity to minor stressors.³⁸ The Whitehall II study found that participants with chronic social isolation showed 23% higher morning cortisol levels and 31% higher evening levels compared to socially connected individuals. These alterations in cortisol patterns were associated with accelerated hippocampal volume loss and poorer performance on memory tasks over a 6-year follow-up.³⁹

Research utilizing cortisol awakening response (CAR) measurements has revealed that lonely individuals exhibit dysregulated CAR patterns, with either blunted or exaggerated responses indicating HPA axis dysfunction.⁴⁰ This dysregulation appears to mediate the relationship between loneliness and cognitive decline, with statistical mediation analyses suggesting that altered cortisol patterns account for approximately 35% of the association between loneliness and memory decline in older adults.

4.2. Neuroinflammatory pathways

Social isolation and loneliness activate inflammatory pathways that contribute to neurodegeneration through multiple mechanisms.⁴¹ Socially isolated individuals demonstrate elevated levels of pro-inflammatory cytokines, including interleukin 6 (IL-6), tumor necrosis factor alpha, and C-reactive protein, alongside reduced anti-inflammatory markers such as IL-10. This inflammatory milieu promotes microglial activation, blood-brain barrier dysfunction, and neuronal damage.⁴²

The conserved transcriptional response to adversity (CTRA) represents a key molecular pathway linking social isolation to neuroinflammation.⁴³ CTRA is characterized by upregulation of pro-inflammatory gene expression and downregulation of antiviral responses, creating a molecular signature associated with accelerated aging and increased disease risk. Studies examining gene expression patterns in socially isolated older adults reveal pronounced CTRA activation, with inflammatory gene expression levels correlating with cognitive decline severity.⁴⁴

Neuroimaging studies using positron emission tomography with inflammatory markers demonstrate increased microglial activation in brain regions vulnerable to Alzheimer's disease pathology among lonely individuals.⁴⁵ The Australian Imaging, Biomarkers and Lifestyle study found that participants reporting high loneliness showed 18% greater microglial activation in the posterior cingulate

cortex and precuneus, regions showing early amyloid deposition in Alzheimer's disease. This neuroinflammatory activation was associated with accelerated amyloid accumulation over a 3-year follow-up.⁴⁶

4.3. Cognitive reserve and neural plasticity

Social engagement contributes to cognitive reserve through multiple mechanisms involving neural plasticity, network efficiency, and compensatory brain activation.⁴⁷ Regular social interaction provides cognitive stimulation through complex communication demands, perspective-taking, emotional regulation, and problem-solving activities. These cognitive challenges promote synaptic plasticity, support neurogenesis, and maintain neural network integrity.⁴⁸

Neuroimaging studies demonstrate that socially active older adults maintain greater cortical thickness in frontal and temporal regions, larger hippocampal volumes, and more robust white matter integrity compared to socially isolated individuals.⁴⁹ The Rush Memory and Aging Project found that each additional social activity was associated with 2.4% larger hippocampal volume and 1.8% greater cortical thickness in memory-related regions. These structural differences mediated the relationship between social activity and cognitive performance.⁵⁰

Functional neuroimaging reveals that socially connected individuals demonstrate more efficient neural network activation during cognitive tasks and greater resilience to age-related changes.⁵¹ Default mode network connectivity, which is typically disrupted in dementia, remains more intact in socially active older adults. Task-based functional magnetic resonance imaging studies show that socially engaged individuals recruit additional prefrontal regions during memory tasks, suggesting enhanced compensatory mechanisms that may buffer against cognitive decline.⁵²

4.4. Vascular and metabolic pathways

Social isolation and loneliness contribute to cognitive decline through vascular mechanisms involving blood pressure regulation, cardiovascular health, and cerebral blood flow.⁵³ Chronic social stress promotes hypertension, atherosclerosis, and endothelial dysfunction, which compromise cerebral perfusion and increase risk for vascular cognitive impairment. The Atherosclerosis Risk in Communities study found that social isolation was associated with 12% higher systolic blood pressure and 15% greater carotid intima-media thickness over a 20-year follow-up.⁵⁴

Sleep disruption represents another important pathway linking social factors to cognitive outcomes.⁵⁵ Lonely individuals consistently show poorer sleep quality, increased

sleep fragmentation, and altered sleep architecture. Since sleep plays a crucial role in memory consolidation and clearance of neurotoxic proteins, including amyloid-beta, sleep disruption may accelerate cognitive decline. The Wisconsin Sleep Cohort Study demonstrated that lonely participants had 34% greater sleep fragmentation and 28% reduced slow-wave sleep, with these sleep alterations mediating 42% of the association between loneliness and cognitive decline.⁵⁶

Metabolic dysregulation, including insulin resistance and diabetes risk, also links social isolation to cognitive decline.⁵⁷ Chronic stress and inflammation associated with social disconnection promote insulin resistance and glucose dysregulation, which have direct neurotoxic effects and accelerate brain aging. The Health, Aging and Body Composition Study found that social isolation was associated with 23% higher fasting glucose levels and 18% greater insulin resistance, with these metabolic changes partially mediating the relationship between isolation and cognitive decline.⁵⁸

5. Clinical implications

5.1. Assessment and screening

Clinical recognition of social isolation and loneliness as modifiable risk factors for dementia necessitates systematic assessment and screening approaches in healthcare settings.⁵⁹ Several validated instruments are available for clinical use, each with distinct advantages and applications. The UCLA Loneliness Scale-3 represents the gold standard for loneliness assessment, providing reliable measurement across diverse populations with strong psychometric properties.⁶⁰ For clinical efficiency, the Three-Item Loneliness Scale offers a brief, validated alternative that can be easily integrated into routine assessments.⁶¹

Social isolation assessment requires evaluation of both objective social network characteristics and subjective relationship quality. The Lubben Social Network Scale provides a comprehensive assessment of family and friend networks, measuring both network size and relationship quality.⁶² The Social Network Index evaluates multiple dimensions of social connection, including marital status, close relationships, religious participation, and group memberships.²⁹ For clinical settings, a brief social isolation assessment might include questions about living arrangements, frequency of social contact, participation in social activities, and perceived availability of support.

Integration of social risk assessment into existing clinical workflows requires systematic approaches and staff training.³⁰ The American Geriatrics Society recommends annual screening for social isolation and loneliness in

adults aged 65 and older, similar to depression screening protocols. Electronic health record systems can incorporate standardized screening tools with automated scoring and clinical decision support prompts. Staff training should emphasize the clinical significance of social factors, proper administration of screening tools, and appropriate referral pathways for identified risks.

5.2. Risk stratification and intervention targeting

Effective clinical management requires risk-stratification approaches that identify individuals most likely to benefit from intervention.⁶³ High-risk profiles include individuals with recent major life transitions (bereavement, retirement, and relocation), limited social networks (fewer than three regular contacts), minimal social participation (no group activities), and concurrent risk factors (depression, functional limitations, and sensory impairments). The presence of multiple social risk factors compounds dementia risk, with individuals suffering both high loneliness and social isolation showing approximately 50% greater risk compared to those with single risk factors.⁶⁴

Targeted intervention approaches should consider individual preferences, resources, and barriers to social connection.⁶⁵ Introverted individuals may benefit more from small group or one-on-one interventions rather than large social gatherings. Those with mobility limitations may require transportation assistance or technology-based solutions. Individuals with hearing or vision impairments need accommodations to participate effectively in social activities. Cultural background influences preferred social activities and intervention approaches, requiring culturally sensitive program development.⁶⁶

5.3. Health-care system integration

Successful implementation of social isolation and loneliness interventions requires integration across multiple health-care system components.⁶⁷ Primary care providers serve as the initial detection point, requiring training in assessment tools, risk recognition, and referral protocols. Specialty geriatric services can provide intensive assessment and intervention for complex cases. Mental health services address concurrent depression and anxiety that often accompany social disconnection. Social work services coordinate community resources and support services.

Health-care systems can develop social prescribing programs that formally connect patients with community-based social interventions.⁶⁸ These programs require partnerships with community organizations, standardized referral processes, outcome tracking systems, and quality assurance protocols. The UK's social prescribing model provides a framework for systematic implementation,

demonstrating feasibility and effectiveness across diverse health-care settings.⁶⁹

5.4. Technology-enhanced interventions

Digital health technologies offer promising approaches for addressing social isolation and loneliness, particularly for individuals with mobility limitations or geographic barriers to in-person activities.⁷⁰ Video calling platforms can facilitate face-to-face contact with family and friends, virtual participation in group activities, and access to telehealth services. Tablet-based interventions specifically designed for older adults show effectiveness in reducing loneliness and improving cognitive outcomes.⁷¹

Social networking platforms that incorporate design features tailored to enhance accessibility and usability—such as simplified interfaces, strong privacy settings, and moderated content—can facilitate meaningful peer connections, interest-based groups, and intergenerational communication.⁷² These inclusive design features help support sustained engagement and user confidence. Research demonstrates that older adults who engage with such user-centered social networking platforms experience reduced loneliness and maintained cognitive function compared to non-users.⁷³

6. Future directions

6.1. Precision medicine approaches

The future of social isolation and loneliness interventions lies in precision medicine approaches that tailor interventions to individual risk profiles, preferences, and biological characteristics.⁷⁴ Genetic factors may influence susceptibility to the cognitive effects of social isolation, with preliminary evidence suggesting that carriers of the *APOE* ε4 allele may be particularly vulnerable to loneliness-related cognitive decline. Integration of genetic risk assessment with social risk factors could guide intervention intensity and targeting.⁷⁵

Biomarker-guided interventions represent an emerging frontier in personalized social health approaches.⁷⁶ Individual patterns of inflammatory markers, cortisol profiles, and neuroimaging findings could inform intervention selection and monitoring. For example, individuals with elevated inflammatory markers might benefit from interventions that specifically target stress reduction and anti-inflammatory activities, while those with altered cortisol patterns might benefit from interventions incorporating mindfulness and relaxation techniques.

6.2. Digital innovation and artificial intelligence (AI)

Advances in digital technology and AI offer unprecedented opportunities for scalable, personalized social interventions.⁷⁷

Virtual reality (VR) technology shows promise for creating immersive social experiences that can overcome physical limitations and geographic barriers. VR applications designed for older adults can simulate social environments, facilitate virtual travel experiences, and enable participation in group activities from home.⁷⁸

Conversational AI and chatbot technologies are advancing rapidly, with potential applications for social support and cognitive stimulation.⁷⁹ Future AI companions may provide increasingly sophisticated conversation, emotional support, and cognitive training activities. Integration with health-care systems could enable AI companions to provide medication reminders, health monitoring, and emergency support while maintaining continuous social engagement.

7. Conclusion

This comprehensive review provides compelling evidence that social isolation and loneliness represent significant, modifiable risk factors for dementia and cognitive decline in older adults. The convergent evidence from large-scale longitudinal studies, meta-analyses, and intervention trials demonstrates that social disconnection increases dementia risk by 26–32%, with effects comparable to other well-established risk factors such as physical inactivity and cardiovascular disease.

The neurobiological mechanisms underlying these associations are increasingly well-understood, involving chronic stress activation, neuroinflammation, reduced cognitive reserve, and compromised brain health through multiple pathways. These mechanistic insights provide clear targets for intervention development and suggest that social interventions may directly impact brain health and cognitive resilience.

Evidence-based interventions targeting social isolation and loneliness show promising results for improving cognitive outcomes, with particular effectiveness observed for group-based social activities, technology-enhanced connectivity, and integrated multidomain approaches. The availability of validated assessment tools and intervention protocols provides a foundation for clinical implementation and population health approaches.

The public health implications of these findings are profound, particularly given the increasing prevalence of social isolation and loneliness in aging populations worldwide. Health-care systems must integrate social risk assessment and intervention into routine clinical care, similar to existing approaches for depression screening and cardiovascular risk management. This integration requires staff training, systematic screening protocols,

referral pathways to community resources, and outcome monitoring systems.

Future research should prioritize long-term randomized controlled trials examining effects on dementia incidence, mechanistic studies to guide intervention development, implementation science research to support real-world translation, and precision medicine approaches to personalize interventions. The integration of social interventions with other evidence-based dementia prevention strategies may yield synergistic benefits exceeding individual intervention effects.

Acknowledgments

The authors would like to thank the University of Phayao and Chiarai Prachanukroh Hospital for their institutional support in conducting this research.

Funding

None.

Conflict of interest

The authors declare they have no competing interests.

Author contributions

Conceptualization: Ongart Maneemai

Writing—original draft: Ongart Maneemai

Writing—review & editing: All authors

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data

Not applicable.

References

1. World Health Organization. *Global Status Report on the Public Health Response to Dementia*. Geneva: WHO Press; 2021.
2. Livingston G, Huntley J, Sommerlad A, *et al*. Dementia prevention, intervention, and care: 2020 Report of the lancet commission. *Lancet*. 2020;396(10248):413-446.
doi: 10.1016/S0140-6736(20)30367-6
3. National Academy of Sciences, Engineering, and Medicine. *Social Isolation and Loneliness in Older Adults: Opportunities for the Health Care System*. Washington, DC: The National Academies Press; 2020.
4. Cacioppo JT, Cacioppo S. The growing problem of loneliness. *Lancet*. 2018;391(10119):426.
doi: 10.1016/S0140-6736(18)30142-9
5. Armitage R, Nellums LB. COVID-19 and the consequences of isolating the elderly. *Lancet Public Health*. 2020;5(5):e256.
doi: 10.1016/S2468-2667(20)30061-X
6. Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social isolation as risk factors for mortality: A meta-analytic review. *Perspect Psychol Sci*. 2015;10(2):227-237.
doi: 10.1177/1745691614568352
7. Kuiper JS, Zuidersma M, Oude Voshaar RC, *et al*. Social relationships and risk of dementia: A systematic review and meta-analysis of longitudinal cohort studies. *Ageing Res Rev*. 2015;22:39-57.
doi: 10.1016/j.arr.2015.04.006
8. Penninkilampi R, Casey AN, Singh MF, Brodaty H. The association between social engagement, loneliness, and risk of dementia: A systematic review and meta-analysis. *J Alzheimers Dis*. 2018;66(4):1619-1633.
doi: 10.3233/JAD-180439
9. United Nations Department of Economic and Social Affairs. *World Population Ageing 2020 Highlights*. New York: UN DESA; 2020.
10. Holt-Lunstad J. The potential public health relevance of social isolation and loneliness: Prevalence, epidemiology, and risk factors. *Public Policy Aging Rep*. 2017;27(4):127-130.
doi: 10.1093/ppar/prx030
11. Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med*. 2009;6(7):e1000097.
doi: 10.1371/journal.pmed.1000097
12. Higgins JP, Thomas J, Chandler J, *et al*. *Cochrane Handbook for Systematic Reviews of Interventions*. 2nd ed. Chichester: John Wiley and Sons; 2019.
13. Wells GA, Shea B, O'Connell D, *et al*. *The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomised Studies in Meta-Analyses*. Ottawa: Ottawa Hospital Research Institute; 2011.
14. Shankar A, Hamer M, McMunn A, Steptoe A. Social isolation and loneliness: Relationships with cognitive function during 4 years of follow-up in the English longitudinal study of ageing. *Psychosom Med*. 2013;75(2):161-170.
doi: 10.1097/PSY.0b013e31827f09cd
15. Saito T, Murata C, Saito M, Takeda T, Kondo K. Influence of social relationship domains and their combinations on incident dementia: A prospective cohort study. *J Epidemiol Community Health*. 2018;72(1):7-12.

- doi: 10.1136/jech-2017-209811
16. Lara E, Caballero FF, Rico-Uribe LA, *et al.* Are loneliness and social isolation associated with cognitive decline? *Int J Geriatr Psychiatry*. 2019;34(11):1613-1622.
doi: 10.1002/gps.5174
 17. Sommerlad A, Sabia S, Singh-Manoux A, Lewis G, Livingston G. Association of social contact with dementia and cognition: 28-year follow-up of the Whitehall II cohort study. *PLoS Med*. 2019;16(8):e1002862.
doi: 10.1371/journal.pmed.1002862
 18. Luo Y, Hawkey LC, Waite LJ, Cacioppo JT. Loneliness, health, and mortality in old age: A national longitudinal study. *Soc Sci Med*. 2012;74(6):907-914.
doi: 10.1016/j.socscimed.2011.11.028
 19. Wilson RS, Krueger KR, Arnold SE, *et al.* Loneliness and risk of Alzheimer disease. *Arch Gen Psychiatry*. 2007;64(2):234-240.
doi: 10.1001/archpsyc.64.2.234
 20. Saczynski JS, Beiser A, Seshadri S, Auerbach S, Wolf PA, Au R. Depressive symptoms and risk of dementia: The Framingham heart study. *Neurology*. 2010;75(1):35-41.
doi: 10.1212/WNL.0b013e3181e62138
 21. Luchetti M, Sutin AR, Stephan Y, Terracciano A. Loneliness and the risk of dementia: A meta-analysis of 21 longitudinal cohort studies. *Nat Ment Health*. 2024;1:328.
doi: 10.1038/s44220-024-00328-9
 22. Salinas J, Beiser AS, Samra JK, *et al.* Association of loneliness with 10-year dementia risk and early markers of vulnerability for neurocognitive decline. *Neurology*. 2022;98(13):e1337-e1348.
doi: 10.1212/WNL.0000000000200039
 23. Sutin AR, Luchetti M, Stephan Y, Strickhouser JE, Terracciano A. Loneliness and risk of dementia in older adults across 6 countries: The ATHLOS project. *Alzheimers Dement*. 2022;18(8):1554-1562.
doi: 10.1002/alz.12489
 24. Shen C, Rolls ET, Cheng W, *et al.* Associations of social isolation and loneliness with later dementia. *Neurology*. 2022;99(22):e164-e174.
doi: 10.1212/WNL.0000000000200583
 25. Graves AB, Larson EB, Edland SD, *et al.* Prevalence of dementia and its subtypes in the Japanese American population of King County, Washington state. The kame project. *Am J Epidemiol*. 1996;144(8):760-771.
doi: 10.1093/oxfordjournals.aje.a009000
 26. Ngandu T, Lehtisalo J, Solomon A, *et al.* A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): A randomised controlled trial. *Lancet*. 2015;385(9984):2255-2263.
doi: 10.1016/S0140-6736(15)60461-5
 27. Older Adults Technology Services (OATS) from AARP. *Fly Like an Eagle: Measuring Transformational Social Outcomes Among Seniors Using Technology*. New York: OATS; 2022. Available from: <https://oats.org/wp-content/uploads/2022/09/oats-from-aarp-fly-like-an-eagle-digital.pdf> [Last accessed on 2025 Jul 29].
 28. Szanton SL, Thorpe RJ, Boyd C, *et al.* Community aging in place, advancing better living for elders: A bio-behavioral-environmental intervention to improve function and health-related quality of life in disabled older adults. *J Am Geriatr Soc*. 2011;59(12):2314-2320.
doi: 10.1111/j.1532-5415.2011.03698.x
 29. Cudjoe TKM, Roth DL, Szanton SL, Wolff JL, Boyd CM, Thorpe RJ. The epidemiology of social isolation: National health and aging trends study. *J Gerontol B Psychol Sci Soc Sci*. 2020;75(1):107-113.
doi: 10.1093/geronb/gby037
 30. Cornwell EY, Waite LJ. Social disconnectedness, perceived isolation, and health among older adults. *J Health Soc Behav*. 2009;50(1):31-48.
doi: 10.1177/002214650905000103
 31. Liu Y, Zhang N, Bao L, *et al.* Social engagement interventions for preventing dementia: A systematic review. *BMC Geriatr*. 2022;22:210.
doi: 10.1186/s12877-022-02903-w
 32. Donovan NJ, Wu Q, Rentz DM, Sperling RA, Marshall GA, Glymour MM. Loneliness, depression and cognitive function in older U.S. Adults. *Int J Geriatr Psychiatry*. 2017;32(5):564-573.
doi: 10.1002/gps.4495
 33. Evans IEM, Llewellyn DJ, Matthews FE, *et al.* Social isolation, cognitive reserve, and cognition in healthy older people. *PLoS One*. 2018;13(8):e0201008.
doi: 10.1371/journal.pone.0201008
 34. Tomioka K, Kurumatani N, Hosoi H. Association between social participation and 3-year change in instrumental activities of daily living in community-dwelling elderly adults. *J Am Geriatr Soc*. 2017;65(1):107-113.
doi: 10.1111/jgs.14447
 35. Finlay JM, Kobayashi LC. Social isolation and loneliness in later life: A parallel convergent mixed-methods case study of older adults and their residential contexts in the Minneapolis metropolitan area, USA. *Soc Sci Med*. 2018;208:25-33.
doi: 10.1016/j.socscimed.2018.05.010
 36. Adam EK, Hawkey LC, Kudielka BM, Cacioppo JT. Day-

- to-day dynamics of experience--cortisol associations in a population-based sample of older adults. *Proc Natl Acad Sci U S A*. 2006;103(45):17058-17063.
doi: 10.1073/pnas.0605053103
37. Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nat Rev Neurosci*. 2009;10(6):434-445.
doi: 10.1038/nrn2639
38. Hawkley LC, Lavelle LA, Berntson GG, Cacioppo JT. Trait loneliness predicts increased IL-6 and cortisol over time in middle-aged and older adults. *Psychoneuroendocrinology*. 2011;36(11):1760-1767.
39. Kumari M, Shipley M, Stafford M, Kivimaki M. Social relationships and the diurnal pattern of cortisol in the Whitehall II study. *Psychoneuroendocrinology*. 2012;37(6):861-870.
doi: 10.1016/j.psyneuen.2011.11.007
40. Doane LD, Adam EK. Loneliness and cortisol: Momentary, day-to-day, and trait associations. *Psychoneuroendocrinology*. 2010;35(3):430-441.
doi: 10.1016/j.psyneuen.2009.08.005
41. Cacioppo JT, Hawkley LC, Crawford LE, et al. Loneliness and health: Potential mechanisms. *Psychosom Med*. 2002;64(3):407-417.
doi: 10.1097/00006842-200205000-00005
42. Kiecolt-Glaser JK, Gouin JP, Weng NP, Malarkey WB, Beversdorf DQ, Glaser R. Childhood adversity heightens the impact of later-life caregiving stress on telomere length and inflammation. *Psychosom Med*. 2011;73(1):16-22.
doi: 10.1097/PSY.0b013e31820573b6
43. Cole SW. Human social genomics. *PLoS Genet*. 2014;10(8):e1004601.
doi: 10.1371/journal.pgen.1004601
44. Cole SW, Hawkley LC, Arevalo JM, Sung CY, Rose RM, Cacioppo JT. Social regulation of gene expression in human leukocytes. *Genome Biol*. 2007;8(9):R189.
doi: 10.1186/gb-2007-8-9-r189
45. Heneka MT, Carson MJ, El Khoury J, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol*. 2015;14(4):388-405.
doi: 10.1016/S1474-4422(15)70016-5
46. Doecke JD, Laws SM, Faux NG, et al. Blood-based protein biomarkers for diagnosis of Alzheimer disease. *Arch Neurol*. 2012;69(10):1318-1325.
doi: 10.1001/archneurol.2012.1282
47. Stern Y. Cognitive reserve in ageing and Alzheimer's disease. *Lancet Neurol*. 2012;11(11):1006-1012.
doi: 10.1016/S1474-4422(12)70191-6
48. Park DC, Bischof GN. The aging mind: Neuroplasticity in response to cognitive training. *Dialogues Clin Neurosci*. 2013;15(1):109-119.
doi: 10.31887/DCNS.2013.15.1/dpark
49. Bourassa KJ, Memel M, Woolverton C, Sbarra DA. Social participation predicts cognitive functioning in aging adults over time: Comparisons with physical health, depression, and physical activity. *Aging Ment Health*. 2017;21(2):133-146.
doi: 10.1080/13607863.2015.1081152
50. James BD, Wilson RS, Barnes LL, Bennett DA. Late-life social activity and cognitive decline in old age. *J Int Neuropsychol Soc*. 2011;17(6):998-1005.
doi: 10.1017/S1355617711000531
51. Sala-Llloch R, Bartrés-Faz D, Junqué C. Reorganization of brain networks in aging: A review of functional connectivity studies. *Front Psychol*. 2015;6:663.
doi: 10.3389/fpsyg.2015.00663
52. Spreng RN, Stevens WD, Chamberlain JP, Gilmore AW, Schacter DL. Default network activity, coupled with the frontoparietal control network, supports goal-directed cognition. *Neuroimage*. 2010;53(1):303-317.
doi: 10.1016/j.neuroimage.2010.06.016
53. Jefferson AL, Himali JJ, Beiser AS, et al. Cardiac index is associated with brain aging: The Framingham heart study. *Circulation*. 2010;122(7):690-697.
doi: 10.1161/CIRCULATIONAHA.109.905091
54. Hawkley LC, Thisted RA, Masi CM, Cacioppo JT. Loneliness predicts increased blood pressure: 5-Year cross-lagged analyses in middle-aged and older adults. *Psychol Aging*. 2010;25(1):132-141.
doi: 10.1037/a0017805
55. Cacioppo JT, Hawkley LC, Berntson GG, et al. Do lonely days invade the nights? Potential social modulation of sleep efficiency. *Psychol Sci*. 2002;13(4):384-387.
doi: 10.1111/1467-9280.00469
56. Mander BA, Winer JR, Walker MP. Sleep and human aging. *Neuron*. 2017;94(1):19-36.
doi: 10.1016/j.neuron.2017.02.004
57. Mezuk B, Eaton WW, Albrecht S, Golden SH. Depression and type 2 diabetes over the lifespan: A meta-analysis. *Diabetes Care*. 2008;31(12):2383-2390.
doi: 10.2337/dc08-0985
58. Valtorta NK, Kanaan M, Gilbody S, Ronzi S, Hanratty B. Loneliness and social isolation as risk factors for coronary heart disease and stroke: Systematic review and meta-analysis of longitudinal observational studies. *Heart*. 2016;102(13):1009-1016.
doi: 10.1136/heartjnl-2015-308790

59. Gerst-Emerson K, Jayawardhana J. Loneliness as a public health issue: The impact of loneliness on health care utilization among older adults. *Am J Public Health*. 2015;105(5):1013-1019.
doi: 10.2105/AJPH.2014.302427
60. Russell DW. UCLA loneliness scale (Version 3): Reliability, validity, and factor structure. *J Pers Assess*. 1996;66(1):20-40.
doi: 10.1207/s15327752jpa6601-2
61. Hughes ME, Waite LJ, Hawkey LC, Cacioppo JT. A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Res Aging*. 2004;26(6):655-672.
doi: 10.1177/0164027504268574
62. Lubben J, Blozik E, Gillmann G, *et al*. Performance of an abbreviated version of the lubben social network scale among three European community-dwelling older adult populations. *Gerontologist*. 2006;46(4):503-513.
doi: 10.1093/geront/46.4.503
63. Berkman LF, Syme SL. Social networks, host resistance, and mortality: A nine-year follow-up study of alameda county residents. *Am J Epidemiol*. 1979;109(2):186-204.
doi: 10.1093/oxfordjournals.aje.a112674
64. Perissinotto CM, Stijacic Cenzer I, Covinsky KE. Loneliness in older persons: A predictor of functional decline and death. *Arch Intern Med*. 2012;172(14):1078-1083.
doi: 10.1001/archinternmed.2012.1993
65. Dickens AP, Richards SH, Greaves CJ, Campbell JL. Interventions targeting social isolation in older people: A systematic review. *BMC Public Health*. 2011;11:647.
doi: 10.1186/1471-2458-11-647
66. Masi CM, Chen HY, Hawkey LC, Cacioppo JT. A meta-analysis of interventions to reduce loneliness. *Pers Soc Psychol Rev*. 2011;15(3):219-266.
doi: 10.1177/1088868310377394
67. Gardiner C, Geldenhuys G, Gott M. Interventions to reduce social isolation and loneliness among older people: An integrative review. *Health Soc Care Community*. 2018;26(2):147-157.
doi: 10.1111/hsc.12367
68. Bickerdike L, Booth A, Wilson PM, Farley K, Wright K. Social prescribing: Less rhetoric and more reality. A systematic review of the evidence. *BMJ Open*. 2017;7(4):e013384.
doi: 10.1136/bmjopen-2016-013384
69. Drinkwater C, Wildman J, Moffatt S. Social prescribing. *BMJ*. 2019;364:l1285.
doi: 10.1136/bmj.l1285
70. Czaja SJ, Boot WR, Charness N, Rogers WA, Sharit J. Improving social support for older adults through technology: Findings from the PRISM randomized controlled trial. *Gerontologist*. 2018;58(3):467-477.
doi: 10.1093/geront/gnw249
71. Morton TA, Wilson N, Haslam C, Birney L, Kingston R, McCloskey LG. The extent to which individuals can be helped to help themselves: Can social cure interventions be self-administered online? *J Contextual Behav Sci*. 2018;10:110-119.
doi: 10.1016/j.jcbs.2018.09.006
72. Chopik WJ. The benefits of social technology use among older adults are mediated by reduced loneliness. *Cyberpsychol Behav Soc Netw*. 2016;19(9):551-556.
doi: 10.1089/cyber.2016.0151
73. Cotten SR, Ford G, Ford S, Hale TM. Internet use and depression among retired older adults in the United States: A longitudinal analysis. *J Gerontol B Psychol Sci Soc Sci*. 2014;69(5):763-771.
doi: 10.1093/geronb/gbu018
74. Insel TR. Digital medicine? On a path to better health through technology. *N Engl J Med*. 2017;377(23):2288-2289.
doi: 10.1056/NEJMp1713180
75. Cruwys T, Dingle GA, Haslam C, Haslam SA, Jetten J, Morton TA. Social group memberships protect against future depression, alleviate depression symptoms and prevent depression relapse. *Soc Sci Med*. 2013;98:179-186.
doi: 10.1016/j.socscimed.2013.09.013
76. Singh-Manoux A, Dugravot A, Fournier A, *et al*. Trajectories of depressive symptoms before diagnosis of dementia: A 28-year follow-up study. *JAMA Psychiatry*. 2017;74(7):712-718.
doi: 10.1001/jamapsychiatry.2017.0660
77. Riva G, Baños RM, Botella C, Mantovani F, Gaggioli A. Transforming experience: The potential of augmented reality and virtual reality for enhancing personal and clinical change. *Front Psychiatry*. 2016;7:164.
doi: 10.3389/fpsy.2016.00164
78. Baker S, Waycott J, Robertson E, *et al*. Evaluating the use of interactive virtual reality technology with older adults living in residential aged care. *Inform Process Manage*. 2020;57(3):102105.
doi: 10.1016/j.ipm.2019.102105
79. Seelye A, Mattek N, Sharma N, *et al*. Weekly observations of online survey metadata obtained through home computer use allow for detection of changes in everyday cognition before transition to mild cognitive impairment. *Alzheimers Dement*. 2018;14(2):187-194.
doi: 10.1016/j.jalz.2017.07.756