

REVIEW ARTICLE

Targeted therapy resistance in EGFR-mutant non-small cell lung cancer: Roles of MET, Trop-2-directed antibody–drug conjugates, and molecular monitoring

Kai Yan^{1†}, Kewei Tian^{1†} , Lailing Du¹ , Xinling Zhang¹ , Jinghua Yuan¹, Xiangquan Song^{2*}, and Xiaoping Li^{1*} 

¹Key Laboratory of Artificial Organs and Computational Medicine in Zhejiang Province, Shulan International Medical College, Zhejiang Shuren University, Hangzhou, Zhejiang, China

²Hangzhou Xihu District Hospital of Traditional Chinese and Western Medicine, Hangzhou, Zhejiang, China

Abstract

Introduction: Epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors (TKIs) have substantially improved outcomes in patients with EGFR-mutant non-small cell lung cancer (NSCLC), but acquired resistance remains a major barrier to durable disease control.

Objective: This narrative review summarizes the major mechanisms of EGFR-TKI resistance and evaluates the roles of mesenchymal–epithelial transition factor (MET)-targeted therapy, trophoblast cell-surface antigen 2 (Trop-2)-directed antibody–drug conjugates (ADCs), and molecular monitoring in post-resistance management.

Methods: Relevant randomized controlled trials, translational studies, and consensus guidelines on targeted therapy resistance in NSCLC were identified from PubMed and Web of Science, with emphasis on recent high-quality evidence and seminal studies.

Results: EGFR-TKI resistance arises through on-target EGFR alterations, bypass pathway activation, histologic transformation, and non-genetic adaptation. *MET* amplification is a clinically relevant bypass mechanism, particularly after third-generation EGFR-TKI exposure, and may support combined EGFR/*MET* inhibition when confirmed using assay-specific criteria. Trop-2-directed ADCs provide a non-genotype-dependent treatment option for patients with heterogeneous or non-actionable resistance, although Trop-2 expression is not yet a validated universal predictive biomarker. Plasma circulating tumor DNA can facilitate rapid detection of emerging resistance, but negative or discordant plasma findings should be interpreted together with tissue biopsy, disease distribution, and the possibility of histologic transformation. Minimal residual disease assessment remains investigational for treatment-guided decision-making.

Conclusion: Post-EGFR-TKI management should follow a mechanism-guided framework integrating progression pattern, tissue and plasma reassessment, biomarker-specific interpretation, matched therapy, and longitudinal molecular monitoring. Prospective studies are needed to standardize testing thresholds and optimize treatment sequencing.

Keywords: Non-small cell lung cancer; Targeted therapy; Epidermal growth factor receptor; Mesenchymal–epithelial transition factor; Trophoblast cell-surface antigen 2; Resistance mechanisms

[†]These authors contributed equally to this work.

*Corresponding authors:

Xiangquan Song
(79219121@qq.com);
Xiaoping Li
(li-xp@zjsru.edu.cn)

Citation: Yan K, Tian K, Du L, *et al.* Targeted therapy resistance in EGFR-mutant non-small cell lung cancer: Roles of MET, Trop-2-directed antibody–drug conjugates, and molecular monitoring. *Eurasian J Med Oncol*. doi: 10.36922/EJMO026260304

Received: June 24, 2026

Revised: August 16, 2026

Accepted: September 8, 2026

Published online: September 23, 2026

Copyright: © 2026 Author(s). This is an Open-Access article distributed under the terms of the Creative Commons Attribution License, permitting distribution, and reproduction in any medium, provided the original work is properly cited.

Publisher's Note: AccScience Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

1. Introduction

Non-small cell lung cancer (NSCLC) accounts for approximately 80–85% of all lung cancer cases and remains one of the leading causes of cancer-related mortality worldwide.¹ Although surgery, radiotherapy, and platinum-based chemotherapy remain important treatment modalities, many patients are still diagnosed at an advanced stage or eventually experience disease recurrence, resulting in unsatisfactory long-term outcomes.² This clinical reality has prompted a gradual shift in research focus from traditional treatment paradigms to the in-depth characterization of tumor molecular biology. The evolution of cancer therapy from conventional chemotherapy toward precision oncology has been extensively reviewed elsewhere.³

With the rapid development of high-throughput sequencing and molecular pathology technologies, the treatment paradigm for NSCLC has progressively shifted from traditional histological subtyping to an era of precision medicine centered on driver genes.⁴ Among these molecular events, epidermal growth factor receptor (EGFR) mutations are among the most established actionable drivers in NSCLC, whereas mesenchymal–epithelial transition factor (MET) amplification may function either as a primary oncogenic alteration or as an important bypass mechanism of acquired resistance to EGFR tyrosine kinase inhibitors (TKIs). In contrast, trophoblast cell-surface antigen 2 (Trop-2) is not a classical driver gene but is increasingly recognized as a therapeutically exploitable cell-surface antigen for antibody–drug conjugates (ADCs) in advanced or treatment-resistant NSCLC.^{5–7} Small-molecule TKIs and ADCs directed against these key targets have significantly improved the progression-free survival and overall survival of patients with specific molecular subtypes, thereby establishing the core role of targeted therapy in the comprehensive treatment landscape of NSCLC.^{8,9}

Although targeted therapies provide substantial initial clinical benefit, acquired resistance remains a major barrier to durable disease control. Resistance evolution is characterized by marked spatial and temporal heterogeneity and may involve on-target secondary alterations, activation of bypass signaling pathways, histologic or phenotypic transformation, and tumor microenvironmental remodeling. Common resistance mechanisms following EGFR-TKI therapy include secondary *EGFR* mutations (such as T790M, C797S), activation of bypass signaling pathways (such as *MET* amplification), and alterations in tumor biological behavior (transformation to small cell lung cancer).¹⁰ Furthermore, patients with certain rare mutation subtypes or coexisting multiple molecular aberrations still

lack highly effective and standardized treatment strategies, which remains a challenge in current clinical practice. In recent years, next-generation EGFR-TKIs have expanded into adjuvant therapy and the treatment of uncommon mutations. In parallel, MET-targeted therapies have shown promising activity in *MET* amplification-driven disease, and Trop-2-directed ADCs have emerged as potential options in TKI-resistant populations. Together, these advances have broadened the therapeutic landscape of NSCLC.^{11–13}

Advances in precision medicine are reshaping treatment decisions for NSCLC. The dynamic monitoring of circulating tumor DNA (ctDNA) and minimal residual disease (MRD) assessments has made it possible to identify emerging resistance and molecular relapse earlier. These approaches may also support treatment adjustments based on real-time molecular changes.^{14,15} Against this backdrop, seamlessly integrating molecular monitoring results with targeted therapy strategies to achieve truly individualized treatment has emerged as a crucial direction for current research. To provide a comprehensive synthesis, relevant literature was primarily retrieved from PubMed and Web of Science, focusing on high-quality clinical trials and peer-reviewed translational studies published predominantly within the last five years. Based on these premises, this narrative review summarizes recent advances in EGFR-, MET-, and Trop-2-directed therapies in NSCLC, with particular emphasis on resistance evolution, mechanism-based post-resistance treatment strategies, and the potential integration of ctDNA- and MRD-based molecular monitoring into individualized therapeutic decision-making.^{5,6,11,13} Our objective is to develop a conceptual framework that integrates the diverse pathways involved in the evolutionary trajectory of NSCLC resistance. This framework aims to synthesize key clinical evidence, elucidate primary resistance mechanisms, and assess the potential utility of molecular monitoring, thereby serving as a crucial reference for optimizing individualized treatment strategies following resistance. In contrast to previous reviews that have examined EGFR, MET, or Trop-2 as discrete therapeutic targets, this review seeks to incorporate these targets within a resistance-evolution framework. This framework connects driver inhibition, suppression of bypass resistance, ADC-based post-resistance therapy, and dynamic molecular monitoring.

In clinical practice, therapeutic resistance in NSCLC rarely follows a linear or single-mechanism pattern. Instead, resistance frequently emerges through the coexistence and dynamic competition of multiple subclones. For example, an EGFR-mutant tumor initially dependent on EGFR signaling may later acquire secondary EGFR alterations,

activate MET-mediated bypass signaling, undergo phenotypic transformation, or lose dependence on a single actionable driver.¹⁶ Under such circumstances, post-progression treatment decisions cannot rely only on the molecular status at baseline. Rather, they require repeated reassessment of the evolving tumor genome and biological phenotype.

The clinical significance of integrating EGFR, MET, Trop-2, and molecular monitoring lies in establishing a dynamic treatment decision-making framework for acquired resistance to targeted therapies in NSCLC. Within this framework, EGFR inhibition represents the foundation of targeted treatment for driver-positive NSCLC; MET inhibition addresses one of the most clinically relevant bypass resistance mechanisms; Trop-2-directed ADCs provide therapeutic options for patients with heterogeneous or non-actionable resistance; and ctDNA/MRD monitoring enables dynamic detection of molecular recurrence and emerging resistance before overt clinical progression.¹⁷ To inform this framework, relevant randomized controlled trials, translational studies, and consensus recommendations were identified through PubMed and Web of Science, emphasizing high-quality recent evidence and landmark studies that have defined current understanding of resistance mechanisms and treatment strategies. This paper aims to summarize the evolving landscape of targeted therapy resistance and discuss emerging precision approaches for optimizing sequential therapeutic strategies in NSCLC.^{18,19}

2. Latest research progress in targeted therapy for core targets

2.1. EGFR mutation target: Clinical breakthroughs in adjuvant therapy and rare mutations

EGFR mutations are the most common driver gene alterations in NSCLC, among which exon 19 deletions (19del) and the exon 21 L858R point mutation account for approximately 80% of all EGFR mutations.²⁰ With the continuous development of third-generation EGFR-TKIs, their clinical value across different treatment stages of EGFR-mutant NSCLC has been gradually validated.¹¹

The therapeutic role of EGFR-TKIs has evolved from the management of advanced metastatic disease to a broader disease-course strategy covering early-stage postoperative recurrence prevention, first-line systemic therapy, central nervous system disease control, and subsequent treatment after acquired resistance.^{11,21} This expansion reflects not only the improved potency and selectivity of newer-generation EGFR-TKIs but also the increasing recognition that EGFR-mutant NSCLC should be managed as a molecularly defined disease throughout its entire clinical

trajectory. In early-stage disease, the major objective of EGFR-TKI therapy is to reduce micrometastatic relapse after complete resection. In advanced disease, the treatment goal is prolonged systemic and intracranial disease control while maintaining quality of life. In uncommon EGFR-mutant disease, therapeutic selection is more complex because sensitivity to EGFR inhibition varies substantially according to the exact mutation subtype and molecular conformation.

From this perspective, EGFR mutation status should not be interpreted simply as positive or negative. Classical sensitizing mutations, major uncommon mutations, exon 20 insertions, compound mutations, and resistance-associated secondary mutations represent biologically and clinically distinct entities. This distinction is particularly important when evaluating clinical trial results, because the efficacy observed in classical EGFR-mutant populations cannot be directly extrapolated to all rare mutation subgroups.²²

In the adjuvant setting, a multicenter, randomized, controlled phase III trial demonstrated that the third-generation EGFR-TKI aumolertinib significantly improved disease-free survival in patients with resected EGFR-mutant NSCLC harboring EGFR-sensitizing mutations.²³ Notably, among patients undergoing MRD testing, molecular clearance after adjuvant EGFR-TKI therapy may provide additional prognostic information. However, whether MRD status should directly guide treatment escalation, de-escalation, or treatment duration remains to be validated in prospective studies.²⁴ The integration of MRD assessment into adjuvant EGFR-TKI therapy is clinically attractive but remains methodologically challenging. A persistently positive MRD result after surgery may indicate the presence of residual microscopic disease and a higher probability of subsequent relapse, whereas MRD clearance during adjuvant therapy may reflect effective suppression of residual tumor clones.²⁵ Nevertheless, several unresolved questions constrain the prompt clinical application of these findings. The optimal management strategy for MRD-positive patients—whether it involves extended EGFR-TKI therapy, enhanced systemic treatment, or increased surveillance—remains indeterminate. Conversely, the safety of treatment de-escalation in MRD-negative patients, without adversely affecting long-term survival outcomes, has yet to be ascertained. Furthermore, variations in assay sensitivity, timing of sample collection, the use of tumor-informed versus tumor-naïve platforms, and the inherent biological heterogeneity of ctDNA shedding may affect the accuracy and reliability of MRD interpretation.²⁶ Therefore, MRD should currently be regarded as a promising risk-stratification tool rather than a definitive

basis for modifying adjuvant EGFR-TKI treatment outside prospective clinical trials.⁴

For uncommon *EGFR* mutations, therapeutic sensitivity varies substantially across mutation subtypes. Major uncommon mutations, such as G719X, L861Q, and S768I, may retain sensitivity to selected EGFR-TKIs, whereas *EGFR* exon 20 insertion mutations generally show limited sensitivity to conventional first- to third-generation EGFR-TKIs. Therefore, treatment strategies should be discussed according to specific mutation categories rather than grouped under a broad definition of “rare *EGFR* mutations.”²⁷ In patients with locally advanced or metastatic NSCLC harboring classical EGFR-sensitizing mutations, limetinib demonstrated prolonged progression-free survival compared with gefitinib in the first-line setting. Its reported intracranial activity also suggests potential value for patients with concurrent brain metastases.^{28,29} Overall, these findings support the expanding role of third-generation EGFR-TKIs across selected clinical settings, including adjuvant therapy, first-line treatment for classical EGFR-sensitizing mutations, and carefully defined uncommon mutation subgroups.

2.2. *MET* amplification target: Clinical progress of selective inhibitors

MET amplification may act as a primary oncogenic driver in a subset of NSCLC and is also a well-recognized bypass mechanism of acquired resistance to EGFR-TKIs. However, its clinical interpretation remains challenging because detection platforms and positivity thresholds differ across studies, including fluorescence *in situ* hybridization (FISH), next-generation sequencing (NGS)-based copy-number assessment, and *MET*/centrosomal protein 7 (CEP7) ratio-based criteria.³⁰

Unlike point mutations, *MET* amplification is a quantitative and context-dependent biomarker.³⁰ Its predictive value may be influenced by absolute gene copy number, *MET*/CEP7 ratio, tumor purity, focality of amplification, and the presence of chromosome 7 polysomy.³¹ High-level focal *MET* amplification is more likely to represent a true oncogenic driver or dominant resistance mechanism, whereas low-level copy-number gain may reflect genomic instability or coexist with other resistance alterations.³² This distinction is clinically relevant because patients with high-level *MET* amplification may derive greater benefit from *MET*-directed therapy, while those with low-level or heterogeneous amplification may show less durable responses.

Another important issue is whether *MET* amplification occurs as a *de novo* driver alteration or as an acquired bypass mechanism after EGFR-TKI exposure. In the former

situation, selective *MET* inhibitor monotherapy may be a rational treatment approach in biomarker-selected patients. In contrast, when *MET* amplification emerges after EGFR-TKI therapy in EGFR-mutant NSCLC, continued EGFR pathway suppression combined with *MET* inhibition may be biologically more appropriate, because the original EGFR-dependent clone and the *MET*-amplified resistant clone may coexist. Therefore, the clinical interpretation of *MET* amplification should integrate treatment history, coexisting genomic alterations, amplification level, and the spatial-temporal pattern of disease progression.

Recently, the selective *MET* inhibitor vebreltinib demonstrated antitumor activity in a phase II clinical study of *MET* amplification-driven advanced NSCLC.³³ These findings underscore the therapeutic significance of *MET* inhibition in patients selected based on specific biomarkers. However, the clinical positioning of *MET* inhibitor monotherapy relative to EGFR/*MET* dual inhibition should be evaluated based on whether *MET* amplification constitutes a primary driver event or an acquired mechanism of bypass resistance.

In addition, molecular monitoring technologies for efficacy evaluation and prognostic assessment are attracting increasing attention. In *MET*-amplified disease, molecular monitoring may help clarify whether *MET* amplification represents a dominant resistance clone or only one component of a heterogeneous resistance landscape. Previous studies have indicated that low baseline ctDNA levels, early ctDNA clearance during treatment, and the absence of co-mutations are associated with longer progression-free survival.^{34,35} These findings suggest that dynamic molecular monitoring holds promise as a crucial tool for predicting efficacy and optimizing treatment in patients with *MET* amplification. Importantly, *MET* amplification should not be interpreted in isolation. Coexisting EGFR secondary mutations, human epidermal growth factor receptor 2 (HER2) amplification, cell-cycle pathway alterations, or histologic transformation may reduce the efficacy of *MET*-directed therapy, highlighting the need for comprehensive molecular profiling at progression.

2.3. Trophoblast cell-surface antigen 2 target: Therapeutic value of antibody–drug conjugates after drug resistance

Trophoblast cell-surface antigen 2 is expressed in a substantial proportion of NSCLC and has been associated with aggressive tumor biology and treatment resistance in selected contexts. However, its role as a predictive biomarker for Trop-2-directed ADCs remains incompletely established.³⁶ Trop-2-directed ADCs consist

of a monoclonal antibody against Trop-2, a chemical linker, and a cytotoxic payload.

Unlike EGFR and MET, Trop-2 does not function primarily as a classical oncogenic driver in NSCLC. Its therapeutic relevance stems mainly from its cell-surface expression and suitability as an antigen for targeted cytotoxic drug delivery. This distinction is important because Trop-2-directed ADCs are not designed to block a single dominant signaling pathway. Instead, they combine antigen recognition with intracellular release of a cytotoxic payload, thereby producing antitumor activity even in tumors with heterogeneous resistance mechanisms. The bystander effect may further enhance activity in tumors with uneven antigen expression, although excessive payload diffusion may also contribute to off-target toxicity. Therefore, Trop-2-directed ADCs occupy a unique therapeutic position between targeted therapy and chemotherapy.

After binding to Trop-2-expressing tumor cells, these agents deliver cytotoxic drugs intracellularly and may exert bystander killing effects, thereby providing a therapeutic strategy for complex resistance contexts that is less dependent on a single oncogenic driver pathway.¹³

OptiTROP-Lung04 is a multicenter randomized controlled trial that enrolled 376 patients with advanced NSCLC harboring EGFR-sensitizing mutations who experienced disease progression following EGFR-TKI therapy. The study compared the efficacy of sacituzumab tirumotecan monotherapy against pemetrexed combined with platinum-based chemotherapy. The results demonstrated that sacituzumab tirumotecan significantly prolonged the median progression-free survival of patients (8.3 months vs. 4.3 months, hazard ratio = 0.49), and also showed a distinct advantage in overall survival (hazard ratio = 0.60, $p = 0.001$).³⁷ Regarding efficacy indicators, the ORR in the sacituzumab tirumotecan group reached 60.6%, which was significantly higher than the 43.1% observed in the chemotherapy group, and the duration of response was also substantially extended.³⁸

Sacituzumab tirumotecan demonstrated consistent efficacy across subgroups, including those previously treated with multiple EGFR-TKIs and those with brain or liver metastases.³⁷ Concurrently, its incidence of grade 3 or higher hematological adverse events was lower than that of the chemotherapy group, leading to improved patient quality of life. Nevertheless, the toxicity profiles of Trop-2-directed ADCs may vary according to payload type, linker stability, and drug-to-antibody ratio.³⁹ Clinical application therefore requires careful monitoring for hematologic toxicity, stomatitis, gastrointestinal adverse events, and interstitial lung disease, particularly in heavily pretreated

patients or those with baseline pulmonary impairment. The aforementioned results indicate that, provided these toxicities are meticulously managed, Trop-2-targeted ADCs represent a valuable therapeutic option in patients who have failed EGFR-TKI therapy, offering a new alternative for post-resistance treatment strategies (Table 1).³⁷ The findings indicate that sacituzumab tirumotecan could serve as a clinically significant later-line treatment option for EGFR-mutant NSCLC following the failure of EGFR-TKI therapy, especially in cases where no clearly actionable resistance alterations are identified or where multiple resistance mechanisms are present. However, its optimal sequencing in relation to platinum-based chemotherapy, EGFR/MET dual inhibition, and other ADCs has yet to be determined.

3. Resistance mechanisms after EGFR-TKI therapy and mechanism-based treatment strategies

3.1. Molecular and biological mechanisms of targeted therapy resistance

Resistance mechanisms in NSCLC targeted therapy are highly heterogeneous and dynamically evolving. Resistance may be present before treatment as primary resistance or may emerge under therapeutic selective pressure as acquired resistance. These two forms are not mutually exclusive and may coexist within spatially heterogeneous tumor subclones.^{40,41} From a molecular biology perspective, currently identified resistance mechanisms can be broadly classified into four categories: on-target resistance, such as secondary *EGFR* mutations; off-target or bypass resistance, such as MET or HER2 amplification; lineage plasticity and histologic transformation, including transformation to small-cell lung cancer (SCLC) or squamous cell carcinoma; and non-genetic mechanisms, including epithelial-mesenchymal transition and tumor microenvironmental remodeling.^{5,42,43} In EGFR-TKI therapy, secondary mutations in the *EGFR* gene remain the most classical mechanism of acquired resistance. Among them, the T790M mutation was once the primary cause of resistance to first- and second-generation EGFR-TKIs. However, with the widespread application of third-generation EGFR-TKIs, the C797S mutation has gradually emerged as a new core resistance event.^{16,44} The allelic configuration of C797S is clinically relevant. When C797S and T790M occur in trans, combined first- and third-generation EGFR-TKI therapy may retain activity in selected cases. In contrast, when these mutations occur in cis, this strategy is generally ineffective, highlighting the importance of allelic-level molecular characterization.^{45,46} Beyond alterations in the target itself, the activation of

Table 1. Key molecular targets and representative precision therapeutic strategies in NSCLC

Molecular target	Alteration	Representative therapy	Indicated population	Major clinical implication	References
EGFR	Exon 19 deletion/L858R	Aumolertinib	Patients with resected EGFR-mutant NSCLC in the adjuvant setting	Phase 3 evidence supports improved disease-free survival in resected EGFR-mutant NSCLC	23
EGFR	Major uncommon mutations (e.g., G719X, L861Q, S768I)	Aumolertinib	Patients with advanced NSCLC harboring uncommon <i>EGFR</i> mutations	Preclinical and early clinical evidence suggest antitumor activity in selected uncommon EGFR-mutant NSCLC	27
EGFR	EGFR-sensitising mutations	Limertinib	Patients with locally advanced or metastatic EGFR-mutant NSCLC receiving first-line treatment	Phase 3 evidence supports prolonged progression-free survival versus gefitinib in the first-line setting.	28
MET	Amplification	Vebreltinib	Patients with advanced <i>MET</i> amplification-driven NSCLC	Phase II data support antitumor activity in biomarker-selected patients, although clinical positioning continues to evolve	33
Trop-2	Trop-2 overexpression	Trop-2-directed ADCs	Previously treated patients with advanced NSCLC	Trop-2-directed ADCs have shown clinical activity in selected populations; however, the predictive value of Trop-2 expression remains to be fully established.	7,39
Trop-2	EGFR-TKI-resistant EGFR-mutant disease	Sacituzumab tirumotecan (sac-TMT)	Patients with locally advanced or metastatic EGFR-mutant non-squamous NSCLC after EGFR-TKI failure	Randomized phase III evidence supports sac-TMT as an effective later-line option, with improved progression-free survival and overall survival versus platinum-pemetrexed chemotherapy.	37

Notes: The therapies listed are representative examples and do not imply class-wide efficacy or uniform regulatory approval across jurisdictions; *MET* amplification, *MET* exon 14 skipping, and *MET* protein overexpression are biologically and clinically distinct biomarkers and should not be used interchangeably; Trop-2 expression is not currently a universally validated predictive biomarker for selecting Trop-2-directed antibody–drug conjugates; clinical evidence and regulatory status should be interpreted according to the specific agent, biomarker definition, study population, treatment setting, and jurisdiction.

Abbreviations: ADC: Antibody–drug conjugate; EGFR: Epidermal growth factor receptor; MET: Mesenchymal–epithelial transition factor; NSCLC: Non-small cell lung cancer; Sac-TMT: Sacituzumab tirumotecan; TKI: Tyrosine kinase inhibitor; Trop-2: Trophoblast cell-surface antigen 2.

bypass signaling pathways also plays a critical role in the resistance process. *MET* amplification is among the most clinically relevant bypass mechanisms after EGFR-TKI therapy. By activating downstream phosphoinositide 3-kinase/protein kinase B, mitogen-activated protein kinase, and related survival pathways independently of EGFR signaling, *MET* amplification allows tumor cells to maintain proliferative and anti-apoptotic signaling despite persistent EGFR inhibition.^{45,47,48} Additionally, molecular events such as HER2 amplification, *BRAF* mutations, and *PIK3CA* mutations can also mediate similar resistance processes in a subset of patients. Furthermore, oncogenic fusions (e.g., *ALK*, *RET*) and cell cycle gene alterations (e.g., *CDK4/CDK6* amplification) have emerged as highly

relevant off-target bypass mechanisms driving third-generation TKI resistance, necessitating comprehensive NGS upon progression.⁴⁹

At a higher level of tumor biological alterations, phenotypic transformation has increasingly garnered attention. A proportion of EGFR-mutant NSCLC can transform into SCLC or squamous cell carcinoma phenotypes following TKI therapy. This resistance pattern is clinically challenging because it is often associated with rapid disease progression, reduced dependence on the original EGFR signaling pathway, and limited responsiveness to subsequent targeted therapy. Therefore, tissue re-biopsy should be considered when clinical progression is inconsistent with the expected pattern

of molecular resistance.⁵⁰⁻⁵² Furthermore, epithelial-mesenchymal transition is also believed to be closely associated with the resistance phenotype and enhanced invasiveness (Figure 1).^{53,54}

3.2. Mechanism-based therapeutic strategies for resistance

A mechanism-based framework has gradually emerged for different resistance patterns. For resistance mediated by *EGFR* C797S and related on-target alterations, fourth-

generation *EGFR*-TKIs are being developed to overcome resistance while preserving activity against activating *EGFR* mutations. However, most agents remain investigational, and their efficacy, safety, optimal sequencing, and activity against compound *EGFR* mutations require further clinical validation.⁵⁵ Although related drugs are still in clinical research, preliminary data suggest this strategy has potential in specific molecular contexts. In addition, for patients with C797S and T790M in a *trans* configuration, the strategy of combining first- and third-generation




Figure 1. Mechanism-guided assessment, treatment, and longitudinal monitoring of acquired EGFR-TKI resistance in EGFR-mutant NSCLC. This figure illustrates an integrated framework connecting acquired resistance mechanisms, molecular and histologic reassessment, mechanism-guided treatment, and longitudinal monitoring after EGFR-TKI progression. Resistance mechanisms are classified into four categories: on-target *EGFR* alterations; off-target or bypass pathway activation, including acquired *MET* amplification and *HER2* (*ERBB2*) amplification; lineage plasticity and histologic transformation; and non-genetic adaptive mechanisms, including epithelial-mesenchymal transition, drug-tolerant persister states, and tumor-microenvironmental remodeling. At progression, plasma ctDNA testing and tissue biopsy provide complementary information. Tissue biopsy is required when plasma findings are negative or discordant and whenever histologic transformation is suspected. Treatment is selected according to the dominant resistance mechanism and the strength of supporting evidence. Options include investigational next-generation *EGFR* inhibitors for selected on-target alterations, *EGFR*/*MET* dual inhibition for confirmed acquired *MET* amplification, histology-directed treatment for transformation, and chemotherapy, drug-specific ADCs, or clinical trials when no actionable mechanism is identified. Serial ctDNA monitoring should be integrated with radiographic and clinical assessment to evaluate clonal evolution.

Abbreviations: ADCs: Antibody-drug conjugates; ctDNA: Circulating tumor DNA; *EGFR*: Epidermal growth factor receptor; *HER2*: Human epidermal growth factor receptor 2; *MET*: Mesenchymal-epithelial transition factor; NSCLC: Non-small cell lung cancer; TKI: Tyrosine kinase inhibitor. This figure was drawn using Figdraw (Export ID: IUTRO42cab; Version 2.0, Home for Researchers, China).

EGFR-TKIs has demonstrated certain efficacy in some studies, although its applicable population and long-term benefits require further validation.^{56,57}

In bypass activation-related resistance, target combination inhibition strategies have become a major focus of current research. In resistance mechanisms driven by MET amplification, combining MET inhibitors with EGFR-TKIs has shown synergistic antitumor activity in multiple clinical studies, significantly improving disease control and progression-free survival.⁵⁸ This strategy provides a clear molecular rationale for resolving acquired resistance to EGFR-TKIs, and also underscores the importance of dynamic molecular testing before and during treatment.⁵⁹

For patients lacking clearly targetable molecular aberrations or those with multiple resistance mechanisms, ADCs offer a relatively “mechanism-independent” therapeutic approach.³⁷ ADC drugs targeting Trop-2, via antigen-mediated intracellular delivery of cytotoxic agents, have shown clinically meaningful activity in patients following EGFR-TKI progression. This effect is not significantly influenced by specific *EGFR* mutation types or bypass activation status, thereby providing a novel option for treatment in complex resistance contexts.³⁷

Regarding resistance related to phenotypic transformation, treatment strategies historically borrowed from standard chemotherapy regimens for SCLC. However, patients typically exhibit limited and transient responses to traditional platinum-etoposide and immunotherapy. Recently, a paradigm shift has occurred with the development of delta-like canonical Notch ligand 3 (DLL3)-targeted bispecific T-cell engagers

(e.g., tarlatamab), which have demonstrated remarkable and durable clinical activity in previously treated *de novo* SCLC.⁶⁰ However, applying these DLL3-directed therapies in EGFR-mutant SCLC transformation remains complex. Recent real-world multi-institutional data suggest that clinical responses to tarlatamab in transformed SCLC may be substantially lower than those observed in *de novo* SCLC, highlighting the distinct underlying biology and intense treatment refractoriness of the transformation phenotype.⁶¹ Future studies should further elucidate the molecular drivers of phenotypic transformation to identify additional novel therapeutic targets (Table 2). Clinically, treatment after EGFR-TKI progression should follow a mechanism-guided algorithm. Tissue re-biopsy and/or plasma-based NGS should first be performed to identify actionable resistance alterations. Patients with *MET* amplification may be considered for EGFR/MET dual inhibition; those with *EGFR* C797S or compound *EGFR* mutations may be eligible for investigational next-generation EGFR inhibitors; patients with histologic transformation require pathology-confirmed treatment adaptation; and those without actionable alterations may be considered for chemotherapy, ADCs, or clinical trials.

3.3. The role of precision monitoring in resistance management

With the widespread application of targeted therapies in NSCLC, resistance management has gradually shifted from relying solely on radiographic and clinical symptom evaluations to a precision monitoring paradigm that integrates molecular-level information. Conventional radiographic assessment remains indispensable for evaluating tumor burden, but it may not capture early

Table 2. Mechanism-guided interpretation and therapeutic options after EGFR-TKI progression

Resistance category	Specific alteration or phenotype	Required confirmation and clinical setting	Representative management strategy	Evidence status	Key clinical implication and limitation	References
On-target <i>EGFR</i> alteration	Acquired <i>EGFR</i> T790M	Tissue- or plasma-confirmed T790M after progression on a first- or second-generation EGFR-TKI; tissue testing should be considered when plasma is negative, and the result would alter treatment	Osimertinib	Established/ approved in this biomarker-defined setting	Randomized phase III evidence supports osimertinib over platinum–pemetrexed in T790M-positive disease after prior first-line EGFR-TKI therapy. This evidence should not be directly extrapolated to resistance after first-line osimertinib.	49,62

(Cont'd...)

Table 2. (Continued)

Resistance category	Specific alteration or phenotype	Required confirmation and clinical setting	Representative management strategy	Evidence status	Key clinical implication and limitation	References
On-target <i>EGFR</i> alteration	C797S or complex/compound <i>EGFR</i> alterations	Comprehensive NGS; allelic phasing should be considered when C797S and T790M coexist, and the cis/trans configuration would affect treatment selection	Investigational fourth-generation <i>EGFR</i> inhibitors or genotype-selected clinical trials	Investigational; phase I/II or early clinical evidence	Fourth-generation <i>EGFR</i> inhibitors remain under clinical investigation. Combined first- and third-generation <i>EGFR</i> inhibition may be considered only in highly selected trans-configured cases and is not a standard treatment.	63,64
Off-target or bypass pathway activation	Acquired <i>MET</i> amplification	Report assay platform, absolute gene copy number, <i>MET</i> /CEP7 ratio when available, amplification focality, and study-specific threshold; confirm plasma-only copy-number findings when feasible	Continued <i>EGFR</i> inhibition combined with a selective <i>MET</i> inhibitor	Phase II study-supported; investigational or jurisdiction-specific	<i>EGFR</i> / <i>MET</i> dual inhibition has demonstrated clinical activity in study-defined acquired <i>MET</i> -mediated resistance. <i>MET</i> amplification must be distinguished from <i>MET</i> exon 14 skipping, <i>MET</i> protein overexpression, chromosome 7 polysomy, and nonspecific low-level copy-number gain.	65
Off-target or bypass pathway activation	<i>HER2</i> amplification; acquired <i>ALK</i> or <i>RET</i> fusion; selected <i>BRAF</i> , <i>PIK3CA</i> , or cell-cycle pathway alterations	Comprehensive genomic profiling and alteration-specific confirmation when the result would change treatment	Alteration-matched therapy or biomarker-selected clinical trial	Early clinical or investigational evidence	These alterations are relatively uncommon and may be subclonal. No universal “ <i>EGFR</i> – <i>HER2</i> dual inhibition” standard has been established for acquired <i>HER2</i> -amplified resistance. Treatment should be based on alteration-specific evidence.	66
Lineage plasticity and histologic transformation	Transformation to small-cell lung cancer or squamous cell carcinoma	Tissue re-biopsy and pathological confirmation are required	Histology-directed treatment; platinum–etoposide is commonly used for SCLC transformation; DLL3-directed therapy should preferably be evaluated in an appropriate clinical trial	Histology-directed chemotherapy: established practice; DLL3-directed therapy in transformed disease: early/retrospective evidence	Plasma genomic testing cannot confirm histologic transformation. Evidence for tarlatamab in <i>de novo</i> SCLC should not be directly extrapolated to <i>EGFR</i> -mutant transformed SCLC.	60,61,67

(Cont'd...)

Table 2. (Continued)

Resistance category	Specific alteration or phenotype	Required confirmation and clinical setting	Representative management strategy	Evidence status	Key clinical implication and limitation	References
Non-genetic adaptive resistance	Epithelial–mesenchymal transition, drug-tolerant persister states, transcriptional reprogramming, or microenvironmental remodeling	Pathological or translational assessment; no standardized clinical diagnostic threshold	Platinum-based chemotherapy or clinical-trial enrollment	Preclinical or early translational evidence; no validated mechanism-specific standard	EMT is a phenotypic resistance mechanism rather than a molecular target. “Targeting EMT pathways” should not be presented as an established clinical treatment.	53,68
No dominant actionable mechanism or multiclonal resistance	No sufficiently validated actionable alteration, or coexistence of several competing resistance mechanisms	Integrate progression pattern, tissue and plasma genomic results, and histology when indicated	Platinum-based chemotherapy, a drug-specific ADC supported in the relevant setting, or clinical-trial enrollment	Treatment-specific evidence; phase III evidence exists for selected ADCs, but class-wide conclusions are inappropriate	Trop-2 expression is not a universally validated predictive biomarker. Trop-2-directed ADC selection should be based on drug-specific trial evidence, prior treatment, toxicity, and regulatory availability rather than “high Trop-2 expression” alone.	39

Notes: “Established,” “study-supported,” “investigational,” and “early evidence” refer to the strength and clinical context of the available evidence and do not imply identical regulatory approval across jurisdictions; regulatory status should be verified according to the region and manuscript submission date; *MET* amplification, *MET* exon 14 skipping, and *MET* protein overexpression are biologically and clinically distinct and should not be used interchangeably; Trop-2 expression is not currently a universally validated predictive biomarker for selecting Trop-2-directed ADC therapy.

Abbreviations: ADC: Antibody–drug conjugate; DLL3: Delta-like canonical Notch ligand 3; EGFR: Epidermal growth factor receptor; EMT: Epithelial–mesenchymal transition factor; HER2: Human epidermal growth factor receptor 2; MET: Mesenchymal–epithelial transition factor; NGS: Next-generation sequencing; NSCLC: Non-small cell lung cancer; SCLC: Small-cell lung cancer; TKI: Tyrosine kinase inhibitor; Trop-2: Trophoblast cell-surface antigen 2.

molecular evolution during targeted therapy. Resistant subclones can emerge before measurable radiographic progression, creating a temporal gap between molecular relapse and imaging-defined disease progression.⁶⁹ Therefore, monitoring tools capable of dynamically reflecting changes in the molecular characteristics of tumors have become key instruments for optimizing resistance management and treatment decision-making.

Nevertheless, ctDNA findings should always be interpreted in clinical context. A positive plasma result may help rapidly identify actionable resistance mechanisms, but a negative result does not exclude molecular progression. Tumors with low shedding, isolated brain metastases, pleural-only disease, or small-volume intrathoracic recurrence may release insufficient ctDNA into the bloodstream. In such cases, tissue re-biopsy remains important, especially when histologic transformation is

suspected. In addition, clonal hematopoiesis may generate false-positive variants unrelated to the tumor, particularly in genes such as *TP53* or other hematologic-associated alterations. Therefore, ctDNA should complement rather than fully replace tissue-based molecular and pathological assessment. However, false-negative results may occur in patients with low tumor DNA shedding, isolated central nervous system progression, or limited intrathoracic disease.¹⁷ Therefore, negative plasma results should not replace tissue biopsy when clinical suspicion of progression remains high.⁷⁰ In early-stage and locally advanced NSCLC, MRD assessment is emerging as a promising tool for relapse risk stratification after curative-intent therapy. Nevertheless, standardized assays, optimal sampling intervals, clinically meaningful thresholds, and MRD-guided treatment strategies remain under active investigation (Figure 2).⁷¹




Figure 2. Integrated framework of core therapeutic targets, resistance mechanisms, and molecular monitoring in non-small cell lung cancer. This figure delineates the complex landscape of precision therapy for NSCLC, structured around three principal therapeutic targets—EGFR, MET, and Trop-2—and their associated treatment modalities, predominant resistance mechanisms, and the significance of molecular monitoring in tailoring individualized treatment strategies. The figure is organized as follows: The upper panel highlights the three primary therapeutic targets (EGFR, MET, and Trop-2) that are currently actionable in clinical settings for NSCLC. Below each target, the respective therapeutic strategies are depicted: EGFR-targeted therapy employing EGFR TKIs, MET-targeted therapy utilizing MET inhibitors, and Trop-2-directed ADCs. The central section provides an overview of the major resistance mechanisms that may develop during or post-treatment. For EGFR-TKIs, resistance can manifest through secondary *EGFR* mutations (e.g., T790M, C797S), activation of bypass pathways (e.g., *MET* amplification), and phenotypic alterations (e.g., epithelial–mesenchymal transition or transformation to small-cell lung cancer). In the context of MET-targeted therapy, resistance may occur via bypass activation through the PI3K/AKT and MAPK pathways. Abbreviations: ADC: Antibody–drug conjugate; AKT: Protein kinase B; ctDNA: Circulating tumor DNA; EGFR: Epidermal growth factor receptor; EMT: Epithelial–mesenchymal transition factor; MAPK: Mitogen-activated protein kinase; MET: Mesenchymal–epithelial transition factor; MRD: Minimal residual disease; NGS: Next-generation sequencing; NSCLC: Non-small cell lung cancer; PI3K: Phosphoinositide 3-kinase; SCLC: Small-cell lung cancer; TKI: Tyrosine kinase inhibitor; Trop-2: Trophoblast cell-surface antigen 2.

4. Discussion

The major challenge in contemporary NSCLC targeted therapy is no longer limited to identifying actionable drivers at diagnosis, but increasingly lies in managing molecular evolution under therapeutic pressure. EGFR-, MET-, and Trop-2-directed strategies represent different but interconnected dimensions of this process: driver inhibition, bypass resistance suppression, and post-resistance cytotoxic delivery.⁷² Future research must

continue to deepen across multiple dimensions—including the development of novel targets, the management of resistance mechanisms, the application of precision biomarkers, and the optimization of treatment for special populations—to further propel the targeted therapy of NSCLC toward a more precise and individualized direction.⁷³

From a clinical decision-making perspective, treatment after EGFR-TKI progression should begin with

classifying the progression pattern. Patients with limited oligoprogression may benefit from local therapy while continuing the original EGFR-TKI, especially when most disease sites remain controlled. In contrast, patients with systemic progression require comprehensive reassessment of resistance mechanisms. Plasma ctDNA testing can serve as an initial noninvasive approach, but tissue biopsy should be considered when plasma testing is negative, when transformation is clinically suspected, or when a progressing lesion is accessible.

After molecular reassessment, treatment should be selected according to the dominant resistance pattern. Patients with *MET* amplification may be candidates for EGFR/*MET* dual inhibition. Patients with *EGFR* C797S or complex *EGFR* compound mutations should preferably be considered for investigational next-generation *EGFR* inhibitors or genotype-specific clinical trials.¹⁶ Patients with histologic transformation require pathology-confirmed treatment adaptation, usually shifting away from standard *EGFR*-TKI-based strategies. For patients without actionable resistance alterations, or those with multiple heterogeneous mechanisms, platinum-based chemotherapy, Trop-2-directed ADCs, or clinical trial enrollment may be reasonable options.⁸ This stepwise framework highlights that post-resistance therapy should not be chosen solely according to treatment line, but according to the biological mechanism driving progression.

In addition to well-characterized bypass mechanisms such as *MET* amplification, rare oncogenic fusions, including *ALK* rearrangements, have emerged as clinically significant drivers of acquired resistance to *EGFR*-TKIs in a subset of patients. An illustrative case report documented that an acquired *EML4-ALK* fusion, specifically Variant 3a/b, mediated resistance to osimertinib in a patient with *EGFR* L858R-positive NSCLC, underscoring that even uncommon fusion events can dominate the resistance phenotype under therapeutic pressure. This observation underscores the necessity for comprehensive NGS at the time of disease progression, as the identification of such fusions may redirect treatment toward alternative targeted strategies, such as *ALK* inhibition, in patients who might otherwise be considered to have “non-actionable” resistance. Nevertheless, the true frequency, clonal architecture, and optimal management of *ALK* fusion-mediated resistance remain largely unknown. Prospective molecular profiling and collaborative case registries are essential to determine whether routine screening for these rare events translates into meaningful clinical benefit and to define their potential impact on treatment paradigms.⁷⁴

Despite the introduction of a mechanism-guided framework, its implementation in routine clinical practice

is impeded by several unresolved operational challenges that are not comprehensively addressed in the current literature. Firstly, *MET* amplification, recognized as one of the most actionable bypass mechanisms, lacks a universally accepted assay-specific threshold for patient selection. Various studies have utilized different cutoffs, such as a FISH *MET*/CEP7 ratio of ≥ 2.0 or an NGS-based copy number of ≥ 5 , without direct comparative analysis of these criteria in prospective head-to-head trials. As a result, clinicians face uncertainty when interpreting a “positive” *MET* amplification result, particularly when the reported level falls within an ambiguous range. We propose that, until standardized criteria are established, treatment decisions should be based on thresholds validated in the pivotal trial of the specific *MET* inhibitor being considered (e.g., FISH gene copy number ≥ 10 or NGS-derived copy number ≥ 10 for savolitinib-based combinations), and that low-level amplification should not automatically guide therapy outside of clinical trials. Secondly, the optimal sequencing of post-resistance therapies remains unclear.

First, exploring novel molecular targets and combination treatment strategies remains a core direction for future research. Although relatively mature treatment systems have been established for classic targets such as *EGFR* and *MET*, efficacy remains limited for certain mutation subtypes (*KRAS* G12C, *BRAF* V600E).^{75,76} In addition, the coexistence of multiple resistance mechanisms suggests that single-target inhibition is often insufficient to achieve durable control.⁶ Therefore, combined inhibition strategies centered on key signaling pathways—such as *EGFR*/*MET* dual-target inhibition, or the rational combination of targeted therapy with anti-angiogenic therapy or immunotherapy—hold promise for delaying the onset of resistance through synergistic effects. However, their optimal combination modalities, sequencing of administration, and toxicity management still require further clarification through prospective clinical studies.^{77,78}

Second, biomarker-guided precision treatment decision-making needs further refinement. Dynamic monitoring of ctDNA and MRD assessment provide vital tools for predicting targeted-therapy efficacy and identifying early resistance. Nevertheless, their detection thresholds, timing, and clinical decision-making criteria remain largely unstandardized. More importantly, the routine clinical utility of ctDNA is hindered by biological limitations, including low tumor DNA shedding (particularly in intrathoracic-only progression), interference from clonal hematopoiesis of indeterminate potential leading to false positives, and the inherent spatial heterogeneity of resistance clones that may cause

discordance between tissue and plasma genotyping.^{69,71} Future research should incorporate large-scale prospective cohorts to validate the predictive value of ctDNA clearance, specific mutational profile changes, and Trop-2 expression across different treatment stages. Such efforts may facilitate the transition of molecular monitoring from an adjunctive tool to a basis for therapeutic decision-making.⁷⁹

Furthermore, the further optimization of post-resistance treatment strategies remains an urgent clinical issue to be resolved. ADCs have demonstrated clinically meaningful activity in EGFR-TKI-resistant NSCLC, but several questions remain unresolved. These include whether Trop-2 expression can reliably predict benefit, how ADCs should be sequenced with chemotherapy or targeted combinations, and whether specific resistance mechanisms influence ADC sensitivity. Differences in payload, linker design, bystander effect, and toxicity profiles should also be considered when comparing ADC agents.^{37,39} However, due to differences in target selection, drug payloads, and toxicity profiles among various ADC drugs, their optimal timing of application and sequential or combination modalities with other treatments still require more clinical data for support. Additionally, for specific resistance mechanisms such as phenotypic transformation (transformation to small cell lung cancer), effective targeted approaches are currently lacking. An in-depth elucidation of the relevant molecular mechanisms may provide a breakthrough for developing novel treatment strategies.^{51,67}

Finally, the systematic evaluation of special populations and real-world treatment outcomes also warrants attention. Special populations require more dedicated evidence. Patients with brain metastases need therapies with reliable intracranial activity; patients with liver metastases often have aggressive disease biology and poorer prognosis; and elderly patients or those with comorbidities may have reduced tolerance to combination therapy or ADC-related toxicity. Prospective subgroup analyses and real-world studies are therefore necessary to refine individualized treatment strategies.^{80,81} Future efforts should strengthen real-world studies and prospective subgroup analyses to evaluate the efficacy, safety, and impact on quality of life of targeted drugs in these populations, thereby providing more generalizable evidence for individualized dose adjustments and optimized treatment strategies.

Overall, the future development of targeted therapy for NSCLC will rely on the deep integration of basic research and clinical practice. By continuously advancing the research and development of drugs for novel targets, refining the molecular biomarker system, and optimizing combination and sequential treatment strategies, it is

expected to further break through the bottleneck of drug resistance, maximize therapeutic benefits, and propel the NSCLC management paradigm toward a true transformation into precision medicine.

5. Conclusion

In recent years, with the continuous refinement of molecular subtyping and the constant emergence of innovative drugs, targeted therapy for NSCLC has entered a new phase of precision and multidimensional development. Treatment strategies revolving around key molecular targets such as EGFR, MET, and Trop-2 have not only significantly prolonged survival in specific populations but also progressively formed a more comprehensive therapeutic paradigm across disease stages (adjuvant therapy, advanced-stage therapy, and post-resistance therapy). Third-generation EGFR-TKIs have demonstrated encouraging efficacy in the adjuvant therapy of patients with EGFR-sensitizing mutations and in rare mutation subtypes, providing new evidence-based rationales for early recurrence risk management and treatment of special populations.^{11,12,24}

Meanwhile, as *MET* amplification represents a crucial driver event and resistance mechanism, successful application of its targeted inhibitors has filled previous treatment gaps, marking significant progress in strategies targeting bypass activation resistance.⁶ Furthermore, ADCs targeting Trop-2 have achieved dual benefits in both progression-free survival and overall survival among patients who experienced EGFR-TKI treatment failure, thereby expanding the available options for post-resistance therapies.³⁷

Despite these remarkable advancements, targeted therapy for NSCLC still faces multiple challenges, including complex resistance mechanisms, the coexistence of multiple molecular aberrations, and the current limitations of predictive biomarkers. Future research should focus on developing novel targets, optimizing multi-target combined inhibition strategies, and establishing dynamic monitoring systems based on ctDNA and MRD to support earlier and more precise treatment decisions.^{24,82} Concurrently, the safety and efficacy of treatments in special populations—such as elderly patients and those with brain or liver metastases—urgently require validation through additional prospective studies.

In summary, ongoing advancements in targeted drug development, coupled with the enhanced integration of molecular biomarkers and precision-monitoring technologies, have the potential to significantly enhance the long-term efficacy of targeted therapies in NSCLC. These developments may ultimately improve both survival

rates and quality of life for patients.

Acknowledgments

None.

Funding

This study was financially supported by the Zhejiang Health Information Association Research Program (2025XHZN-Z04). The funder did not participate in the design, performance, or reporting of the current study.

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Author contributions

Conceptualization: Kai Yan, Kewei Tian

Funding acquisition: Xiangquan Song, Xiaoping Li

Visualization: Kai Yan, Kewei Tian

Writing—original draft: Kai Yan, Kewei Tian

Writing—review & editing: Lailing Du, Jinghua Yuan, Xinling Zhang, Xiangquan Song, Xiaoping Li

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data

Not applicable.

References

- Bray F, Laversanne M, Sung H, *et al.* Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2024;74(3):229-263.
doi: 10.3322/caac.21834
- Fox AH, Nishino M, Osarogiagbon RU, *et al.* Acquiring tissue for advanced lung cancer diagnosis and comprehensive biomarker testing: A National Lung Cancer Roundtable best-practice guide. *CA Cancer J Clin.* 2023;73(4):358-375.
doi: 10.3322/caac.21774
- Sonkin D, Thomas A, Teicher BA. Cancer treatments: Past, present, and future. *Cancer Genet.* 2024;286-287:18-24.
doi: 10.1016/j.cancergen.2024.06.002
- Rolfo C, Mack P, Scagliotti GV, *et al.* Liquid Biopsy for Advanced NSCLC: A Consensus Statement From the International Association for the Study of Lung Cancer. *J Thorac Oncol.* 2021;16(10):1647-1662.
doi: 10.1016/j.jtho.2021.06.017
- Zhou F, Guo H, Xia Y, *et al.* The changing treatment landscape of EGFR-mutant non-small-cell lung cancer. *Nat Rev Clin Oncol.* 2025;22(2):95-116.
doi: 10.1038/s41571-024-00971-2
- Wu YL, Guarneri V, Voon PJ, *et al.* Tepotinib plus osimertinib in patients with EGFR-mutated non-small-cell lung cancer with MET amplification following progression on first-line osimertinib (INSIGHT 2): a multicentre, open-label, phase 2 trial. *Lancet Oncol.* 2024;25(8):989-1002.
doi: 10.1016/s1470-2045(24)00270-5
- Kuo P, Elboudwarej E, Zavodovskaya M, *et al.* Trop-2 expression in non-small cell lung cancer. *PLoS One.* 2025;20(4):e0321555.
doi: 10.1371/journal.pone.0321555
- Planchard D, Jänne PA, Cheng Y, *et al.* Osimertinib with or without Chemotherapy in EGFR-Mutated Advanced NSCLC. *N Engl J Med.* 2023;389(21):1935-1948.
doi: 10.1056/NEJMoa2306434
- Hendriks LE, Kerr KM, Menis J, *et al.* Non-oncogene-addicted metastatic non-small-cell lung cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. *Ann Oncol.* 2023;34(4):358-376.
doi: 10.1016/j.annonc.2022.12.013
- Tian Z, Cen L, Wei F, *et al.* EGFR mutations in non-small cell lung cancer: Classification, characteristics and resistance to third-generation EGFR-tyrosine kinase inhibitors (Review). *Oncol Lett.* 2025;30(2):375.
doi: 10.3892/ol.2025.15121
- Tsuboi M, Herbst RS, John T, *et al.* Overall Survival with Osimertinib in Resected EGFR-Mutated NSCLC. *N Engl J Med.* 2023;389(2):137-147.
doi: 10.1056/NEJMoa2304594
- Okuma Y, Kubota K, Shimokawa M, *et al.* First-Line Osimertinib for Previously Untreated Patients With NSCLC and Uncommon EGFR Mutations: The UNICORN Phase 2 Nonrandomized Clinical Trial. *JAMA Oncol.* 2024;10(1):43-51.
doi: 10.1001/jamaoncol.2023.5013
- Sands J, Ahn MJ, Lisberg A, *et al.* Datopotamab Deruxtecan in Advanced or Metastatic Non-Small Cell Lung Cancer With Actionable Genomic Alterations: Results From the Phase II TROPION-Lung05 Study. *J Clin Oncol.* 2025;43(10):1254-1265.
doi: 10.1200/jco-24-01349
- Pellini B, Chaudhuri AA. Circulating Tumor DNA Minimal

- Residual Disease Detection of Non-Small-Cell Lung Cancer Treated With Curative Intent. *J Clin Oncol.* 2022;40(6):567-575.
doi: 10.1200/jco.21.01929
15. Tran HT, Heeke S, Sujit S, *et al.* Circulating tumor DNA and radiological tumor volume identify patients at risk for relapse with resected, early-stage non-small-cell lung cancer. *Ann Oncol.* 2024;35(2):183-189.
doi: 10.1016/j.annonc.2023.11.008
 16. Zalaquett Z, Catherine Rita Hachem M, Kassiss Y, *et al.* Acquired resistance mechanisms to osimertinib: The constant battle. *Cancer Treat Rev.* 2023;116:102557.
doi: 10.1016/j.ctrv.2023.102557
 17. Liu HR. Liquid biopsy – driven cancer genetics in the ultrasensitive and hyper-data-integration eras. *Cancer Adv.* 2026;9:e26001-e26032.
doi: 10.53388/2026926005
 18. Soria JC, Ohe Y, Vansteenkiste J, *et al.* Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med.* 2018;378(2):113-125.
doi: 10.1056/NEJMoa1713137
 19. Engelman JA, Zejnullahu K, Mitsudomi T, *et al.* MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science.* 2007;316(5827):1039-1043.
doi: 10.1126/science.1141478
 20. Batra U, Biswas B, Prabhash K, Krishna MV. Differential clinicopathological features, treatments and outcomes in patients with Exon 19 deletion and Exon 21 L858R EGFR mutation-positive adenocarcinoma non-small-cell lung cancer. *BMJ Open Respir Res.* 2023;10(1):e001492.
doi: 10.1136/bmjresp-2022-001492
 21. Ramalingam SS, Vansteenkiste J, Planchard D, *et al.* Overall Survival with Osimertinib in Untreated, EGFR-Mutated Advanced NSCLC. *N Engl J Med.* 2020;382(1):41-50.
doi: 10.1056/NEJMoa1913662
 22. Robichaux JP, Le X, Vijayan RSK, *et al.* Structure-based classification predicts drug response in EGFR-mutant NSCLC. *Nature.* 2021;597(7878):732-737.
doi: 10.1038/s41586-021-03898-1
 23. Zhang L, Zhang X, Wu L, *et al.* Aumolertinib as adjuvant therapy in resected EGFR-mutated non-small-cell lung cancer (ARTS): a double-blind, multicentre, randomised, controlled, phase 3 trial. *Lancet Oncol.* 2026;27(2):159-168.
doi: 10.1016/s1470-2045(25)00643-6
 24. Herbst RS, John T, Grohé C, *et al.* Molecular residual disease analysis of adjuvant osimertinib in resected EGFR-mutated stage IB-IIIa non-small-cell lung cancer. *Nat Med.* 2025;31(6):1958-1968.
doi: 10.1038/s41591-025-03577-y
 25. Abbosh C, Birkbak NJ, Wilson GA, *et al.* Phylogenetic ctDNA analysis depicts early-stage lung cancer evolution. *Nature.* 2017;545(7655):446-451.
doi: 10.1038/nature22364
 26. Cui Q, Zhang JT, Yin H, *et al.* Comparison of circulating tumor DNA assays for molecular residual disease detection in non-small cell lung cancer. *J Thorac Oncol.* 2026;104090.
doi: 10.1016/j.jtho.2026.104090
 27. Shi C, Zhang C, Fu Z, *et al.* Antitumor activity of aumolertinib, a third-generation EGFR tyrosine kinase inhibitor, in non-small-cell lung cancer harboring uncommon EGFR mutations. *Acta Pharm Sin B.* 2023;13(6):2613-2627.
doi: 10.1016/j.apsb.2023.03.007
 28. Shi Y, Wu L, Ji Y, *et al.* Efficacy and safety of limertinib versus gefitinib as first-line treatment for locally advanced or metastatic non-small-cell lung cancer with EGFR-sensitising mutation: a randomised, double-blind, double-dummy, phase 3 trial. *Lancet Respir Med.* 2025;13(8):677-686.
doi: 10.1016/s2213-2600(25)00121-3
 29. Shi Y, Li B, Wu L, *et al.* Efficacy and Safety of Limertinib (ASK120067) in Patients With Locally Advanced or Metastatic EGFR Thr790Met-Mutated NSCLC: A Multicenter, Single-Arm, Phase 2b Study. *J Thorac Oncol.* 2022;17(10):1205-1215.
doi: 10.1016/j.jtho.2022.05.011
 30. Qin K, Hong L, Zhang J, Le X. MET Amplification as a Resistance Driver to TKI Therapies in Lung Cancer: Clinical Challenges and Opportunities. *Cancers (Basel).* 2023;15(3):612.
doi: 10.3390/cancers15030612
 31. Peng LX, Jie GL, Li AN, *et al.* MET amplification identified by next-generation sequencing and its clinical relevance for MET inhibitors. *Exp Hematol Oncol.* 2021;10(1):52.
doi: 10.1186/s40164-021-00245-y
 32. Caparica R, Yen CT, Coudry R, *et al.* Responses to Crizotinib Can Occur in High-Level MET-Amplified Non-Small Cell Lung Cancer Independent of MET Exon 14 Alterations. *J Thorac Oncol.* 2017;12(1):141-144.
doi: 10.1016/j.jtho.2016.09.116
 33. Wu YL, Yao Y, Yang JJ, *et al.* Vebreltinib in MET amplification-driven advanced non-small-cell lung cancer (KUNPENG): A single-arm, multi-cohort, multicentre, phase 2 study. *Lancet Oncol.* 2026;27(1):36-44.
doi: 10.1016/s1470-2045(25)00594-7
 34. Rolfo C, O'Brate A, Menzel C, *et al.* Liquid and Tissue Biopsies for Identifying MET Exon 14 Skipping NSCLC:

- Analyses from the Phase II VISION Study of Tepotinib. *Clin Cancer Res.* 2025;31(13):2675-2684.
doi: 10.1158/1078-0432.Ccr-24-4097
35. Hartmaier RJ, Markovets AA, Ahn MJ, *et al.* Osimertinib + Savolitinib to Overcome Acquired MET-Mediated Resistance in Epidermal Growth Factor Receptor-Mutated, MET-Amplified Non-Small Cell Lung Cancer: TATTON. *Cancer Discov.* 2023;13(1):98-113.
doi: 10.1158/2159-8290.Cd-22-0586
 36. Sun X, Jia L, Wang T, *et al.* Trop2 binding IGF2R induces gefitinib resistance in NSCLC by remodeling the tumor microenvironment. *J Cancer.* 2021;12(17):5310-5319.
doi: 10.7150/jca.57711
 37. Fang W, Wu L, Meng X, *et al.* Sacituzumab Tirumotecan in EGFR-TKI-Resistant, EGFR-Mutated Advanced NSCLC. *N Engl J Med.* 2026;394(1):13-26.
doi: 10.1056/NEJMoa2512071
 38. Killock D. Sacituzumab tirumotecan improves OS in EGFR TKI-resistant NSCLC. *Nat Rev Clin Oncol.* 2025;22(12):899.
doi: 10.1038/s41571-025-01090-2
 39. Ahn MJ, Tanaka K, Paz-Ares L, *et al.* Datopotamab Deruxtecán Versus Docetaxel for Previously Treated Advanced or Metastatic Non-Small Cell Lung Cancer: The Randomized, Open-Label Phase III TROPION-Lung01 Study. *J Clin Oncol.* 2025;43(3):260-272.
doi: 10.1200/jco-24-01544
 40. Johnson M, Garassino MC, Mok T, Mitsudomi T. Treatment strategies and outcomes for patients with EGFR-mutant non-small cell lung cancer resistant to EGFR tyrosine kinase inhibitors: Focus on novel therapies. *Lung Cancer.* 2022;170:41-51.
doi: 10.1016/j.lungcan.2022.05.011
 41. Vokes NI, Chambers E, Nguyen T, *et al.* Concurrent TP53 Mutations Facilitate Resistance Evolution in EGFR-Mutant Lung Adenocarcinoma. *J Thorac Oncol.* 2022;17(6):779-792.
doi: 10.1016/j.jtho.2022.02.011
 42. Coleman N, Hong L, Zhang J, Heymach J, Hong D, Le X. Beyond epidermal growth factor receptor: MET amplification as a general resistance driver to targeted therapy in oncogene-driven non-small-cell lung cancer. *ESMO Open.* 2021;6(6):100319.
doi: 10.1016/j.esmoop.2021.100319
 43. Shiba-Ishii A, Takemura N, Kawai H, Matsubara D. Histologic transformation of non-small-cell lung cancer in response to tyrosine kinase inhibitors: Current knowledge of genetic changes and molecular mechanisms. *Cancer Sci.* 2024;115(7):2138-2146.
doi: 10.1111/cas.16192
 44. Chhouri H, Alexandre D, Grumolato L. Mechanisms of Acquired Resistance and Tolerance to EGFR Targeted Therapy in Non-Small Cell Lung Cancer. *Cancers (Basel).* 2023;15(2):504.
doi: 10.3390/cancers15020504
 45. Chmielecki J, Gray JE, Cheng Y, *et al.* Candidate mechanisms of acquired resistance to first-line osimertinib in EGFR-mutated advanced non-small cell lung cancer. *Nat Commun.* 2023;14(1):1070.
doi: 10.1038/s41467-023-35961-y
 46. Lu C, Wei XW, Wang Z, *et al.* Allelic Context of EGFR C797X-Mutant Lung Cancer Defines Four Subtypes With Heterogeneous Genomic Landscape and Distinct Clinical Outcomes. *J Thorac Oncol.* 2024;19(4):601-612.
doi: 10.1016/j.jtho.2023.11.016
 47. Wang R, Chen Y, Li L, Zhang L, Zhang S. Osimertinib acquired resistance among patients with EGFR-mutated NSCLC: from molecular mechanisms to clinical therapeutic strategies. *Cancer Drug Resist.* 2025;8:61.
doi: 10.20517/cdr.2025.140
 48. Tamiya A, Osuga M, Harada D, *et al.* Mechanisms of resistance and correlation between pre-treatment co-alterations and p-prognosis to osimertinib in chemo-naïve advanced non-small cell lung cancer. *Lung Cancer.* 2024;195:107917.
doi: 10.1016/j.lungcan.2024.107917
 49. Chmielecki J, Mok T, Wu YL, *et al.* Analysis of acquired resistance mechanisms to osimertinib in patients with EGFR-mutated advanced non-small cell lung cancer from the AURA3 trial. *Nat Commun.* 2023;14(1):1071.
doi: 10.1038/s41467-023-35962-x
 50. Catania C, Liu SV, Garassino M, *et al.* Correlation between treatments and outcomes of patients with EGFR-mutated non-small-cell lung cancer that transitioned into small-cell lung cancer: an international retrospective study. *ESMO Open.* 2025;10(7):105326.
doi: 10.1016/j.esmoop.2025.105326
 51. Saalfeld FC, Möller J, Christopoulos P, *et al.* Small cell transformation in EGFR-mutated non-small cell lung cancer: DLL3 expression and efficacy of immune checkpoint inhibitors or tyrosine kinase inhibitors combined with chemotherapy. *Eur J Cancer.* 2024;213:115065.
doi: 10.1016/j.ejca.2024.115065
 52. Chen Y, He M, Dai Z, *et al.* Clinical and molecular profiling of EGFR-mutant lung adenocarcinomas transformation to small cell lung cancer during TKI treatment. *Front Oncol.* 2023;13:1308313.
doi: 10.3389/fonc.2023.1308313
 53. Dela Cruz MCP, Medina PMB. Epithelial-mesenchymal

- transition (EMT) and its role in acquired epidermal growth factor receptor-tyrosine kinase inhibitor (EGFR-TKI) chemoresistance in non-small cell lung cancer (NSCLC). *Cancer Pathog Ther.* 2025;3(3):215-225.
doi: 10.1016/j.cpt.2024.07.001
54. Lailier C, Didelot A, Garinet S, *et al.* PrP(C) controls epithelial-to-mesenchymal transition in EGFR-mutated NSCLC: implications for TKI resistance and patient follow-up. *Oncogene.* 2024;43(37):2781-2794.
doi: 10.1038/s41388-024-03130-0
 55. Lim SM, Fujino T, Kim C, *et al.* BBT-176, a Novel Fourth-Generation Tyrosine Kinase Inhibitor for Osimertinib-Resistant EGFR Mutations in Non-Small Cell Lung Cancer. *Clin Cancer Res.* 2023;29(16):3004-3016.
doi: 10.1158/1078-0432.Ccr-22-3901
 56. Su C, Sun SY. Fourth-generation epidermal growth factor receptor-tyrosine kinases inhibitors: hope and challenges. *Transl Cancer Res.* 2024;13(8):3929-3934.
doi: 10.21037/tcr-24-406
 57. Xu J, Zhu A, Xu Y, *et al.* Real-world analysis of subsequent-line strategies for EGFR C797S mutant non-small cell lung cancer patients acquired resistance to third-generation EGFR tyrosine kinase inhibitors. *BMC Cancer.* 2025;25(1):1879.
doi: 10.1186/s12885-025-15335-x
 58. Hu D, Hu Y, Lei S, Wu D, Wang Y. MET tyrosine kinase inhibitors in combination with EGFR tyrosine kinase inhibitors in NSCLC patients with EGFR mutations and acquired *MEt* alterations: a systematic review and meta-analysis. *BMC Cancer.* 2025;25(1):732.
doi: 10.1186/s12885-025-14145-5
 59. Zhou X, Zeng L, Huang Z, *et al.* Strategies Beyond 3rd EGFR-TKI Acquired Resistance: Opportunities and Challenges. *Cancer Med.* 2025;14(9):e70921.
doi: 10.1002/cam4.70921
 60. Ahn MJ, Cho BC, Felip E, *et al.* Tarlatamab for Patients with Previously Treated Small-Cell Lung Cancer. *N Engl J Med.* 2023;389(22):2063-2075.
doi: 10.1056/NEJMoa2307980
 61. Cooper AJ, Herzberg B, Parma M, *et al.* Multi-Institution Analysis of Tarlatamab for the Treatment of EGFR-Mutant Transformed Small Cell Lung Cancer. *JCO Precis Oncol.* 2025;9:e2500538.
doi: 10.1200/po-25-00538
 62. Papadimitrakopoulou VA, Mok TS, Han JY, *et al.* Osimertinib versus platinum-pemetrexed for patients with EGFR T790M advanced NSCLC and progression on a prior EGFR-tyrosine kinase inhibitor: AURA3 overall survival analysis. *Ann Oncol.* 2020;31(11):1536-1544.
doi: 10.1016/j.annonc.2020.08.2100
 63. Zhang D, Zhao J, Yang Y, *et al.* Fourth-generation EGFR-TKI to overcome C797S mutation: past, present, and future. *J Enzyme Inhib Med Chem.* 2025;40(1):2481392.
doi: 10.1080/14756366.2025.2481392
 64. Corvaja C, Passaro A, Attili I, *et al.* Advancements in fourth-generation EGFR TKIs in EGFR-mutant NSCLC: Bridging biological insights and therapeutic development. *Cancer Treat Rev.* 2024;130:102824.
doi: 10.1016/j.ctrv.2024.102824
 65. Yamaoka T, Ohmori T, Ohba M, *et al.* Acquired Resistance Mechanisms to Combination Met-TKI/EGFR-TKI Exposure in Met-Amplified EGFR-TKI-Resistant Lung Adenocarcinoma Harboring an Activating EGFR Mutation. *Mol Cancer Ther.* 2016;15(12):3040-3054.
doi: 10.1158/1535-7163.Mct-16-0313
 66. Takezawa K, Pirazzoli V, Arcila ME, *et al.* HER2 amplification: a potential mechanism of acquired resistance to EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFR T790M mutation. *Cancer Discov.* 2012;2(10):922-933.
doi: 10.1158/2159-8290.Cd-12-0108
 67. Rock A, Mambetsariev I, Pareek S, *et al.* Small Cell Transformation of EGFR-Mutant NSCLC Treated with Tyrosine Kinase Inhibition. *Curr Oncol.* 2025;32(10):554.
doi: 10.3390/curroncol32100554
 68. Zhu X, Chen L, Liu L, Niu X. EMT-Mediated Acquired EGFR-TKI Resistance in NSCLC: Mechanisms and Strategies. *Front Oncol.* 2019;9:1044.
doi: 10.3389/fonc.2019.01044
 69. Gray JE, Markovets A, Reungwetwattana T, *et al.* Longitudinal Analyses of Circulating Tumor DNA for the Detection of EGFR Mutation-Positive Advanced NSCLC Progression During Treatment: Data From FLAURA and AURA3. *J Thorac Oncol.* 2024;19(11):1525-1538.
doi: 10.1016/j.jtho.2024.07.008
 70. Leonetti A, Verzè M, Minari R, *et al.* Resistance to osimertinib in advanced EGFR-mutated NSCLC: a prospective study of molecular genotyping on tissue and liquid biopsies. *Br J Cancer.* 2024;130(1):135-142.
doi: 10.1038/s41416-023-02475-9
 71. Gale D, Heider K, Ruiz-Valdepenas A, *et al.* Residual ctDNA after treatment predicts early relapse in patients with early-stage non-small cell lung cancer. *Ann Oncol.* 2022;33(5):500-510.
doi: 10.1016/j.annonc.2022.02.007
 72. Jeon H, Wang S, Song J, Gill H, Cheng H. Update 2025: Management of Non-Small-Cell Lung Cancer. *Lung.*

- 2025;203(1):53.
doi: 10.1007/s00408-025-00801-x
73. Ernst SM, Aldea M, von der Thüsen JH, *et al.* Utilizing ctDNA to discover mechanisms of resistance to targeted therapies in patients with metastatic NSCLC: towards more informative trials. *Nat Rev Clin Oncol.* 2025;22(5):371-378.
doi: 10.1038/s41571-025-01011-3
 74. Yamaguchi T, Masago K, Sasaki E, Kuroda H, Matsushita H, Horio Y. EML4-ALK Variant 3a/b as a mechanism of osimertinib resistance in a patient with EGFR L858R positive NSCLC. *Cancer Genet.* 2024;280-281:13-16.
doi: 10.1016/j.cancergen.2023.12.001
 75. de Langen AJ, Johnson ML, Mazieres J, *et al.* Sotorasib versus docetaxel for previously treated non-small-cell lung cancer with KRAS(G12C) mutation: a randomised, open-label, phase 3 trial. *Lancet.* 2023;401(10378):733-746.
doi: 10.1016/s0140-6736(23)00221-0
 76. Jänne PA, Riely GJ, Gadgeel SM, *et al.* Adagrasib in Non-Small-Cell Lung Cancer Harboring a KRAS(G12C) Mutation. *N Engl J Med.* 2022;387(2):120-131.
doi: 10.1056/NEJMoa2204619
 77. de Marinis F, Kim TM, Bonanno L, *et al.* Savolitinib plus osimertinib in epidermal growth factor receptor (EGFR)-mutated advanced non-small cell lung cancer with MET overexpression and/or amplification following disease progression on osimertinib: primary results from the phase II SAVANNAH study. *Ann Oncol.* 2025;36(8):920-933.
doi: 10.1016/j.annonc.2025.04.003
 78. Nogami N, Barlesi F, Socinski MA, *et al.* IMpower150 Final Exploratory Analyses for Atezolizumab Plus Bevacizumab and Chemotherapy in Key NSCLC Patient Subgroups With EGFR Mutations or Metastases in the Liver or Brain. *J Thorac Oncol.* 2022;17(2):309-323.
doi: 10.1016/j.jtho.2021.09.014
 79. Trontzas IP, He M, Wurtz A, *et al.* Quantitative Protein Expression of Antibody-Drug Conjugate Targets in EGFR Mutated and Wild-type Non-Small Cell Lung Cancer. *Clin Cancer Res.* 2025;31(13):2767-2776.
doi: 10.1158/1078-0432.Ccr-24-3347
 80. Sakata Y, Saito G, Sakata S, *et al.* Osimertinib as first-line treatment for elderly patients with advanced EGFR mutation-positive non-small cell lung cancer in a real-world setting (OSI-FACT-EP). *Lung Cancer.* 2023;186:107426.
doi: 10.1016/j.lungcan.2023.107426
 81. Nguyen TNH, Van QL, Thi Bich PN, *et al.* Therapeutic Outcomes of Osimertinib in EGFR-Mutant Non-Small Cell Lung Cancer With Brain Metastases: Results From a Retrospective Study at Vietnam National Cancer Hospital. *Cancer Control.* 2025;32:10732748251348429.
doi: 10.1177/10732748251348429
 82. Areses Manrique MC, León-Mateos L, Silva-Díaz S, *et al.* Real-world clinical experience with osimertinib in advanced EGFR-mutated non-small cell lung cancer. *Clin Transl Oncol.* 2025;28(2):494-503.
doi: 10.1007/s12094-025-04015-8