

Research Article

Maladaptive Smartphone use among Jordanian Adolescents: Prevalence, Correlations, and the Role of Cognitive Emotion Regulation

Fatimah S. Tarawneh¹, Alaa Abuattallah², Latefa Ali Dardas^{3*}

¹Princess Muna College of Nursing, Mutah University, Al-kark 61710, Jordan

²Department of Maternal and Child Health Nursing, Faculty of Nursing, Applied Science Private University, Faculty of Nursing, Amman 11937, Jordan

³Department of Community Health Nursing, School of Nursing, University of Jordan, Amman 11942, Jordan

Keywords: Maladaptive smartphone use, School adolescents, Jordan, Internet addiction

Health Psychology Research

Vol. 13, 2025

Background

Maladaptive smartphone use (MSU) has emerged as a significant concern, particularly among adolescents. In Jordan, a country with a youthful population and high rates of smartphone engagement, understanding MSU and its psychological impacts is crucial.

Objective

The study aims to (i) determine the point prevalence and patterns of MSU among Jordanian adolescents, and (ii) examine whether cognitive emotion regulation strategies mediate the relationship between MSU and psychological health.

Methods

A cross-sectional, descriptive, correlational survey was conducted. Using a multistage sampling approach, the study targeted students aged 13 – 17 in Jordanian public and private schools, with schools randomly selected from each region. A structured questionnaire assessed sociodemographic data, smartphone addiction, cognitive emotion regulation, and psychological health targeting depression, anxiety, and stress.

Results

Among the 960 students who participated, 36% exhibited signs of MSU. Adolescents with high MSU reported significantly higher maladaptive cognitive emotion regulation strategies than those with low MSU. Factors such as age, gender, socioeconomic status, and academic performance influenced MSU levels. Maladaptive emotion regulation strategies partially mediated the relationship between MSU and psychological health, with high MSU linked to greater levels of emotional distress.

Conclusion

This study reveals a significant prevalence of MSU among Jordanian adolescents, with notable demographic variations. It proposes a link between MSU and psychological health, emphasizing the mediating role of cognitive emotion regulation. These results contribute to the growing body of literature on smartphone use behaviors and their psychological implications, particularly within the context of Jordanian adolescents. The study's insights are relevant for developing targeted interventions to mitigate MSU and its psychological impacts.

*Corresponding author:

Latefa Ali Dardas

Department of Community Health Nursing, School of Nursing, University of Jordan, Amman 11942, Jordan

Email: l.dardas@ju.edu.jo

1. INTRODUCTION

In 2023, the number of smartphone users worldwide reached approximately 6.92 billion, accounting for 86.29% of the global population. This represents 86.29% of the world's population.¹ In 2020, the user base expanded by 93 million, reflecting a 1.8% annual growth rate. Mobile subscriptions have surpassed six billion, and the daily addition of new smartphones in use exceeds one million.² The emergence of multipurpose smartphones and their subsequent widespread adoption have altered the communication and information landscape, reshaped the values and interests of a large number of users, and raised concerns about usage addiction globally.³

Many researchers use the term “smartphone addiction” to describe excessive use of smartphones. However, this term is not included in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, or the upcoming International Classification of Diseases, 11th Revision.³ Instead, the term maladaptive smartphone use (MSU) is increasingly used, indicating excessive behavior with less severe impairment than addiction.⁴ Regardless of the terminology, this phenomenon is widely recognized due to its potentially harmful consequences.⁵

Among adolescents, MSU is particularly evident, and its pattern demonstrates associations with several negative psychosocial and physical consequences, including impulsivity, anxiety, depression, attention deficits,^{6,7} dysfunctional family dynamics, strained relationships with friends,^{8,9} insomnia, neck pain, and other somatic problems.^{10,11} In addition, the heightened utilization of smartphones among adolescents has been linked to compromised cognitive functions and altered brain structure.¹² Together, these findings ought to raise concerns over MSU for clinicians and educators in contemporary society.⁷

Several factors are associated with MSU among adolescents. Cognitive emotion regulation strategies have been reported to play a key role in the process of adaptation and maladaptation to smartphone use.^{13,14} Cognitive emotion regulation refers to the cognitive mechanisms involved in handling emotionally charged information, playing a vital role in the human experience by facilitating the management of emotions following stressful events.¹⁵ As children progress in age, there is a developmental shift in their emotion regulation strategies from external, behavior-oriented approaches to internal, cognition-based methods. Children acquire the ability to regulate their emotions through cognitive processes, including but not limited to self-blame, rumination, catastrophizing, acceptance, planning, and redirecting focus toward positive aspects.¹⁶ A notable finding in existing research is how individuals employ cognitive processes to manage their emotions and how this relates to the extent of MSU. Evidence suggests that individuals who struggle with effective emotion regulation may be more prone to MSU, potentially using it as a coping mechanism to escape or alleviate emotional distress.¹⁷ Understanding these cognitive emotion regulation strategies is crucial in comprehending the intricate relationship between psychological tendencies and excessive smartphone dependency.

Jordan, with 63% of its population under the age of 30, possesses one of the youngest demographics globally. Notably, 32% of individuals aged 15 – 30 in the country are currently unemployed.¹⁸ Jordan, classified by the World Health Organization as a nation requiring assistance, initiated the Mental Health Gap Action Programme in 2021.¹⁹

Although the program has been incorporated into Jordan's health policies and the National Mental Health and Substance Use Action Plan, the availability of mental health services remains constrained. This limitation is attributed to the scarcity of primary health-care clinics, the substantial population of Palestinian and Syrian refugees, and the prevalent cultural stigma surrounding mental health issues.²⁰ A recent study revealed that 59% of Jordanian children engage with YouTube, 25% indulge in gaming, and 23% spend over 5 h daily on the Internet.²¹ Atoum *et al.*,²² called for further investigation on this issue to improve health outcomes in light of the excessive utilization of electronic devices by adolescents in Jordan.

Despite the increasing attention from both scientific literature and mainstream media on MSU as a noteworthy health concern, there remains a paucity of research on this issue in the Arab region in general, and Jordan in particular. The exploration of MSU among Jordanian adolescents and its connection with cognitive emotion regulation strategies emerges as an innovative and crucial research area. Given the widespread integration of smartphones into daily routines, there is a compelling need for a thorough examination of the potential psychological implications within this demographic. Therefore, this study aimed to (i) determine the point prevalence and patterns of MSU among Jordanian adolescents, and (ii) examine whether cognitive emotion regulation strategies mediate the relationship between MSU and psychological health.

2. METHODS

2.1. STUDY DESIGN

A cross-sectional, descriptive, correlational survey design was employed to systematically investigate the study's objectives. Data were collected using a valid and reliable self-reported questionnaire administered to high school students, capturing information related to the study's main variables. The data collection process spanned from February to May 2024, ensuring a comprehensive representation of participants within the designated timeframe. The selection of this research design aligns with the study's objectives, offering a methodologically sound approach to examining the associations among key variables.

2.2. SAMPLING AND POWER ANALYSIS

A multistage sampling method was employed to ensure a representative and methodologically rigorous selection process. In the first stage, the country was stratified into three main regions: northern, central, and southern districts. To maximize school representation, the researcher identified the three governorates with the highest population distribution, as reported by the Department of Statistics in 2023.²³ The selected governorates were Irbid (north), Amman (center), and Al-Karak (south). Based on proportional allocation, approximately 50% of the randomly selected schools were from the central district, 30% from the northern district, and 20% from the southern district. In the second stage, educational directorates within these governorates were randomly chosen from the Ministry of Education's official school list using a simple random selection technique. Specifically, in the northern district, schools were randomly selected from the Irbid Qasabah directorate. In the central district, selections

were made from the Aljamaah and Marka directorates, whereas in the southern district, schools were chosen from the North Mazar directorate. Following this, secondary public schools were stratified by gender, ensuring the inclusion of both male and female schools. Private schools, which accommodate both male and female students within the same institution, were also included. The selection of schools was based on high student density, identified from the Ministry of Education's records. Schools were then randomly selected using a simple random selection method, resulting in an initial total of 16 schools. However, the number of schools was subsequently increased to ensure that the target sample size was met, adjusting based on student response rates. Within each selected school, convenience sampling was employed to recruit participants. The target population comprised male and female students aged 13 – 17 who met the predefined inclusion criteria. This multistage approach ensured a balanced and representative sample, enhancing the generalizability and validity of the study findings. Figure 1 summarizes the sampling procedure.

Cochran's formula, in Equation I,

$$n = (Z^2pq)/e^2 \quad (I)$$

It is particularly well-suited for large populations, such as school students. In this formula, p represents the estimated proportion of the population possessing the characteristic of interest, e denotes the desired level of precision (i.e., the margin of error), and q is defined as $1-p$.²⁴ The Z -value, which corresponds to the desired confidence level, is obtained from a standard Z -table. For this study, using a 95% confidence level ($Z = 1.96$), an estimated proportion (p) of 0.5, and a margin of error (e) of 0.05, the sample size was calculated as follows (Equation II):

$$n = (1.96^2 \times 0.5 \times 0.5)/(0.05)^2 = 384 \quad (II)$$

This calculation indicates that a minimum sample size of 384 participants is required to achieve a 95% confidence level, ensuring that the true population value falls within $\pm 5\%$ of the surveyed value. To account for potential replication errors, attrition, and data entry inaccuracies, a 10% buffer was added, increasing the minimal required sample size to 435 participants.




Figure 1. Sampling flowchart

2.3. MEASUREMENT

A structured questionnaire comprising three sections was used. The first section gathered sociodemographic data, including gender, age, educational level (grades 7 – 11), type of school, parental educational level, place of residence, grade point average (GPA), monthly family income, and perceived parental control over smartphone use both at home and outside. In addition, it collected information on the average hours spent on smartphones during weekends and weekdays, as well as the ownership status of smartphones.

The second section included the smartphone addiction inventory (SPAI), a 26-item self-report measure designed to assess smartphone addiction.²⁵ Each item was rated on a four-point Likert scale: 1 = strongly disagree, 2 = somewhat disagree, 3 = somewhat agree, and 4 = strongly agree. The cutoff score for SPAI is typically set at 64, distinguishing individuals with potential smartphone addiction.²⁵ The SPAI is structured around four key factors: (i) Compulsive behavior, (ii) functional impairment, (iii) withdrawal, and (iv) tolerance. It has been validated across different cultures and maintains a reliable factor structure across various populations,^{26–28} particularly among adolescents.²⁹ The authors of SPAI established internal consistency, reporting a Cronbach's alpha of 0.94 for the total scale, with a 2-week test-retest reliability ranging from 0.80 to 0.91 across its subscales. In the present study, the Cronbach's alpha correlation coefficient was 0.92, confirming strong reliability.

The third section included the Cognitive Emotion Regulation Questionnaire (CERQ),³⁰ a 36-item measure designed to assess cognitive emotion regulation strategies in response to stressful or threatening life events. The CERQ consists of nine distinct subscales, including five adaptive strategies (positive refocusing, acceptance, positive reappraisal, refocusing on planning, and putting into perspective) and four maladaptive strategies (rumination, self-blame, blaming others, and catastrophizing). Each subscale was assessed using four items, rated on a five-point Likert scale from 1 ("Almost never") to 5 ("Almost always"), indicating the frequency of use of each strategy. Higher subscale scores reflect a greater tendency to use a specific cognitive regulation strategy.³¹ This study employed the validated Arabic version of the CERQ,³² which demonstrated strong psychometric properties, with a Cronbach's alpha correlation coefficient of 0.86.

The fourth section included the Depression, Anxiety, and Stress Scale-21 Items (DASS-21), a widely used psychometric instrument designed to measure three psychological states: Depression, anxiety, and stress.³³ The scale consists of 21 items, with seven items per subscale, each assessing core symptoms associated with these emotional states. The depression subscale evaluates low mood, lack of interest in life, low self-esteem, and low motivation. The anxiety subscale measures fear responses and physiological symptoms of anxiety, whereas the stress subscale assesses difficulties in relaxation, heightened nervous arousal, and tendencies to become easily upset or irritable. Each item was rated on a four-point scale, ranging from 0 ("Did not apply to me at all") to 3 ("Applied very much or most of the time"). The DASS-21 categorizes scores into normal, mild, moderate, severe, and extremely severe for each psychological state.³³ This scale has been extensively validated and is widely recognized for its reliability and validity. The Arabic version, validated by Parkitny and McAuley,³⁴ was used in this study, demonstrating a Cronbach's alpha correlation coefficient of 0.920, ensuring strong internal consistency.

2.4. TRANSLATION OF STUDY MEASURES

To ensure linguistic and cultural accuracy, the translation of study instruments followed a rigorous five-stage process. The initial translation involved two bilingual experts independently translating the original English versions of the instruments into Arabic. This was followed by a synthesis process, where a review panel compared the translations and reconciled discrepancies to produce a unified Arabic version. The back-translation stage required two independent translators with no prior exposure to the original instruments to translate the Arabic version back into English. The back-translated English version was then compared with the original English instruments to identify any semantic inconsistencies, and discrepancies were resolved through discussion with linguistic and subject matter experts. Finally, the validation and reliability testing stage involved expert review for linguistic clarity, cultural appropriateness, and religious sensitivity. In addition, Cohen's kappa coefficient and intraclass correlation coefficient were computed to evaluate inter-rater agreement and ensure the reliability of the translation process.³⁵

2.5. ETHICAL CONSIDERATIONS

Approval for ethical clearance was obtained from the University of Jordan's Deanship of Scientific Research Institutional Review Board Committee, followed by formal authorization from the Ministry of Education to conduct data collection. In addition, ethical assurances were secured from the original authors of the research instruments used in this study. To uphold ethical integrity and participant protection, multiple safeguards were implemented.

To ensure ethical compliance and participant protection, parental consent and child assent were obtained following the guidelines of the American Academy of Pediatrics (1976).³⁶ The consent form provided detailed information about the study, including its title, significance, purpose, and the researcher's contact information for any inquiries. Participants were assured that their responses would remain confidential, no personal identifying data would be recorded, and all collected information would be used solely for research purposes.

In collaboration with school counselors, the consent form was distributed to parents through a WhatsApp group for adolescents meeting the inclusion criteria. To enhance parental understanding of the study's objectives and procedures, the form explicitly outlined the study's aim, participation requirements, and contact details for further inquiries. A passive consent approach was adopted, whereby only parents who objected to their child's participation informed the school principals or counselors.

Adolescents were also provided with assent forms to ensure they were fully aware of their participation and could give voluntary consent. Both parents and students were clearly informed about the voluntary nature of the study, any potential risks, and their right to withdraw at any time without consequences. To maintain participant anonymity, all names were coded for data analysis, and confidentiality was strictly upheld. Physical copies of the questionnaires were securely stored in a locked location accessible only to the researcher. Electronic data, including survey files and recordings, was stored on a password-protected computer. Personally identifiable information was maintained in a separate coded file, with access restricted to the researcher and supervisor to ensure the highest level of data security and participant privacy.

2.6. STATISTICAL ANALYSIS

Data were analyzed using both descriptive and inferential statistical techniques. Descriptive statistics, including means, standard deviations, frequencies, and percentages, were used to summarize participants' demographic characteristics and key study variables. Inferential analyses were conducted to examine group differences and relationships among variables.

Specifically, one-way Analysis of Variance (ANOVA) with Scheffé *post hoc* tests was used to assess differences in MSU scores across sociodemographic subgroups such as age, gender, GPA, income, and sleep quality. Multivariate ANOVA (MANOVA) was performed to evaluate the impact of these factors on MSU subscale scores (compulsive behavior, functional impairment, withdrawal, and tolerance). Independent-sample t-tests were used to compare cognitive emotion regulation strategy scores between adolescents with high and low levels of MSU. Assumptions for each test were assessed and met.

To test the hypothesized mediation model, Partial Least Squares Structural Equation Modeling was conducted using SmartPLS version 4.0, which allowed for the analysis of complex relationships between MSU, cognitive emotion regulation strategies, and psychological health (depression, anxiety, and stress). Model fit was assessed using multiple fit indices, including the Chi-square test, root mean square error of approximation (RMSEA), and comparative fit index (CFI).

All statistical analyses were performed using IBM Statistical Package for the Social Sciences statistics version 28 and SmartPLS version 4.0. $p < 0.05$ was considered to be statistically significant.

3. RESULTS

3.1. PARTICIPANTS' DEMOGRAPHIC CHARACTERISTICS

A total of 960 students, comprising 550 girls (57.3%) and 410 boys (42.7%), completed and returned the study questionnaire. Participants were distributed across Jordan as follows: 24.7% in the north ($n = 237$), 62.5% ($n = 600$) in the middle, and 12.8% ($n = 123$) in the south. The mean age of the students was 14.94 (standard deviation [SD] = 1.2) with a range of 13 – 17 years. The majority of the participants were ninth-grade students (32.5%). Of the total sample, 58.9% attended public schools, whereas 41.1% were from private schools. About 43.2% had an excellent GPA of 90% and more (mean $n = 85.7$, SD = 10.2), and approximately 59.4% of them studied for an average of 1 – 3 h/day. Regarding parental education levels, 50.6% of the parents held a Bachelor's degree or higher, whereas 3.9% were illiterate. In terms of maternal education, 44% of the mothers had attained a Bachelor's degree or higher, whereas 1.8% were illiterate. Overall, maternal education levels were higher compared to paternal education levels. For this survey result, 33.4% had a family monthly income between 501 and 1,000 Jordanian Dinar (JD)/month. Table 1 details all demographic characteristics of respondents.

3.2. SMARTPHONE USAGE PROFILE

The mean score for the MSU was 59.07 (SD = 16.07), ranging from 26 to 103 based on the total score's distribution.

Table 1. Demographic characteristics of respondents

Variables	Mean	Standard deviation
Age	14.94	1.2
Grade point average	85.7	10.2
Variables	Frequency (n)	Percentage (%)
Gender		
Female	550	57.3
Male	410	42.7
Age		
13 years	136	14.2
14 years	212	22
15 years	285	29.7
16 years	228	23.8
17 years	99	10.3
Residence		
North	237	24.7
Middle	600	62.5
South	123	12.8
Type of school		
Public	565	58.9
Private	395	41.1
Fathere education		
Illiterate	37	3.9
11 th grade and below	93	9.7
Tawjihi (high school)	232	24.2
Diploma	112	11.7
Bachelor's degree or higher	486	50.6
Mother's education		
Illiterate	17	1.8
11 th grade and below	81	8.4
Tawjihi (high school)	264	27.5
Diploma	176	18.3
Bachelor's degree or higher	422	44.0
Family monthly income		
<150 JD	34	3.5
150 – 300 JD	121	12.6
301 – 500 JD	207	21.6
501 – 1,000 JD	321	33.4
1001 – 1,500 JD	122	12.7
1501 – 2,000 JD	155	16.1
Grade point average		
Excellent (90 – 100%)	415	43.2
Very good (80 – 89%)	333	34.7
Good (70 – 79%)	139	14.5
Moderate (60 – 69%)	58	6
Weak (e59%)	15	1.6
Practices a hobby		
Yes	633	65.9
No	327	34.1
Average hours of study per day		
<1 h	153	15.9
1 – 3 h	570	59.4
4 – 6 h	207	21.6
7 – 9 h	26	2.7
More than 9 h	4	0.4

Abbreviation: JD: Jordanian Dinar.

Almost 64% of the sample was under the cutoff score of 64, as the prevalence of MSU among school adolescents was 36%, where 57.3% were females (Table 2).

One-way ANOVA was conducted to examine differences in MSU scores across various demographic factors, including gender, school type, age, GPA, family income, and perceived quality of sleep (Table S1). Scheffe *post hoc* tests were used for multiple comparisons where significant differences were observed. The results indicated that age significantly impacted MSU scores ($F [4, 955] = 2.51, p=0.040$), with students aged 15 – 17 reporting significantly higher MSU scores than those aged 13 – 14 ($p<0.01$). Gender differences were also significant, with female students exhibiting higher MSU scores than male students ($F [1, 955] = 60.42, p<0.001$). Academic performance was inversely related to MSU, as students with lower GPAs reported significantly higher MSU scores than those with excellent GPAs ($F [4, 955] = 5.86, p<0.001$). Similarly, family income was a significant factor ($F [5, 954] = 3.09, p=0.009$), with students from families earning 301 – 500 JDs/month displaying higher MSU scores than those from families earning 1,001 – 1,500 JDs/month ($p=0.044$). Perceived sleep quality showed a strong association with MSU ($F [2, 957] = 73.22, p<0.001$), with students reporting poor sleep quality exhibiting significantly higher MSU scores than those with moderate or good sleep quality ($p<0.001$). Conversely, school type, grade level, residence, and parental education level did not show a significant impact on MSU scores ($p>0.05$).

A MANOVA was conducted to analyze differences in MSU subscales (compulsive behavior, functional impairment, withdrawal, and tolerance) based on sociodemographic characteristics (Table S2). Preliminary assumption testing confirmed no serious violations in normality, linearity, variance homogeneity, or multicollinearity. School type significantly influenced compulsive behavior, with public school students scoring higher than private school students ($F [1, 958] = 4.34, p<0.05$). Age also had a significant effect on compulsive behavior ($F [4, 955] = 4.42, p<0.001$), with older adolescents (15 – 17 years) exhibiting higher scores than younger students. Gender differences were evident across all MSU subscales, with female students scoring significantly higher than male students in compulsive behavior ($F [1, 958] = 47.17, p<0.001$), functional impairment ($F [1, 958] = 42.22, p<0.001$), withdrawal ($F [2, 958] = 47.71, p<0.001$), and tolerance ($F [2, 958] = 40.55, p<0.001$). GPA was inversely correlated with compulsive behavior, functional impairment, and withdrawal symptoms, with students having lower GPAs exhibiting

higher MSU subscale scores ($p<0.001$). *Post hoc* tests indicated that students with weak, moderate, and good GPAs had significantly higher scores in compulsive behavior, functional impairment, and withdrawal symptoms compared to those with excellent GPAs ($p<0.01$). Residence also played a role, particularly in functional impairment ($F [2, 957] = 5.90, p<0.001$). Students from southern regions exhibited significantly higher functional impairment than those from central regions ($p<0.05$). Overall, these findings highlight that age, gender, GPA, family income, and sleep quality significantly influence MSU prevalence and severity, with older adolescents, females, lower academic achievers, and those with poor sleep quality being at higher risk for compulsive smartphone behaviors and psychological impairments.

3.3. PARTICIPANTS' PSYCHOLOGICAL HEALTH AND COPING STRATEGIES

The mean score for the DASS-21 scale was 24.87 (SD = 13.79), ranging from 0 to 63. In particular, the mean scores for depression, anxiety, and stress were 15.62 (SD = 10.02), 16.14 (SD = 10.26), and 18.00 (SD = 10.12), respectively. Based on the DASS classification of depression severity, 29% of the students did not report any depressive symptoms, 17.1% reported mild depression, 25% reported moderate depression, 13% reported severe depression, and 15.9% reported extremely severe depression. As for the classifications of anxiety, 22.5% did not report any anxiety symptoms, 8.4% reported mild anxiety, 18.9% reported moderate anxiety, 13.6% reported severe anxiety, and 36.6% reported extremely severe anxiety. Regarding reports on stress levels, 42.4% did not report any stress symptoms, 14.5% reported mild stress levels, 19.3% reported moderate stress levels, 14.2% reported severe stress levels, and 9.7% reported having extremely severe stress symptoms. Based on the reports on the CERQ, the mean score of the adaptive subscale score was 69.10 (SD = 12.91), ranging from 20 to 99. The mean score of the maladaptive subscale was 51.31 (SD = 12.25). Refocusing on planning, positive reappraisal, and rumination were among the most commonly used coping mechanisms. Table 3 details the students' responses on all subscales.

3.4. COGNITIVE REGULATION STRATEGIES AMONG ADOLESCENTS

An independent sample *t*-test was used to explore differences in mean scores of cognitive emotion regulation strategies between Jordanian school adolescents with high and low levels of MSU. All test assumptions were first examined to ensure that the data were compatible with the test. The analysis revealed a statistically significant difference ($p<0.001$) in the mean scores of maladaptive cognitive regulation strategies between those who had a high level of MSU (mean $n = 55.31$, SD = 10.98) and a low level of MSU (mean $n = 47.52$, SD = 12.20). In particular, those with high MSU levels reported significantly higher maladaptive cognitive regulation strategies mean scores than those who had low MSU levels.

3.5. COPING STRATEGIES AS POTENTIAL MEDIATORS BETWEEN MSU AND PSYCHOLOGICAL HEALTH

Preliminary and correlation analyses between the study variables were conducted before model testing. The conceptual

Table 2. Prevalence of maladaptive smartphone use among school adolescents

Prevalence of maladaptive smartphone use	Statistical values
Mean±standard deviation	59.07±16.07
Median	58
25 th , 75 th percentile	46, 70
Minimum, maximum values	26, 103
Scores	<i>n</i> (%)
>64 (cutoff score)	347 (36.2)
<64 (cutoff score)	613 (63.8)
<46	222 (23.1)
46 – 58	271 (28.2)
59 – 70	227 (23.6)
>70	240 (25)

Table 3. Reports on study variables from students (n=960)

Study measure	(Mean±standard deviation)	P25	Median	P75	Minimum value	Maximum value
The Smartphone Addiction Inventory Scale	59.07±16.07	46	58	70	26	103
Compulsive behavior subscale	19.74±5.51	15	19	24	9	36
Functional impairment subscale	18.09±5.73	14	18	22	8	32
Withdrawal subscale	13.69±4.52	10	13	17	6	24
Tolerance subscale	7.56±2.35	6	7	9	3	12
Adaptive CER Questionnaire	69.10±12.91	61	70	78	20	99
Maladaptive CER Questionnaire	51.31±12.25	43	52	60	16	79
Self-blame subscale	12.74±4.00	10	13	16	4	20
Acceptance subscale	12.42±3.78	10	13	15	4	20
Rumination subscale	14.63±4.14	12	15	18	4	20
Positive refocusing subscale	13.25±4.30	10	13	17	4	20
Refocus on the planning subscale	14.62±4.01	12	15	18	4	20
Positive reappraisal subscale	14.84±3.89	12	15	18	4	20
Putting into perspective subscale	13.97±3.63	12	14	17	4	20
Catastrophizing subscale	12.46±4.21	9	13	16	4	20
Other-blame subscale	11.48±4.43	8	12	15	4	20
Depression, Anxiety, and Stress Scale	24.87±13.79	14	23	34	0	63
Depression subscale	7.81±5.01	4	7	11	0	21
Anxiety subscale	8.07±5.13	4	8	12	0	21
Stress subscale	9.00±5.06	5	9	12	0	21

Abbreviation: CER: Cognitive emotion regulation.

mediation model was then fitted to the data with SmartPLS 4 (version 4.0) to examine coping strategies as potential mediators in the relationship between MSU and psychological health (i.e., depression, anxiety, and stress). The model demonstrated satisfactory model fit, with the following indices: χ^2 of 6.195, $p=0.001$, RMSEA of 0.074, and CFI of 0.925. As shown in Figure 2, MSU had a significant positive direct effect on psychological health ($b = 0.255$, $p<0.001$) and maladaptive coping strategies ($b = 0.301$, $p<0.001$). In addition, maladaptive coping strategies had a significant positive direct effect on psychological health ($b = 0.811$, $p<0.001$). In contrast, adaptive coping strategies had a significant negative effect on psychological health ($b = -0.287$, $p<0.001$). Moreover, the findings showed that maladaptive coping strategies partially mediated the effects of MSU on psychological health ($b = 0.244$, $p<0.001$).

4. DISCUSSION

The primary aim of this study was to examine MSU among Jordanian school adolescents, focusing on its prevalence, psychosocial correlations, and the role of cognitive emotion regulation strategies. This study revealed that the prevalence of MSU among Jordanian school adolescents was 36%. To accurately interpret these findings, they should be considered within the broader context of both Eastern and Western literature. Extremera *et al.*¹⁵ reported a 41.9% prevalence of MSU among Spanish adolescents (ages 14 – 18), with no significant gender differences. In Southern Thailand, Kim *et al.*³⁷ found an alarming 70.3% prevalence of MSU among Thai Muslim adolescents. In India, MSU rates ranged from 39% to 67%,³⁸ while in Indonesia, 11.4% of junior high students exhibited MSU symptoms.³⁹

Among South Korean adolescents, 35.2% were classified as smartphone addicts, with 7.6% at high risk and 27.6% at potential risk.⁴⁰ In the Philippines, the MSU rate was 62.6%,

with a higher prevalence among males (66.2%) than females (60.2%).⁴¹ In Iran, Esmailpour *et al.*⁴² reported a 38.9% prevalence, while in Turkey, Çağan and Koca⁴³ found a 36.9% prevalence, with females (68%) exhibiting higher rates than males (32%). In Saudi Arabia, Al-Amri *et al.*⁴⁴ reported that 49.5% of middle-school adolescents exhibited MSU, with a higher prevalence among females (56%) than males (44%). Similarly, studies in Morocco indicated a high prevalence, with 57.8% of adolescents identified as maladaptive users.⁴⁵ Another study reported higher susceptibility among girls (69.1%) compared to boys (63%).⁴⁶ In Lebanon, 43.4% of adolescents exhibited MSU,⁴⁷ while in Kuwait, the prevalence reached 64.6%.⁴⁸

Despite extensive literature on MSU among adolescents worldwide, no studies have specifically examined its prevalence among Jordanian school students using self-report questionnaires. Existing research focuses primarily on university students and adults.^{49,50} However, an artificial intelligence-based analysis of online surveys indicated that Jordanian parents perceive a growing trend of MSU among adolescents, with signs of addictive behavior becoming a significant social concern.²¹ Regardless of the terminologies and measurement tools used across studies, the prevalence of MSU among Jordanian adolescents appears comparably high, reinforcing the global nature of smartphone addiction and the urgent need for culturally sensitive interventions.

4.1. MSU BASED ON THE DEMOGRAPHIC CHARACTERISTICS OF JORDANIAN ADOLESCENTS

MSU appeared to be significantly higher among older female adolescents from families with low monthly income and those with poor academic performance. Consistent with these findings, other studies have also identified a higher prevalence of MSU among females.^{51,52} Girls were found to be between 2 and 7 times more likely to develop MSU than boys.^{37,53} While the exact reasons for this difference are




Figure 2. Coping strategies partially mediate the relationship between MSU and psychological health

Abbreviations: CER: Cognitive emotion regulation; DASS: Depression, Anxiety, and Stress Scale; MSU: Maladaptive smartphone use.

not fully understood, it may be related to gender-specific patterns of smartphone use. Girls are more inclined to use their devices for communication functions and social networking services, whereas boys are more likely to engage in activities such as gaming, watching videos, and listening to music.⁵⁴ Female adolescents are known to take pleasure in forming and sustaining social relationships, which demands frequent communication and emotional involvement. They also tend to use smartphones to handle emotionally intense matters, resulting in more frequent use and checking of their devices compared to males.⁵⁵ In addition, females are less inclined to participate in outdoor activities and events, preferring instead to build relationships with friends on social media.⁵⁶ They tend to use their smartphones more frequently in public places to avoid feelings of loneliness. Conversely, males typically use their phones more for professional and technological purposes rather than social interactions. Furthermore, females are more prone to anxiety if they cannot keep up with social media updates.⁵⁷

Smartphone use is widespread across all age groups. Behera⁵⁸ found similar results to this study, indicating that 90% of adolescents with MSU were in the 16 – 17 age group. Positive correlations suggest that MSU increases with an increase in student age.⁵⁹ Older adolescents exhibit a higher level of smartphone addiction compared to younger ones.^{60–62} Age is the most effective predictor variable, determining that the level of MSU increases with age.⁶³ These results may be due to social media policies and guidelines. Accessibility to major platforms like Facebook, Instagram, and TikTok requires users to meet certain age requirements. In addition, teenagers often seek independence, and their parents may struggle to enforce rules and restrictions. In contrast, a study conducted in Switzerland found a higher prevalence of smartphone addiction among younger adolescents.⁶⁴

The Department of Statistics in Jordan in 2010 estimated that the poverty line for Jordanian families was JD 336 (\$516 USD) per month.⁶⁵ In this study, MSU was more prevalent

among adolescents from families with a monthly income of 301 – 500 JD. In this context, lower economic conditions might heighten the risk of MSU.⁵⁷ Low-income families often face challenges in making ends meet and may spend more time away from home. Consequently, parents in these families might not have sufficient opportunity to oversee their children's smartphone usage, increasing the risk of MSU among low-income youth.⁶⁶ Smartphone addiction is more prevalent among individuals with low income and socioeconomic status, as it provides an escape from their financial difficulties.⁶⁷ In addition, limited family resources may prevent adolescents from participating in outdoor activities, leading them to spend a significant amount of time on their smartphones. However, this finding contrasts with studies indicating that MSU is more common among individuals with high income, where greater financial resources and parental permissiveness allow for increased smartphone access and usage.^{53,68} Furthermore, in this study, an analysis of the mean differences between MSU and GPA, an indicator of academic performance, revealed a significant relationship. Specifically, students with higher levels of MSU tended to have lower GPAs. This finding underscores a potential inverse relationship, suggesting that increased smartphone use may negatively impact academic performance. A meta-analysis study revealed that MSU detrimentally affects students' learning and overall academic performance by impairing the skills and cognitive abilities necessary for academic success.⁶⁹ When students spend excessive time on their phones, neglecting daily responsibilities, engaging in extensive smartphone use, or excessive texting, their GPAs tend to decline.⁶⁹ Excessive smartphone use distracts students and impedes learning, thereby decreasing academic performance.⁷⁰

Sleep quality also emerged as a significant factor influencing MSU among adolescents in this study. Participants reporting poor sleep quality had significantly higher MSU scores than those with moderate or good sleep. This finding is consistent with prior research demonstrating that

excessive smartphone use, particularly during nighttime hours, disrupts sleep patterns and impairs restfulness.¹⁰ Disrupted sleep may, in turn, increase emotional vulnerability and reliance on smartphones as a coping mechanism, creating a negative feedback loop between poor sleep and problematic use. These results highlight the importance of incorporating sleep hygiene education into intervention programs aimed at reducing MSU among adolescents.

Overall, these findings underscore the intricate interplay of sociodemographic factors in shaping MSU behaviors among adolescents. The study highlights how age, gender, economic conditions, academic performance, and sleep quality are all interconnected in influencing the extent and impact of MSU. These results emphasize the need for targeted interventions that consider these sociodemographic factors. Tailored strategies could include educational programs for parents and adolescents about the risks of excessive smartphone use, policies to regulate smartphone usage in schools, and community initiatives to provide alternative recreational activities. Further investigation into contextual influences on MSU behavior is essential to develop a comprehensive understanding of how different environments and circumstances affect smartphone use among adolescents. This research could inform the development of more effective intervention strategies that are sensitive to the specific needs and challenges faced by various demographic groups.

4.2. THE ROLE OF COGNITIVE EMOTION REGULATION STRATEGIES IN SHAPING MSU PATTERNS

This study found that students with high MSU reported significantly higher maladaptive cognitive regulation scores than those with low MSU. Maladaptive cognitive strategies such as rumination, catastrophizing, and self-blame⁷¹ are unhelpful thought patterns that may contribute to problematic smartphone use as a coping mechanism for negative emotions. These findings align with prior research indicating that poor emotion regulation and reliance on negative regulation strategies are associated with MSU.⁷²

Furthermore, maladaptive emotion regulation mediates the relationship between social anxiety and MSU, suggesting that smartphones serve as external emotion regulation tools, reinforcing problematic usage.⁷³ Similar patterns have been observed among Spanish adolescents, where difficulties in regulating emotions correlate with compulsive behaviors and MSU as a coping strategy.¹⁵ Problematic users scored higher in all maladaptive cognitive emotion regulation strategies – including self-blame, rumination, blaming others, and catastrophizing – compared to non-problematic users.¹⁵

Adolescents who rely on maladaptive coping strategies may use smartphones to escape negative emotions or prolong positive ones when alternative strategies are lacking.^{74,75} Deficits in self-regulation contribute to MSU, as emotions act as both warning signals and behavior reinforcers. Effective cognitive emotion regulation is essential for adaptive functioning and emotional resilience.⁷⁶

A meta-analysis confirmed that habitual use of maladaptive emotion regulation strategies is strongly associated with depressive and anxiety symptoms.⁷⁷ While adaptive strategies correlate with fewer symptoms, maladaptive strategies are linked to more frequent symptoms, mediating the psychological impact of MSU. In addition, cognitive emotion regulation strategies influence the extent to which

smartphone use affects mental health and daily functioning, particularly through emotion suppression and cognitive reappraisal.⁷⁸

Arrivillaga *et al.*⁷⁹ found that MSU is significantly associated with depression and anxiety, driven by maladaptive cognitive and emotional processes, such as repetitive negative thinking, susceptibility to boredom, and fear of missing out.⁸⁰ Rumination, a key component of maladaptive cognitive emotion regulation, mediates the relationship between MSU severity and psychological maladjustment.^{81,82} Psychological flexibility and self-compassion, when mediated by cognitive emotion regulation, can predict MSU risk, highlighting their crucial role in mitigating problematic smartphone use.⁸³ This study supports findings from prior research, demonstrating that cognitive emotion regulation strategies significantly impact the extent to which smartphone use affects mental health and daily functioning, underscoring the need for targeted interventions.

5. CONCLUSION

Schools, parents, and community organizations should collaborate to develop educational initiatives that address the risks associated with excessive smartphone use. These programs should focus on promoting healthy digital habits, enhancing self-regulation skills, and educating students about the potential negative impacts of MSU on their academic performance and mental health. In addition, interventions should be tailored to address demographic differences, such as age and gender, to effectively reach and engage diverse groups of adolescents. The findings also highlight the need for incorporating emotional regulation skills into interventions aimed at reducing MSU. Programs should focus on helping adolescents develop adaptive coping strategies and improve their emotional resilience.

ACKNOWLEDGMENTS

None.

FUNDING

This study was funded by the University of Jordan Deanship of Scientific Research (Grant number 18/2023).

CONFLICT OF INTEREST

The authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Conceptualization: Fatimah S. Tarawneh, Latefa Ali Dardas

Data curation: Fatimah S. Tarawneh, Latefa Ali Dardas

Methodology: Fatimah S. Tarawneh, Latefa Ali Dardas

Resources: Fatimah S. Tarawneh

Writing – original draft: All authors

Writing – review & editing: All authors

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Approval for ethical clearance was obtained from the University of Jordan's Deanship of Scientific Research Institutional Review Board Committee (295), followed by

formal authorization from the Ministry of Education to conduct data collection. Parental consent and child assent to participate were obtained following the guidelines of the American Academy of Pediatrics (1976).

CONSENT FOR PUBLICATION

All participants and their legal guardians provided written informed consent for the anonymized data to be used for publication purposes. Consent forms clearly stated that the data collected would remain confidential and would

be used strictly for research and academic dissemination. All names were coded for data analysis to maintain anonymity.

DATA AVAILABILITY STATEMENT

Data will be made available upon request to the corresponding author.

Submitted: 10 February 2025; Accepted: 06 March 2025;
Published: 04 August 2025

REFERENCES

1. Sangeetha S, Inturi R, Srikanth A, Hariprasath T, Abishavarthana P, Samyuktha S. *On-Device Application Categorization using Application Resources*. United States: IEEE; 2023. p. 1-6.
2. Jay A. *Number of Smartphone and Mobile Phone Users Worldwide in 2022/2023: Demographics, Statistics, Predictions*. *FinancesOnline*. Available from: <https://financesonline.com/number-of/2023> [Last accessed on 2023 Jan 15].
3. Panova T, Carbonell X. Is smartphone addiction really an addiction? *J Behav Addict*. 2018;7(2):252-259. doi: 10.1556/2006.7.2018.49
4. Fischer-Grote L, Kothgassner OD, Felnhöfer A. Risk factors for problematic smartphone use in children and adolescents: A review of existing literature. *Neuropsychiatr*. 2019;33(4):179-190. doi: 10.1007/s40211-019-00319-8
5. Rahmillah FI, Tariq A, King M, Oviedo-Trespalacios O. Is distraction on the road associated with maladaptive mobile phone use? A systematic review. *Accid Anal Prev*. 2023;181:106900. doi: 10.1016/j.aap.2022.106900
6. Körmendi A. Smartphone usage among adolescents. *Psychiatr Hung*. 2015;30(3):297-302.
7. Wacks Y, Weinstein AM. Excessive smartphone use is associated with health problems in adolescents and young adults. *Front Psychiatry*. 2021;12:669042. doi: 10.3389/fpsy.2021.669042
8. Kim HJ, Min JY, Min KB, Lee TJ, Yoo S. Relationship among family environment, self-control, friendship quality, and adolescents' smartphone addiction in South Korea: Findings from nationwide data. *PLoS One*. 2018;13(2):e0190896. doi: 10.1371/journal.pone.0190896
9. Lee H, Kim J. A structural equation model on Korean adolescents' excessive use of smartphones. *Asian Nurs Res (Korean Soc Nurs Sci)*. 2018. doi: 10.1016/j.anr.2018.03.002
10. Kim SY, Han S, Park EJ, et al. The relationship between smartphone overuse and sleep in younger children: A prospective cohort study. *J Clin Sleep Med*. 2020;16(7):1133-1139. doi: 10.5664/jcsm.8446
11. Zhuang L, Wang L, Xu D, Wang Z, Liang R. Association between excessive smartphone use and cervical disc degeneration in young patients suffering from chronic neck pain. *J Orthop Sci*. 2021;26(1):110-115. doi: 10.1016/j.jos.2020.02.009
12. Tymofiyeva O, Yuan JP, Kidambi R, et al. Neural correlates of smartphone dependence in adolescents. *Front Hum Neurosci*. 2020;14:564629. doi: 10.3389/fnhum.2020.564629
13. Liu RD, Hong W, Ding Y, et al. Psychological distress and problematic mobile phone use among adolescents: The mediating role of maladaptive cognitions and the moderating role of effortful control. *Front Psychol*. 2019;10:1589. doi: 10.3389/fpsyg.2019.01589
14. Wanqing H, Fenqing L, Solodukho A. Smartphone addiction and cross-cultural adjustment among overseas Chinese students: The role of emotion regulation beliefs and strategies. *Front Psychol*. 2022;13:1009347. doi: 10.3389/fpsyg.2022.1009347
15. Extremera N, Quintana-Orts C, Sánchez-Álvarez N, Rey L. The role of cognitive emotion regulation strategies on problematic smartphone use: Comparison between problematic and non-problematic adolescent users. *Int J Environ Res Public Health*. 2019;16(17):3142. doi: 10.3390/ijerph16173142
16. Vink M, Gladwin TE, Geeraerts S, et al. Towards an integrated account of the development of self-regulation from a neurocognitive perspective: A framework for current and future longitudinal multi-modal investigations. *Dev Cogn Neurosci*. 2020;45:100829. doi: 10.1016/j.dcn.2020.100829
17. Giordano C, Coco GL, Salerno L, Di Blasi M. The role of emotion dysregulation in adolescents' problematic smartphone use: A study on adolescent/parents triads. *Comput Hum Behav*. 2021;115:106632. doi: 10.1016/j.chb.2020.106632
18. Unicef J. *Youth a Successful Transition to Adulthood for Every Child*. United States: Unicef; 2020.
19. WHO. *Country Cooperation Strategy for WHO and Jordan 2021-2025*. Geneva: WHO; 2021.
20. Al-Krenawi A. *Syrian Refugees in Jordan and Mental Health: Challenges and Culturally Sensitive Practice*. Netherlands: Brill; 2024.
21. Abu-Taieh EM, AlHadid I, Kaabneh K, et al. Predictors of smartphone addiction and social isolation among Jordanian children and adolescents using SEM and ML. *Big Data Cognit Comput*. 2022;6(3):92. doi: 10.3390/bdcc6030092
22. Atoum M, Al-Rawashdeh S, Atoum D, Atoum H,

- Atoum R. Electronic devices use association with psychological distress and sleep among adolescents. *Electron J Gen Med*. 2021;18(6):em327. doi: 10.29333/ejgm/11314
23. Statistics Do. *Jordanian Statistical Yearbook*; 2023. Available from: https://dosweb.dos.gov.jo/ar/product/statistical_year_book2023 [Last accessed on 2023 Jan 15].
24. Nanjundeswaraswamy T, Divakar S. Determination of sample size and sampling methods in applied research. *Proc Eng Sci*. 2021;3(1):25-32. doi: 10.24874/PES03.01.003
25. Lin YH, Chang LR, Lee YH, Tseng HW, Kuo TB, Chen SH. Development and validation of the smartphone addiction inventory (SPAI). *PLoS One*. 2014;9(6):e98312. doi: 10.1371/journal.pone.0098312
26. Khoury JM, De Freitas AAC, Roque MAV, Albuquerque MR, Das Neves MCL, Garcia FD. Assessment of the accuracy of a new tool for the screening of smartphone addiction. *PLoS One*. 2017;12(5):e0176924. doi: 10.1371/journal.pone.0176924
27. Pavia L, Cavani P, Di Blasi M, Giordano C. Smartphone addiction inventory (SPAI): Psychometric properties and confirmatory factor analysis. *Comput Hum Behav*. 2016;63:170-178. doi: 10.1016/j.chb.2016.05.039
28. Simó-Sanz C, Ballestar-Tarín ML, Martínez-Sabater A. Smartphone addiction inventory (SPAI): Translation, adaptation and validation of the tool in Spanish adult population. *PLoS One*. 2018;13(10):e0205389. doi: 10.1371/journal.pone.0205389
29. Andrade ALM, Spritzer DT, Scatena A, et al. Psychometric properties of the smartphone addiction inventory-short form (SPAI-SF) in Brazilian adolescents. *Psychiatr Res*. 2023;319:115001. doi: 10.1016/j.psychres.2022.115001
30. Chamizo Nieto MT, Rey L, Sánchez Álvarez N. Validation of the Spanish version of the cognitive emotion regulation questionnaire in adolescents. *Psicothema*. 2020;32:153-159. doi: 10.7334/psicothema2019.156
31. Garnefski N, Kraaij V. The cognitive emotion regulation questionnaire. *Eur J Psychol Assess*. 2007;23(3):141-149. doi: 10.1027/1015-5759.23.3.141
32. Ouerchefani R, Ouerchefani N, Rejeb MR, Le Gall D. Cognitive emotion regulation questionnaire: Psychometric properties of the Tunisian version. *Encephale*. 2021;47(5):406-412. doi: 10.1016/j.encep.2020.11.007
33. Lovibond S. Manual for the depression anxiety stress scales. *Psychology Foundation of Australia*. Australia: NovoPsych; 1995.
34. Parkitny L, McAuley J. The depression anxiety stress scale (DASS). *J Physiother*. 2010;56(3):204. doi: 10.1016/s1836-9553(10)70030-8
35. Hussain NM, AlMansouri DH, AlGhareeb M, et al. Translating and validating the hoarding rating scale-self report into Arabic. *BMC Psychol*. 2023;11(1):233. doi: 10.1186/s40359-023-01277-1
36. Permission P. Informed consent, parental permission, and assent in pediatric practice. Committee on bioethics, American academy of pediatrics. *Pediatrics*. 1995;95(2):314-317.
37. Kim Y, Dhammasaccakarn W, Laeheem K, Rinthaisong I. The impact of family functioning factors on smartphone addiction and phubbing among muslim adolescents in Thailand. *Children (Basel)*. 2024;11(5):522. doi: 10.3390/children11050522
38. Davey S, Davey A. Assessment of smartphone addiction in Indian adolescents: A mixed method study by systematic-review and meta-analysis approach. *Int J Prev Med*. 2014;5(12):1500-1511.
39. Fatkuriyah L, Sun-Mi C. The relationship among parenting style, self-regulation, and smartphone addiction proneness in Indonesian junior high school students. *IJNP Indones J Nurs Pract*. 2021;5(1):51-59. doi: 10.18196/ijnp.v5i1.11186
40. Lee C, Lee SJ. Prevalence and predictors of smartphone addiction proneness among Korean adolescents. *Child Youth Serv Rev*. 2017;77:10-17. doi: 10.1016/j.childyouth.2017.04.002
41. Buctot DB, Kim N, Kim JJ. Factors associated with smartphone addiction prevalence and its predictive capacity for health-related quality of life among Filipino adolescents. *Child Youth Serv Rev*. 2020;110:104758. doi: 10.1016/j.childyouth.2020.104758
42. Esmaeilpour F, Letafatkar A, Baker JS, et al. Reliability and construct validity of the smartphone addiction scale short version (SAS-SV) in Iranian students. *J Public Health*. 2021;31:345-353. doi: 10.1007/s10389-021-01529-7
43. Çağan Ö, Koca B. Evaluation of high school students' smartphone addiction and insomnia level. *Evaluation*. 2020;7(1):45-51. doi: 10.4274/jtsm.galenos.2020.84755

44. Al-Amri A, Abdulaziz S, Bashir S, Ahsan M, Abualait T. Effects of smartphone addiction on cognitive function and physical activity in middle-school children: A cross-sectional study. *Front Psychol.* 2023;14:1182749. doi: 10.3389/fpsyg.2023.1182749
45. Sadiq S, Anasse K, Slimani N. Scrolling through sleep: Moroccan high school students smartphone uses and sleep quality. *Int J Innov Res Sci Stud.* 2023;6(3):702-709. doi: 10.53894/ijirss.v6i3.1795
46. Louraghi I, Ahami A, Khadmaoui A, Mammad K, Lamrani AC. Evaluation of the nomophobia's prevalence and its impact on school performance among adolescents in Morocco. *Prob Psychol Cent.* 2018;12(2):84-94. doi: 10.33225/ppc/18.12.84
47. Fekih-Romdhane F, Malaeb D, Sarray El Dine A, Obeid S, Hallit S. The relationship between smartphone addiction and aggression among lebanese adolescents: The indirect effect of cognitive function. *BMC Pediatr.* 2022;22(1):735. doi: 10.1186/s12887-022-03808-y
48. Buabbas AJ, Al-Mass MA, Al-Tawari BA, Buabbas MA. The detrimental impacts of smart technology device overuse among school students in Kuwait: A cross-sectional survey. *BMC Pediatr.* 2020;20:1-12. doi: 10.1186/s12887-020-02417-x
49. Alabdallat Y, Albakri KA, Al-Hanaqtah BM, Al-Dajani MH, Saleh OM, Harvey H. The association between smartphone addiction, depression and anxiety among medical students in Jordan. *Jordan Med J.* 2023;57(1):60-61. doi: 10.35516/jmj.v57i1.1102
50. Albursan IS, Al Qudah MF, Dutton E, et al. National, sex and academic discipline difference in smartphone addiction: A study of students in Jordan, Saudi Arabia, Yemen and Sudan. *Community Ment Health J.* 2019;55:825-830. doi: 10.1007/s10597-019-00368-x
51. De-Sola Gutiérrez J, Rodríguez De Fonseca F, Rubio G. Cell-phone addiction: A review. *Front Psychiatry.* 2016;7:175. doi: 10.3389/fpsyg.2016.00175
52. Yang SY, Chen MD, Huang YC, Lin CY, Chang JH. Association between smartphone use and musculoskeletal discomfort in adolescent students. *J Community Health.* 2017;42:423-430. doi: 10.1007/s10900-016-0271-x
53. Jahng KE. Maternal abusive parenting and young South Korean adolescents' problematic smartphone use: The moderating effects of time spent hanging out with peers and trusting peer relationships. *Child Youth Serv Rev.* 2019;98:96-104. doi: 10.1016/j.childyouth.2018.12.028
54. Chen B, Liu F, Ding S, Ying X, Wang L, Wen Y. Gender differences in factors associated with smartphone addiction: A cross-sectional study among medical college students. *BMC Psychiatry.* 2017;17:341. doi: 10.1186/s12888-017-1503-z
55. Yang SY, Lin CY, Huang YC, Chang JH. Gender differences in the association of smartphone use with the vitality and mental health of adolescent students. *J Am Coll Health.* 2018;66(7):693-701. doi: 10.1080/07448481.2018.1454930
56. Slaih MA, Khader YS, Amarneh BH, Alyahya MS, Al-Adwan NT. Patterns of smartphone use among university students in Jordan. *Arab J Psychiatr.* 2019;30(1):54-61. doi: 10.12816/0052936
57. Mallya NV, Sunil Kumar DR, Mashal S. A study to evaluate the behavioral dimensions of nomophobia and attitude toward smartphone usage among medical students in Bengaluru. *Nat J Physiol Pharm Pharmacol.* 2018;8(11):1553-1553. doi: 10.5455/njppp.2018.8.0827603092018
58. Behera D. *A Study to Assess the Prevalence of Mobile Phone Addiction and its Impact Among Adolescents in Selected Pre-University Colleges at Bangalore with View to Prepare Information Booklet.* India: Rajiv Gandhi University of Health Sciences; 2018.
59. Gezgin DM. Understanding patterns for smartphone addiction: Age, sleep duration, social network use and fear of missing out. *Cypr J Educ Sci.* 2018;13(2):409-421.
60. Bhandari DJ, Pandya YP, Sharma DB. Smartphone use and its addiction among adolescents in the age group of 16-19 years. *Indian J Community Med.* 2021;46(1):88-92. doi: 10.4103/ijcm.IJCM_263_20
61. Fırat S, Gül H, Sertçelik M, Gül A, Gürel Y, Kılıç BG. The relationship between problematic smartphone use and psychiatric symptoms among adolescents who applied to psychiatry clinics. *Psychiatry Res.* 2018;270:97-103. doi: 10.1016/j.psychres.2018.09.015
62. Lee EJ, Ogbolu Y. Does parental control work with smartphone addiction?: A cross-sectional study of children in South Korea. *J Addict Nurs.* 2018;29(2):128-138. doi: 10.1097/JAN.0000000000000222
63. Durak HY. Investigation of nomophobia and smartphone addiction predictors among adolescents in Turkey: Demographic variables and academic performance. *Soc Sci J.* 2019;56(4):492-517. doi: 10.1016/j.soscij.2018.09.003
64. Haug S, Castro RP, Kwon M, Filler A,

- Kowatsch T, Schaub MP. Smartphone use and smartphone addiction among young people in Switzerland. *J Behav Addict*. 2015;4(4):299-307. doi: 10.1556/2006.4.2015.037
65. Statistics Do. *Report on the Situation of Poverty in Jordan*. Available from: https://dosweb.dos.gov.jo/databank/analytical_reports/economic/poverty_report/poverty_report_2010.pdf [Last accessed on 2023 Jan 01].
66. Lee J, Lim H, Allen J, Choi G, Jung J. Smartphone addiction and depression among low-income boys since COVID-19: The moderating effect of being an only child. *Healthcare (Basel)*. 2021;9:1350. doi: 10.3390/healthcare9101350
67. Khalily MT, Loona MI, Bhatti MM, Ahmad I, Saleem T. Smartphone addiction and its associated factors among students in twin cities of Pakistan. *J Pak Med Assoc*. 2020;70(8):1357-1362. doi: 10.5455/JPMA.23054
68. Sarman A, Çiftci N. Relationship between smartphone addiction, loneliness, and depression in adolescents: A correlational structural equation modeling study. *J Pediatr Nurs*. 2024;76:150-159. doi: 10.1016/j.pedn.2024.02.019
69. Sunday OJ, Adesope OO, Maarhuis PL. The effects of smartphone addiction on learning: A meta-analysis. *Comput Hum Behav Rep*. 2021;4:100114. doi: 10.1016/j.chbr.2021.100114
70. Mahsud M, Khalaf AJM, Mahsud Z, Afzal A, Afzal F. Addiction to smartphones leading to distraction in the classrooms: Effect of different cultures. *J Stat Manag Syst*. 2021;24(4):741-754. doi: 10.1080/09720510.2020.1834660
71. Garnefski N, Kraaij V, Spinhoven P. Negative life events, cognitive emotion regulation and emotional problems. *Person Individ Differ*. 2001;30(8):1311-1327. doi: 10.1016/S0191-8869(00)00113-6
72. Van Deursen AJ, Bolle CL, Hegner SM, Kommers PA. Modeling habitual and addictive smartphone behavior: The role of smartphone usage types, emotional intelligence, social stress, self-regulation, age, and gender. *Comput Hum Behav* 2015;45:411-420. doi: 10.1016/j.chb.2014.12.039
73. Zsido AN, Arato N, Lang A, Labadi B, Stecina D, Bandi SA. The role of maladaptive cognitive emotion regulation strategies and social anxiety in problematic smartphone and social media use. *Person Individ Differ*. 2021;173:110647. doi: 10.1016/j.paid.2021.110647
74. Casale S, Caplan SE, Fioravanti G. Positive metacognitions about Internet use: The mediating role in the relationship between emotional dysregulation and problematic use. *Addict Behav*. 2016;59:84-88. doi: 10.1016/j.addbeh.2016.03.014
75. Estévez A, Urbiola I, Iruarizaga I, Onaindia J, Jauregui P. Emotional dependency in dating relationships and psychological consequences of internet and mobile abuse. *Anal Psicol*. 2017;33(2):260-268. doi: 10.6018/analesps.33.2.255111
76. Maleki L, Kazemi A, Alimehdi M. The relationship between cognitive emotional regulation and mindfulness with problematic use of cell-phones among medical students. *Family Health*. 2023;12(4):114-127.
77. Schäfer JÖ, Naumann E, Holmes EA, Tuschen-Caffier B, Samson AC. Emotion regulation strategies in depressive and anxiety symptoms in youth: A meta-analytic review. *J Youth Adolesc*. 2017;46:261-276. doi: 10.1007/s10964-016-0585-0
78. Fortes AB, Broilo PL, Lisboa CSM. Smartphone use and psychological well-being: The moderating role of emotion regulation. *Trends Psychol*. 2021;29:189-203. doi: 10.1007/s43076-020-00051-1
79. Arrivillaga C, Elhai JD, Rey L, Extremera N. Depressive symptomatology is associated with problematic smartphone use severity in adolescents: The mediating role of cognitive emotion regulation strategies. *Cyberpsychol J Psychosoc Res Cybers*. 2023;17(3):2. doi: 10.5817/CP2023-3-2
80. Elhai JD, Yang H, Montag C. Cognitive-and emotion-related dysfunctional coping processes: Transdiagnostic mechanisms explaining depression and anxiety's relations with problematic smartphone use. *Curr Addict Rep*. 2019;6:410-417. doi: 10.1007/s40429-019-00260-4
81. Arrivillaga C, Rey L, Extremera N. Psychological distress, rumination and problematic smartphone use among Spanish adolescents: An emotional intelligence-based conditional process analysis. *J Affect Disord*. 2022;296:1-8. doi: 10.1016/j.jad.2021.09.021
82. Vally Z, Alghraibeh AM, Elhai JD. Severity of depression and anxiety in relation to problematic smartphone use in the United Arab Emirates: The mediational roles of rumination and fear of missing out. *Hum Behav Emerg Technol*. 2021;3(3):423-431. doi: 10.1002/hbe2.259
83. Zavardeh SZ, Salimi F, Amani O. The mediating role of cognitive emotion regulation in the relationship between psychological flexibility and self-compassion with problematic use of smartphone in none-clinical population. *J Psychol N Ideas*. 2024;20(24):1-12.