

RESEARCH ARTICLE

Gradient hydroxyapatite nanoparticles with spatial distribution facilitate the healing of tendon-to-bone interface

Supplementary File
Table S1. Primer sequences and product size used for polymerase chain reaction (PCR) analysis

Gene	Primer	Sequence (5'-3')	Size (mer)
GAPDH	Forward	CAAGTTCAACGGCACAGTCAAG	22
	Reverse	CGCCAGTAGACTCCACGACATA	22
Scx	Forward	GTTGAGCAAAGACCGTGACAGA	22
	Reverse	ATAGCCCGCCTGTCCATCTC	20
Tnmd	Forward	ACACTTTCTACAGCAATGGCGA	22
	Reverse	GACTTCCAATGTTTCATCGGTGC	23
Acan	Forward	CCTTGGACACTTTCACATGCTTAT	24
	Reverse	TCCACTGGTAGTCTTGAGCATT	23
Col II	Forward	GAAGAGCGGAGACTACTGGATTG	23
	Reverse	TCTCCAAACCAGATGTGCTTCTT	23
Bmp2	Forward	GCGGTCTCCTAAAGGTCCTC	20
	Reverse	AGCCTCAACTCAAACCTCGCT	20
Runx2	Forward	CTGGCCTTCCTCTCTCAGTAAGA	23
	Reverse	TGTAAGTGAAGGTGGCTGGATAG	23