

# Integration of topological indices and QSPR modeling with VIKOR-based multi-criteria decision analysis for evaluating eye drug candidates

Muhammad Kamran<sup>1,2,3†</sup>, Shereen Iqbal<sup>4†</sup>, Hifza Iqbal<sup>4\*</sup>, Muhammad Akhtar Tarar<sup>5†</sup>, Muhammad Farman<sup>1,6</sup>, Evren Hincal<sup>1,7</sup> and Mohamad Hafez<sup>3,8</sup>

<sup>1</sup>Mathematics Research Center, Department of Mathematics Near East University, Mersin, Turkey

<sup>2</sup>International Center for Interdisciplinary Research in Sciences, The University of Lahore, Lahore, Pakistan

<sup>3</sup>Department of Mathematics, Faculty of Engineering and Quantity Surviving, INTI International University Colleges, Nilai, Negeri Sembilan, Malaysia

<sup>4</sup>Department of Mathematics and Statistics, The University of Lahore, Lahore, Punjab, Pakistan

<sup>5</sup>Department of Civil Engineering, The University of Lahore, Lahore, Punjab, Pakistan

<sup>6</sup>Department of Computer Engineering, Biruni University, Istanbul, Turkey

<sup>7</sup>Research Center of Applied Mathematics, Khazar University, Baku, Azerbaijan

<sup>8</sup>Department of Management, Faculty of Management, Shinawatra University, Pathum Thani, Thailand  
[kamrankfueit@gmail.com](mailto:kamrankfueit@gmail.com); [shereeniqbal321@gmail.com](mailto:shereeniqbal321@gmail.com); [iqbalhifza3@gmail.com](mailto:iqbalhifza3@gmail.com); [akhtartarar2000@yahoo.com](mailto:akhtartarar2000@yahoo.com);  
[farmanlink@gmail.com](mailto:farmanlink@gmail.com); [evren.hincal@neu.edu.tr](mailto:evren.hincal@neu.edu.tr); [mohdahmed.hafez@newinti.edu.my](mailto:mohdahmed.hafez@newinti.edu.my)

## ARTICLE INFO

### Article History:

Received: January 19, 2026

Revised: March 12, 2026

Accepted: March 19, 2026

Published Online: May 13, 2026

### Keywords:

Graph theory

Multi-criteria decision-making

Multiple linear regression

Public health

Quantitative structure–property relationship ranking

VIKOR

AMS Classification 2010:

26A33; 34A08; 35H15;

34K50 47H10; 60H10

## ABSTRACT

Topological indices encode molecular structure into numerical descriptors, enabling prediction of physicochemical properties without extensive experimentation. This study evaluates the implementation of graph-theoretical topological indices in combination with multi-criteria decision-making methods to compute physicochemical properties and ranking of eye drugs. A set of classical topological indices, consisting of  $NSK_1$ ,  $NSK_2$ ,  $M_1$ ,  $M_2$ ,  $NM_1$ ,  $NM_2$ ,  $NSK$ ,  $NF$ , and  $NH$ , are computed using an edge-partition approach to describe the structural aspects of 12 selected drugs. Multiple linear regression models were developed to develop quantitative structure–property relationship models for important physicochemical properties, including molar volume, polarizability, molar weight, and molar refraction. The multiple linear regression models showed reliable predictive output, with statistical indicators and correlation coefficients higher than 0.89, confirming the value of the selected indices. The VIKOR technique enables the ranking of drugs based on their estimated physicochemical properties. The combined application of quantitative structure–property relationship and multi-criteria decision-making framework enhances robustness in discriminating structurally similar eye drugs. This combination offers a systematic framework for prioritizing compounds in pharmacological research and drug optimization. Such approach also supports public health by facilitating more efficient drug development and therapeutic decision-making for eye diseases.



\*Corresponding Author

†These authors contributed equally to this work.

## 1. Introduction

A wide range of eye disorders encompasses conditions that effect vision and eye health, ranging from mild, manageable problems to serious diseases that can impair sight.<sup>1</sup> Notably, progressive clouding of the ocular lens, known as cataract, is the most prevalent condition, resulting in limited sight. The therapeutic approach involves implantation of a synthetic intraocular lens<sup>2</sup> after removal of the affected lens. Glaucoma is one of the most serious eye disorder, which is a set of clinical conditions linked to degeneration of optic nerves. This disorder is commonly caused by an increase in pressure within the eye. Patients with glaucoma commonly report complaints such as blurred vision, discomfort in the eyes, restriction of peripheral vision, and a progressive decline in eyesight.<sup>3</sup> Glaucoma is treated clinically by reducing intraocular pressure using pharmacological therapy to prevent progressive optic nerve damage, as well as through laser-based interventions.<sup>4</sup>

Age-related oracular degeneration is recognized as a major cause of vision loss in elderly individuals. This disorder arises from the degeneration of central retina and is commonly demonstrated by blind spots, challenges in reading, and visual distortion<sup>5</sup>. Current treatment techniques, including photo-dynamic therapy, anti-vascular endothelial growth factor injections, and laser interventions, are primarily aimed at delaying disease propagation and preserving remaining vision. Diabetic eye disease, a well-known micro vascular disorder, develops due to retinal blood vessel damage. Symptoms may include progressive eyesight loss, floaters, and blurred vision.<sup>6</sup> Conjunctivitis, commonly known as pink eye, is an inflammatory condition affecting the thin, translucent membrane that covers the sclera and lines the inner surface of the eyelids.<sup>7</sup> Its clinical manifestation include eyelid crusting, itching, redness, discharge, and tearing.<sup>8</sup> Allergic conjunctivitis is treated with supportive techniques such as cold compresses and antihistamine drops, whereas bacterial conjunctivitis is usually cured by antibiotic eye drops.<sup>9</sup>

A collection of eye drugs plays an important role in eye ailments management by using various pharmacological techniques. Carbolic anhydrase modulating agents such as acetazolamide,<sup>10</sup> and brinzolamide<sup>11</sup> lower intraocular pressure proficiently, thereby contributing to glaucoma supervision and reducing aqueous humor secretion. Acetylcysteine<sup>12</sup> acts as an antioxidant for dry eye conditions, thereby improving corneal health in inflammatory keratitis,

while aciclovir<sup>13</sup> and ganciclovir<sup>14</sup> serve as antiviral agents against human cytomegalovirus diseases. Tropicamide<sup>15</sup> and cyclopentolate,<sup>16</sup> both parasympathetic drugs, are commonly used for indicative cyclopedia and mydriasis. xylometazoline,<sup>17</sup> and apraclonidine,<sup>18</sup> act to alleviate eye congestion. Adrenergic receptor agonists decrease intraocular pressure by decreasing aqueous humor production and inducing vasoconstrictive effects on ocular blood vessels. Bromfenac,<sup>19</sup> a nonsteroidal anti-inflammatory drug, effectively controls postoperative inflammation following ocular surgery. Carteolol,<sup>20</sup> a  $\beta$ -blocker, reduces intraocular pressure with minimal systemic effects, while lodoxamide<sup>21</sup> stabilizes mast cells thereby preventing allergic conjunctivitis. These drugs form a dynamic part of modern pharmacotherapy, targeting infectious, inflammatory, and pressure-related ocular diseases with high therapeutic precision. This is supported by studies highlighting the impact of infection and parasitic diseases on self-rated health, as well as the role of socioeconomic and environmental factors, including housing conditions, in the spread and mitigation of infection diseases.<sup>22,23</sup>

Euler, in 1736 first formulated graph theory by constituting a fundamental branch of discrete mathematics.<sup>24,25</sup> Since its inception, it has found widespread use in technological fields, such as chemistry, physics, biology, and computer science, and in multiple scientific techniques.<sup>26,27</sup> The development of chemical graph theory has emerged through the integration of molecular structures with mathematical frameworks. The main purpose of this field lies in the study of topological indices, which perform as numerical tools strongly correlated with the functional and structural properties of molecules. These indices are widely applied in quantitative structure-activity relationship (QSAR) and quantitative structure-property relationship (QSPR) modeling, where they provide a reliable basis for predicting biological activities and physicochemical properties of chemical entities.<sup>28</sup>

Topological indices act as mathematical tool that efficiently capture the structural properties of chemical compounds.<sup>29</sup> The concept of topological indices was first introduced through the Wiener index in 1947, which played a pioneer role in correlating physicochemical properties with the molecular structure of petroleum.<sup>30</sup> Since then, in chemical graph theory, these indices have become fundamental tools, enabling the analysis and calculation of physicochemical properties. A molecular graph  $G$  is constructed from the structural framework of hydrocarbons. In such graph, the

vertices correspond to atoms other than hydrogen, and this set of collection is depicted as  $V(G)$ . The bonds between these atoms are represented as edges, and set of edges is denoted by  $E(G)$ .<sup>31,32</sup>

The first and second Zagreb indices, originally proposed by Trinajstić and Gutman,<sup>33</sup> are among the earliest and most fundamental topological descriptors in chemical graph theory. These indices have since played a pivotal role in correlating molecular structure with various physicochemical and biological properties.

Mathematically, these indices are expressed as:

$$M_1(G) = \sum_{ab \in E(G)} d(a) + d(b), \quad (1)$$

$$M_2(G) = \sum_{ab \in E(G)} d(a) \times d(b), \quad (2)$$

where  $d(a)$  and  $d(b)$  represent the degrees of vertices  $a$  and  $b$ , respectively.

$$NM_1(G) = \sum_{ab \in E(G)} [S(a) + S(b)], \quad (3)$$

$$NM_2(G) = \sum_{ab \in E(G)} [S(a) \times S(b)], \quad (4)$$

where  $S(a)$  and  $S(b)$  represent the sum of neighborhood degrees of vertices  $a$  and  $b$ , respectively. The Hyper Zagreb index, denoted as  $HM(G)$ ,<sup>34</sup> is mathematically defined as follows:

$$HM(G) = \sum_{ab \in E(G)} (d(a) + d(b))^2 \quad (5)$$

Shirdel et al. proposed a revised formulation of the Zagreb indices.<sup>35</sup> In this modification, the expression for the second Zagreb index of a graph  $G$  is reformulated as follows:

$$ReZG_2(G) = \sum_{ab \in E(G)} \frac{d(a) \times d(b)}{d(a) + d(b)} \quad (6)$$

The square root maximum–minimum degree index  $mMsde(G)$  is defined as:

$$mMsde(G) = \sum_{ab \in E(G)} \sqrt{\frac{\max\{d(a), d(b)\}}{\min\{d(a), d(b)\}}} \quad (7)$$

The neighborhood Sombor–Kulli  $NSK$  index is defined as:

$$NSK(G) = \sum_{ab \in E(G)} \sqrt{S(a)^2 + S(b)^2} \quad (8)$$

The  $SK_1$  index (first Sombor-type index) is described as:

$$SK_1(G) = \sum_{ab \in E(G)} \frac{1}{\sqrt{d(a)^2 + d(b)^2}} \quad (9)$$

The  $SK_2$  index (second Sombor-type index) described as:

$$SK_2(G) = \sum_{ab \in E(G)} \sqrt{d(a)^2 + d(b)^2} \times [d(a) + d(b)] \quad (10)$$

The neighborhood harmonic index is defined as:

$$NH(G) = \sum_{ab \in E(G)} [d(a)^2 + d(b)^2] \quad (11)$$

The neighborhood forgotten index is defined as:


$$NF(G) = \sum_{b \in V(G)} d(b)^3 \quad (12)$$

Topological indices for the compounds depicted in **Figure 1** were derived using an edge partitioning approach. In this method, the indices are computed by systematically counting edges that join vertices with the same degree classification ( $d(a), d(b)$ ).

## 2. Methodology

This study employed a combined computational framework integrating graph-theoretical topological indices, QSPR modeling, and multi-criteria decision-making (MCDM) analysis. Molecular structures of 12 ophthalmic drug candidates were initially retrieved from established pharmacological databases and translated into molecular graphs, where atoms represent vertices and bonds represent edges. A set of classical and newly defined topological descriptors  $M_1, M_2, NM_1, NM_2, NSK, NSK_1, NSK_2, NH, NF, HM, mMsde$ , and  $ReZG_2$  was calculated using an edge-partitioning algorithm. Subsequently, a multiple linear regression (MLR) analysis was performed to develop QSPR models correlating these descriptors with experimental physicochemical parameters, including polarizability, molar volume, molar refraction, and molecular weight. Statistical metrics, including correlation coefficient, coefficient of determination, Fisher's statistic, standard error, and root mean square error were used to evaluate model performance and predictive accuracy. For MCDM, the VIKOR method was used based on their predicted physicochemical behavior to prioritize the studied drugs. The topological indices and weights assigned to these indices served as decision criteria, which enable compromise-based ranking of drugs that identified the most structurally efficient candidates.

**Figure 1** depicts the molecular structures of the 12 drugs for ocular disorders analyzed in this study. These chemical structures serve as the foundation for constructing molecular graphs



**Figure 1.** Molecular structure of (A) acetazolamide, (B) acetylcysteine, (C) aciclovir, (D) tropicamide, (E) xylometazoline, (F) apraclonidine, (G) brinzolamide, (H) bromfenac, (I) carteolol, (J) cyclopentolate, (K) lodoxamide, and (L) ganciclovir

**Table 1.** Physicochemical properties of ophthalmic drugs

Eye drugs	Polarizability	Molar volume	Molar refraction	Molar weight
Acetylcysteine	15.2	126.1	38.3	163.2
Acetazolamide	18.9	110.6	47.7	222.3
Aciclovir	20.8	127.2	52.4	225.2
Apraclonidine	23.5	150	59.2	245.11
Tropicamide	32.6	244.8	82.2	284.35
Bromfenac	31.4	213.5	79.1	334.16
Brinzolamide	35.8	255.4	90.4	383.5
Cartelol	32.3	258.6	81.4	292.3
Iodoxamide	26	174.6	65.5	311.63
Cyclopentolate	32.7	256.5	82.4	291.4
Ganciclovir	23	140.6	57.9	255.23
Xylometazoline	30.5	243.6	76.9	244.37

used to compute topological indices. Each chemical structure reflects a distinct atomic connectivity and arrangement, which influence its physicochemical properties and ranking.

**Table 1** presents the experimentally obtained physicochemical parameters—polarizability, molar volume, molar refraction, and molecular weight—of the selected drugs, sourced from PubChem. These physicochemical parameters provide insight into molecular size, optical properties, and inter molecular interactions of compounds, serving as dependent variables in the QSPR modeling framework.

Multiple linear regression is a most commonly used technique for predicting a target variable based on two or more independent variables.<sup>36</sup> The mathematical form of multiple linear regression model is:

$$Z = a_1I_1 + a_2I_2 + a_3I_3 + \cdots + k,$$

where  $Z$  denotes the target variable,  $a_i$  represents the coefficients of regression corresponding to the input variables  $I_i$ , and  $k$  is known as regression constant. In MLR framework, several modeling approaches have been developed, including

step-wise regression, the standard method, sequential forward selection, and sequential backward approach. In this study, the standard method was employed, where physicochemical properties served as target variables, while topological indices served as input variables. To evaluate model performance, key statistical metrics were employed, including sample size ( $n$ ), correlation coefficient ( $r$ ), standard error, coefficient of determination ( $R^2$ ), Fisher's statistic ( $F$ ), and significance level ( $p$ ). These metrics describe the reliability and predictive capacity of the regression model.

### 3. Results and discussion

This section integrates topological indices with MCDM methods and QSPR gives a systematic approach for quantitative drug validation. The high degree of correlation between the physicochemical properties and computed indices underscores the structural interpretability of these indices. Topological indices such as  $NM_1$ ,  $NH$ , and  $NSK$  effectively captured structural topology that influenced major physicochemical attributes relevant to eye drugs design.

#### Illustrative computation:

Let  $G$  represent molecular graph of acetylcysteine, then

$M_1 = 40$ ,  $M_2 = 41$ ,  $NM_1 = 238$ ,  $NM_2 = 216$ ,  $NSK = 212$ ,  $NSK_1 = 534$ ,  $NSK_2 = 610$ ,  $NH = 620$ ,  $NF = 479$ ,  $mMsde = 13.02$ ,  $HM = 184$ , and  $ReZG_2 = 8.77$ .

Based on **Figure 1**,  $|V(G)| = 9$  and  $|E(G)| = 9$ . Degree-based edge partitions of acetylcysteine are as follow:

$$E_{3,4} = \{e = ab \in E(G) \mid d(a) = 3, d(b) = 4\},$$

$$E_{3,5} = \{e = ab \in E(G) \mid d(a) = 3, d(b) = 5\},$$

$$E_{4,6} = \{e = ab \in E(G) \mid d(a) = 4, d(b) = 6\},$$

$$E_{4,7} = \{e = ab \in E(G) \mid d(a) = 4, d(b) = 7\},$$

$$E_{5,7} = \{e = ab \in E(G) \mid d(a) = 5, d(b) = 7\},$$

$$E_{6,7} = \{e = ab \in E(G) \mid d(a) = 6, d(b) = 7\}.$$

such that  $|E_{3,4}| = 3$ ,  $|E_{3,5}| = 2$ ,  $|E_{4,6}| = 1$ ,  $|E_{4,7}| = 1$ ,  $|E_{5,7}| = 1$ , and  $|E_{6,7}| = 1$ .

The required results were obtained using **Equations 1–12**.

Similarly, the results for the remaining drugs are summarized in **Table 2**.

The predictive performance of the topological indices in modeling molecular activity was validated using the MLR models, which demonstrated satisfactory accuracy in estimating physicochemical properties. The potential

of these models as computational substitutes for preliminary screening is supported by the agreement between predicted and observed values. An additional decision-support approach based on VIKOR method was introduced to integrate multiple physicochemical parameters into a single evaluation model. Evidence-based drug candidate selection was facilitated by the resulting framework, which showed differences between structurally isomorphic molecules. Brinzolamide obtained a lower ranking due to its complex structure and higher molecular weight, while acetylcysteine's ranking at the top can be attributed to its balanced molecular characteristics and favorable physicochemical profile.

Overall, the combined QSPR-VIKOR approach shows a successful strategy for selection in ophthalmic research and rational drug ranking. It not only provides a reproducible mathematical foundation for early-phase drug screening but also reduces dependency on experimental assays. Future studies may incorporate additional graph descriptors, larger molecular datasets, and advanced machine-learning approaches for generalization across drug classes and to further enhance predictive accuracy.

**Table 2**, presents the quantitative measures of the computed graph-theoretical indices ( $NSK$ ,  $NSK_1$ ,  $NSK_2$ ,  $M_1$ ,  $M_2$ ,  $NH$ ,  $NF$ ,  $NM_1$ ,  $NM_2$ ,  $mMsde$ ,  $ReZG_2$ , and  $HM$ ) for every drug. The diversity in these values captures variations in molecular topology, which are crucial for correlating structure with properties. **Table 3** presents the correlation coefficients between the topological indices and the measured physicochemical properties. The consistently high correlation values ( $r \geq 0.89$ ) affirm that the chosen descriptors accurately reflect structural influence on the compounds' physicochemical characteristics.

#### 3.1. Regression models

##### (a) Model 1

$$P = -4.3447 + 0.6661(mMsde) - 1.4494(ReZG_2) + 0.3203(NM_1) - 0.0562(NM_2) + 3.3858(NH)$$

$$n = 12, \quad F = 10.1165, \quad p = 0.0069, \quad R = 0.9455, \quad R^2 = 0.894, \quad SE = 30.65, \quad VIF = 4.9809.$$

##### (b) Model 2

$$MV = -36.0271 + 16.7369(M_1) - 9.2227(M_2) -$$

**Table 2.** Topological indices of ophthalmic drugs

Eye drugs	$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mMsde$	$HM$	$ReZG_2$
Acetazolamide	60	70	140	383	70	191.5	392	2.53	802	18.97	322	13.81
Acetylcysteine	40	41	83	195	41.5	97.5	202.25	2.06	419	13.02	184	8.77
Aciclovir	80	93	186	534	93	267	545	3.39	1,112	20.45	388	19.07
Tropicamide	100	114	228	610	114	305	626	4.53	1,286	25.36	470	24.08
Xylometazoline	96	114	216	620	108	310	634.5	3.46	1,298	24.68	496	21.96
Apraclonidine	76	87	174	479	87	239.5	489	3.04	998	19.99	368	17.85
Brinzolamide	106	125	284	886	142	443	910.5	4.56	1,876	28.45	556	23.75
Bromfenac	96	109	224	606	112	303	623	4.11	1,280	24.10	452	23.05
Cartelol	106	118	254	706	127	353	722	4.29	1,478	28.90	520	24.03
Cyclopentolate	104	119	238	676	119	338	697.5	4.38	1,438	27.08	510	24.23
Iodoxamide	90	101	216	590	108	295	605.5	3.86	1,242	25.47	438	20.45
Ganciclovir	90	105	212	609	106	304.5	625	3.65	1,282	23.54	440	21.33

**Table 3.** Correlation values of various indices for different properties of ophthalmic drugs

Property	$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mMsde$	$ReZG_2$	$HM$
Polarizability	0.9275	0.9172	0.9208	0.8891	0.9208	0.8891	0.8909	0.9208	0.8929	0.9116	0.9081	0.9225
Molar volume	0.8333	0.8109	0.8043	0.7659	0.8043	0.7659	0.7683	0.8234	0.7713	0.8274	0.8013	0.9225
Molar refraction	0.9268	0.9167	0.9206	0.8893	0.9206	0.8893	0.8911	0.9202	0.8930	0.9113	0.9072	0.9222
Molar weight	0.8123	0.8051	0.8845	0.8773	0.8845	0.8773	0.8796	0.8582	0.8815	0.8397	0.7882	0.8183

$$3.8381(NM_1) + 1.6575(NSK_1)$$

$$n = 12, F = 5.2851, p = 0.0279, R = 0.8667, R^2 = 0.7512, SE = 36.94, VIF = 2.29516.$$

### (c) Model 3

$$MR = 7.1373 + 1.1012(NSK) - 1.6544(NSK_2) + 0.7702(NF) - 0.3407(mMsde)$$

$$n = 12, F = 14.695, p = 0.001, R = 0.9453, R^2 = 0.8936, SE = 6.77, VIF = 4.9632.$$

### (d) Model 4

$$MW = 127.3426 - 0.2588(HM) - 4.7906(NM_2) + 5.1019(NSK_2)$$

$$n = 12, F = 12.1937, p = 0.002, R = 0.9058, R^2 = 0.8206, SE = 28.83, VIF = 3.0617.$$

Topological indices are not selected randomly; initially, these indices were screened using correlation analysis. Only those indices which show strong correlation coefficients ( $|r| \geq 0.70$ ) with the target physicochemical property were considered for model construction. Subsequently, MLR models were developed using step-wise selection to retain only statistically significant descriptors ( $p < 0.05$ ).

To examine the presence of multicollinearity among the selected descriptors, the variance inflation factor (VIF) was calculated for each independent variable. In the present study, the calculated

VIF values for all selected topological indices, fall within the acceptable range, as shown in **Table 4**. This suggests that the independent variables do not exhibit strong inter correlations. It further indicates that each index contributes to the regression model and that the estimated coefficients of models are reliable. Therefore, the developed models can be considered free from significant multicollinearity effects and statistically stable.

**Table 5** presents the predicted values of the physicochemical properties obtained through MLR models. The close agreement between predicted and observed values as described in **Tables 1** and **5** validates the robustness and generalizability of the developed QSPR models.

**Figure 2** compares the actual and estimated physicochemical parameters graphically. The near-overlapping data trends confirm that the QSPR models effectively capture the relationship between molecular structure and physicochemical properties.

### 3.2. Multi-criteria decision-making technique


Multi-criteria decision-making refers to a systematic approach used to analyze and resolve decision-making and planning problems that incorporate multiple evaluation criteria.<sup>37,38</sup> In many real-world cases, a single optimal solution may not exist; therefore, decision makers must differentiate among alternatives based on their preferences and objectives.<sup>39</sup> The notion of solution in this context can vary; it may imply

**Table 4.** Variation inflation factor values of various indices for different properties of ophthalmic drugs

Property	$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mM_{sde}$	$ReZG_2$	$HM$
Polarizability	7.153	6.297	6.570	4.773	6.570	4.773	4.874	6.570	4.933	5.917	5.701	6.711
Molar volume	3.272	2.919	2.832	2.418	2.831	2.418	2.441	3.105	2.468	3.170	2.794	6.711
Molar refraction	7.090	6.263	6.557	4.781	6.557	4.781	4.855	6.526	4.893	5.898	5.650	6.686
Molecular weight	2.939	2.842	4.594	4.341	4.594	4.341	4.419	3.795	4.485	3.391	2.640	3.026

**Table 5.** Estimated values of physicochemical properties of selected ophthalmic drugs

Eye drugs	Polarizability	Molar volume	Molar refraction	Molar weight
Acetazolamide	20.15	102.69	46.93	209.15
Acetylcysteine	13.02	98.36	36.51	177.41
Tropicamide	31.16	216.73	78.85	277.23
Aciclovir	22.68	173.88	57.39	249.28
Apraclonidine	22.22	156.01	55.78	232.23
Brinzolamide	34.23	187.43	87.75	384.25
Bromfenac	29.89	200.44	77.42	285.74
Carteolol	34.16	209.91	81.02	294.17
Lodoxamide	28.12	176.37	72.23	276.73
Ganciclovir	26.42	173.27	69.24	284.68
Cyclopentolate	31.63	213.84	82.55	315.48
Xylometazoline	26.32	204.12	67.66	265.96

**Figure 2.** Comparison of actual and estimated values of physicochemical properties

selecting the most preferred option from a set of alternatives, identifying a subset of practical choices, or ranking them according to priority. In its broadest sense, MCDM aims to determine all efficient or non-dominated alternatives that best satisfy the established criteria.<sup>36</sup>

The QSPR approach, integrated with the VIKOR method, offers a systematic framework for assessing and prioritizing compounds based on their biological and pharmacological characteristics.<sup>37</sup> MCDM technique, VIKOR,<sup>38</sup> was employed to rank 12 ophthalmic drugs. In this method, the weighting factors were determined through the ratio approach (**Figure 3**), and a comparable procedure was followed for assigning weights to other molecular properties.

### 3.3. VIKOR technique

In order to obtain results that approach the optimal solution, the VIKOR technique was applied. This method considers each investigated drug as an alternative and evaluates them using a set of predefined criteria derived from the QSPR analysis conducted in this study. By combining multiple criteria, VIKOR enables systematic comparison of alternatives and provides a statistically supported ranking of the drugs. Crucially, the method identifies a solution that approximates the optimal result, showing a balanced trade-off between conflicting factors and providing insight for decision-making in drug prioritization. Zeleny<sup>39</sup> proposed the notion of compromise-based solutions within MCDM with practical applications identified in 1998.<sup>40</sup> The following algorithm explained the proposed methodology steps, which are described as follows:

[H] VIKOR algorithm Decision matrix  $f_{ij}$ , weight vector  $w_i$ , compromise parameter  $v \in [0, 1]$   $S_j$ ,  $R_j$ ,  $Q_j$  and final ranking

Initialize parameters

Determine number of alternatives  $m$  and criteria  $n$

$i = 1$   $n$  Determine the best value:

$$f_i^+ = \begin{cases} \max_j f_{ij}, & \text{if criterion } i \text{ is benefit type} \\ \min_j f_{ij}, & \text{if criterion } i \text{ is cost type} \end{cases}$$

Determine the worst value:

$$f_i^- = \begin{cases} \min_j f_{ij}, & \text{if criterion } i \text{ is benefit type} \\ \max_j f_{ij}, & \text{if criterion } i \text{ is cost type} \end{cases}$$

$j = 1$   $m$  Compute the utility measure:

$$S_j = \sum_{i=1}^n w_i \frac{f_i^+ - f_{ij}}{f_i^+ - f_i^-}$$

Compute the regret measure:

$$R_j = \max_i \left[ w_i \frac{f_i^+ - f_{ij}}{f_i^+ - f_i^-} \right]$$

Determine:

$$S^+ = \min_j S_j, \quad S^- = \max_j S_j$$

$$R^+ = \min_j R_j, \quad R^- = \max_j R_j$$

$j = 1$   $m$  Compute the VIKOR index:

$$Q_j = v \frac{S_j - S^+}{S^- - S^+} + (1 - v) \frac{R_j - R^+}{R^- - R^+}$$

Rank alternatives in ascending order of  $Q_j$

Select the best alternative based on minimum  $Q_j$  and check VIKOR acceptance conditions

The relative weight  $w_i$  of each criteria is calculated by dividing its mean by the sum of all criterion means:

$$w_i = \frac{\bar{Y}_i}{\sum_{k=1}^m \bar{Y}_k},$$

where  $m$  is the number of criteria. This ensures:

$$\sum_{i=1}^m w_i = 1$$

**Table 6** presents the results considering polarizability as the evaluation criterion, with relative importance assigned to each topological index. The assign weights are evenly distributed, which shows that, in prediction of polarizability, no single index dominates. However, indices such as  $M_1$ ,  $HM$ , and  $NH$  contributes slightly higher, suggesting that overall cumulative neighborhood effects and molecular connectivity play a remarkable role in determining the susceptibility to polarization of ophthalmic drugs.

The weights described in **Table 7** reflect the sensitivity of molar volume to molecular structure. In defining molecular size, the  $HM$  index gains the highest weight, emphasizing the role of vertex connectivity. Indices such as  $M_1$  and  $mMsde$  also contribute significantly, indicating that variations in degree distribution that affect the volumetric occupation. The indices  $NSK_1$  and  $NM_2$  gained relatively lower weights, implying that extreme neighborhood interactions have a reduced impact on volumetric behavior.

**Table 8**, presents the weight distribution across indices, indicating that molar refractivity depends on a combination of neighborhood interactions, branching, and overall connectivity. Higher weight values associated with  $HM$ ,  $M_1$ , and  $NM_1$  suggest that degree-based interactions and molecular compactness influence optical response. Based on **Table 9**, indices related to



**Table 6.** Weights allocated for polarizability

$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mMsde$	$ReZG_2$	$HM$
0.08501	0.08407	0.08440	0.08150	0.08440	0.08150	0.08166	0.08440	0.08184	0.08357	0.08325	0.08456

**Table 7.** Weights allocated for molar volume

$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mMsde$	$ReZG_2$	$HM$
0.08601	0.08367	0.08298	0.07906	0.08298	0.07906	0.07930	0.08500	0.07963	0.08541	0.08267	0.09523

**Table 8.** Weights allocated for molar refractivity

$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mMsde$	$ReZG_2$	$HM$
0.08497	0.08404	0.08440	0.08153	0.08440	0.08153	0.08170	0.08436	0.08187	0.08354	0.08316	0.08458

**Table 9.** Weights allocated for molecular weight

$M_1$	$M_2$	$NM_1$	$NM_2$	$NSK$	$NSK_1$	$NSK_2$	$NH$	$NF$	$mMsde$	$ReZG_2$	$HM$
0.07956	0.07890	0.08666	0.08596	0.086666	0.08596	0.08618	0.08407	0.08637	0.08228	0.07722	0.08031

cumulative degree effects and neighborhood size, particularly  $NSK$ ,  $NM_1$ , and  $NF$ , gain slightly higher weights, indicating that molecular weight is sensitive to the total structural mass within the graph rather than edge variations. The contribution of  $M_1$  and  $M_2$  suggest that simple degree sums are insufficient to capture molecular weight differences among the studied drugs.

**Figure 3**, illustrates the influence of physical and chemical properties used in VIKOR analysis. The balanced weighting among polarizability, molecular weight, molar volume, and refractivity shows that selected properties contribute significantly to the decision-making process.

**Table 10** summarizes the ranking of drugs used for ocular treatment according to polarizability. Acetylcysteine highlighted as the top-ranked drug, reflecting its favorable balance between ideal and worst-case solutions. Conversely, brinzolamide gains the lowest rank, which can be associated to its variation from the ideal values of polarizability within the evaluated data set.

The results in **Table 11** describe a similar pattern, with acetylcysteine again achieved the top position due to its best adjustment between structural efficiency and molecular size. Acetazolamide and apraclonidine gain relatively strong rankings, while compounds with large molecular structures, such as cyclopentolate and brinzolamide, are placed at the lower end. This result implies that larger molar volume that may decrease suitability under the selected evaluation criteria. **Table 12**, demonstrates that in terms of molar refractivity drugs with moderate structural complexity tend to give better outcome. Because

of stable optical behavior of acetylcysteine consistently gains first rank across properties. Apraclonidine and aciclovir also exhibit favorable rankings, whereas brinzolamide again appears at the lower rank, showing less refractive properties relative to other drugs.

The results presented in **Table 13** highlight the influence of molecular weight on drug ranking. Compounds with lower molecular weight and efficient structural distribution, particularly acetazolamide and acetylcysteine, achieve superior ranks. Molecules with higher molecular weight, including carteolol and brinzolamide, are ranked lower, which may adversely affect pharmaceutical behavior of these drugs.

**Figure 4** displays a visual comparison of overall VIKOR rankings in accordance with all physicochemical criteria. The consistency observed in the drug positions highlights the stability of the applied decision-making technique. Acetylcysteine consistently stands out as the most favorable drug, while brinzolamide occupies the lowest rank. This visualization increase the valuable results and supports the credibility of integrated QSPR-VIKOR framework.

**Table 14** presents the final rankings obtained by evaluating properties into a combined decision matrix. The results in the table show favorable constancy across molar volume, polarizability, molecular weight, and molar refractivity. Acetylcysteine obtains the top rank position across all criterion, indicating a well-balanced physicochemical properties. Acetylcysteine consistently gains first rank because of its simple and compact molecular structure produces balanced degree distributions. Its low molecular weight and limited structural complexity lead to enhance efficacy as

**Table 10.** VIKOR ranking of ophthalmic drugs based on polarizability


Ophthalmic drugs	$S_j$	$R_j$	$Q_j$	Polarizability rank
Acetazolamide	0.296179	0.031369	0.3264008	2
Acetylcysteine	0.000000	0.000000	0.000000	1
Aciclovir	0.464481	0.055467	0.5473903	4
Tropicamide	0.763575	0.083391	0.8552645	9
Xylometazoline	0.740913	0.081028	0.8309866	8
Apraclonidine	0.4620425	0.048896	0.5089676	3
Brinzolamide	0.995244	0.085010	0.9813049	12
Bromfenac	0.7179826	0.076899	0.7959829	7
Cartelol	0.868049	0.088312	0.9360985	11
Cyclopentolate	0.828406	0.083254	0.8875452	10
Iodoxamide	0.669179	0.065523	0.7074112	5
Ganciclovir	0.666756	0.067637	0.7179592	6

**Table 11.** VIKOR ranking of ophthalmic drugs based on molar volume


Ophthalmic drugs	$S_j$	$R_j$	$Q_j$	Molar volume rank
Acetazolamide	0.297575	0.035330	0.3297500	2
Acetylcysteine	0.000000	0.000000	0.0000000	1
Aciclovir	0.535419	0.062463	0.5877380	4
Tropicamide	0.766094	0.083981	0.8133848	9
Xylometazoline	0.710151	0.079877	0.7643475	8
Apraclonidine	0.463231	0.048558	0.4804811	3
Brinzolamide	0.996071	0.095238	0.9863080	12
Bromfenac	0.719843	0.076367	0.7512919	7
Cartelol	0.866037	0.086021	0.8739701	10
Cyclopentolate	0.831103	0.097919	0.9171902	11
Iodoxamide	0.671122	0.066966	0.6788280	6
Ganciclovir	0.668227	0.067169	0.6784134	5

**Table 12.** VIKOR ranking of ophthalmic drugs based on molar refractivity

Ophthalmic drugs	$S_j$	$R_j$	$Q_j$	Molar refractivity rank
Acetazolamide	0.305659	0.039021	0.3835859	2
Acetylcysteine	0.000000	0.000000	0.0000000	1
Aciclovir	0.533018	0.055406	0.5945314	4
Tropicamide	0.760831	0.082356	0.8678717	9
Xylometazoline	0.705582	0.073039	0.7852226	7
Apraclonidine	0.460946	0.048843	0.5196086	3
Brinzolamide	0.992516	0.084975	1.0000000	12
Bromfenac	0.715700	0.076816	0.8125378	8
Cartelol	0.860836	0.084970	0.9336320	11
Cyclopentolate	0.825830	0.083163	0.9053644	10
Iodoxamide	0.667183	0.065496	0.7215320	5
Ganciclovir	0.664989	0.067563	0.7324594	6



**Figure 3.** Weights of physicochemical properties of ophthalmic drugs



**Figure 4.** VIKOR ranking of Ophthalmic drugs based on molecular weight  
Abbreviations: MR: Molar refractivity; MV: Molar volume; MW: Molecular weight

an antioxidant. Additionally, its well-established mucolytic agent aligns with high desirability in MCDM analysis, reinforcing its top ranking. Conversely, brinzolamide remains the least ranked candidate due to its higher molecular weight and complexity. Brinzolamide ranks last primarily because its heterocyclic and highly irregular graph topology creates uneven degree distributions and

weaker performance in degree-based and Sombor-type topological indices, which favor structurally balanced molecules. Its specialized pharmacological use for glaucoma management limits its broader therapeutic desirability in multi-criteria ranking frameworks compared with drugs exhibiting wider clinical applicability. Additionally, the presence of multiple heteroatoms and rigid ring

**Table 13.** VIKOR ranking of eye drugs based on molar weight.

Ophthalmic drugs	$S_j$	$R_j$	$Q_j$	Molecular weight
Acetazolamide	0.294772	0.030830	0.3259944	2
Acetylcysteine	0.000000	0.000000	0.000000	1
Aciclovir	0.531909	0.051451	0.5641377	4
Tropicamide	0.757596	0.083069	0.8597464	9
Xylometazoline	0.702177	0.068569	0.7482064	7
Apraclonidine	0.459445	0.045396	0.4925774	3
Brinzolamide	0.995443	0.086666	1.0000000	12
Bromfenac	0.713226	0.071332	0.7697867	8
Cartelol	0.858819	0.082283	0.9060582	11
Cyclopentolate	0.818184	0.078024	0.8610233	10
Iodoxamide	0.665888	0.064510	0.7062766	6
Ganciclovir	0.663442	0.062739	0.6951228	5

**Table 14.** VIKOR ranking of ophthalmic drugs based on physicochemical properties

Ophthalmic drugs	$P$	$MV$	$MR$	$MW$
Acetazolamide	2	2	2	2
Acetylcysteine	1	1	1	1
Aciclovir	4	4	4	4
Tropicamide	9	9	9	9
Xylometazoline	8	8	7	7
Apraclonidine	3	3	3	3
Brinzolamide	12	12	12	12
Bromfenac	7	7	8	8
Cartelol	11	10	11	11
Cyclopentolate	10	11	10	10
Iodoxamide	5	6	5	6
Ganciclovir	6	5	6	5

Abbreviations: P: Polarizability; MR: Molar refractivity;  
MV: Molar volume; MW: Molecular weight.

systems reduces structural efficiency in graph descriptors, leading to lower scores beyond simple considerations of molecular weight. The stability of rankings in accordance with properties confirms the potency of VIKOR method in delivering a valid and compromise-based drug priority.

#### 4. Conclusion

This study highlights that the combination of graph-theoretical topological indices with the QSPR modeling framework, and VIKOR-based MCDM analysis provides a reliable and systematic framework for the evaluation of ophthalmic drug candidates. The computed topological indices effectively capture slight divergences in molecular structure, supporting accurate prediction of key physicochemical properties such as polarizability, molar refractivity, molar volume,

and molecular weight. The strong agreement between predicted and observed values confirms the robustness of the regression models and highlights the suitability of neighborhood-based indices in representing molecular behavior. The application of the VIKOR method further enhances the analysis by combining multiple physicochemical factors into a single compromise-based ranking scheme. The resulting hierarchy consistently identifies acetylcysteine as the most structurally efficient compound, reflecting its balanced molecular size, connectivity, and electronic characteristics.

Overall, the proposed computational framework reduces dependence on laboratory screening and offers a reproducible decision-support tool for early-stage drug evaluation. By connecting mathematical modeling with pharmaceutical analysis,

this approach contributes insights into drug priority and may be extended to other classes of remedial agents in future investigations.

## Acknowledgments

None.

## Funding

This work is funded by Faculty of Engineering and Quantity Surviving, INTI International University Colleges, Nilai, Malaysia.

## Conflict of interest

The authors declare they have no competing interests.

## Author contributions

*Conceptualization:* Muhammad Kamran, Shereen Iqbal, Hifza Iqbal, Muhammad Akhtar Tarar, Mohamad Hafez

*Investigation:* Shereen Iqbal, Hifza Iqbal, Muhammad Akhtar Tarar, Muhammad Kamran

*Methodology:* Shereen Iqbal, Hifza Iqbal, Muhammad Akhtar Tarar

*Formal analysis:* Muhammad Kamran, Muhammad Farman, Evren Hincal

*Writing—original draft:* Shereen Iqbal, Hifza Iqbal

*Writing—review & editing:* Muhammad Kamran, Muhammad Farman, Mohamad Hafez, Evren Hincal

## Availability of data

Not applicable.







## AI tools statement

All authors confirm that no AI tools were used in the preparation of this manuscript.

## References

1. Richter Grace M, et al. Trabecular procedures combined with cataract surgery for open-angle glaucoma: a report by the American Academy of Ophthalmology. *Ophth.* 2024;131(3):370-382. <https://doi.org/10.1016/j.ophtha.2023.09.010>
2. Steinmetz Jaimie D, et al. Causes of blindness and vision impairment in 2020 and trends over 30 years, and prevalence of avoidable blindness in relation to VISION 2020: the Right to Sight: an analysis for the Global Burden of Disease Study. *Lancet Glob Health* 2021; 9(2):e144-e160. [https://doi.org/10.1016/S2214-109X\(20\)30489-7](https://doi.org/10.1016/S2214-109X(20)30489-7)
3. Stein Joshua D, Anthony P Khawaja, Jennifer S Weizer. Glaucoma in adults screening, diagnosis, and management: a review. *Jama.* 2021;325(2):164-174. <https://doi.org/10.1001/jama.2020.21899>
4. Wong Wan Ling, et al. Global prevalence of age-related macular degeneration and disease burden projection for 2020 and 2040: a systematic review and meta-analysis. *Lancet Glob Health.* 2014;2(2):e106-e116. [https://doi.org/10.1016/S2214-109X\(13\)70145-1](https://doi.org/10.1016/S2214-109X(13)70145-1)
5. Gyawali Rajendra, et al. Quality of 2019 American optometric association clinical practice guideline for diabetic eye care. *Ophthalmic Physiol Opt.* 2021;41(1):165-170. <https://doi.org/10.1111/opo.12763>
6. Alhazmi Muflih, Sayed Saber. Glucose-insulin regulatory system: Chaos control and stability analysis via Atangana Baleanu fractal-fractional derivatives. *Alex Eng J.* 2025;122:77-90. <https://doi.org/10.1016/j.aej.2025.02.066>
7. Maqbool Saadia, et al. Acute conjunctivitis through a public health lens: a review. *Biomedica.* 2024;40(4):153-158. <https://doi.org/10.24911/BioMedica/5-1165>
8. Azari Amir A, Neal P Barney. Conjunctivitis: a systematic review of diagnosis and treatment. *Jama.* 2013;310(16). <https://doi.org/10.1001/jama.2013.280318>
9. Althubiani Mohammed, Sayed Saber. Hyers Ulam stability of fractal fractional computer virus models with the Atangana Baleanu operator. *Fractal Fract.* 2025;9(3):158. <https://doi.org/10.3390/fractalfract9030158>
10. Leung, Alexander KC, et al. Bacterial conjunctivitis in childhood: etiology, clinical manifestations, diagnosis, and management. *Recent Pat Inflamm Allergy Drug Discov.* 2018;12(2):120-127. <https://doi.org/10.2174/1872213X12666180129165718>
11. Rf Brubaker. Flow of aqueous humor in humans. *Invest Ophthalmol Vis Sci.* 1991;32:3145-3166. <https://doi.org/10.1167/iovs.32.13.3145>
12. Greig Sarah L, Emma D. Deeks. Brinzolamide/brimonidine: a review of its use in patients with open-angle glaucoma or ocular hypertension. *Drugs & aging* 2015;32(3):251-260. <https://doi.org/10.1007/s40266-015-0250-4>
13. Irfan Sameera. Original research The Efficacy & Safety of 0. 3% Acetylcysteine eyedrops in Filamentary keratitis. *Pak J ophthalmol.* 2019;35(4):263-270. <https://doi.org/10.36351/pjo.v35i4.890>
14. Brandt Curtis R, Leeanne M Coakley, Dorene R Grau. A murine model of herpes simplex virus-induced ocular disease for antiviral drug testing. *J Virol Methods.* 1992;36(3):209-222. [https://doi.org/10.1016/0166-0934\(92\)90052-F](https://doi.org/10.1016/0166-0934(92)90052-F)
15. Chou Timothy Y, Bennett Y Hong. Ganciclovir ophthalmic gel 0.15% for the treatment of acute

- herpetic keratitis: background, effectiveness, tolerability, safety, and future applications. *Ther Clin Risk Manag.* 2014;665-681.  
<https://doi.org/10.2147/TCRM.S58242>
16. Yilmaz Medine, et al. The effects of tropicamide and cyclopentolate hydrochloride on laser flare meter measurements in uveitis patients: a comparative study. *Int Ophthalmol.* 2021;41(3):853-857.  
<https://doi.org/10.1007/s10792-020-01639-3>
17. Contreras-Salinas Homero, et al. Ocular cyclopentolate: a mini review concerning its benefits and risks. *Clin Ophthalmol.* 2022;16:3753.  
<https://doi.org/10.2147/OPTH.S388982>
18. Hussein Ola G, et al. Exquisite integration of quality-by-design and green analytical approaches for simultaneous determination of xylometazoline and antazoline in eye drops and rabbit aqueous humor, application to stability study. *J Pharm Biomed Anal.* 2023;235:115598.  
<https://doi.org/10.1016/j.jpba.2023.115598>
19. Grob Seanna R, et al. Pharmacologic agents used in the assessment or correction of blepharoptosis: a report by the American Academy of ophthalmology. *Ophthalmology.* 2025;132(6):718-730.  
<https://doi.org/10.1016/j.optha.2025.01.018>
20. Schechter Barry A. Use of topical bromfenac for treating ocular pain and inflammation beyond cataract surgery: a review of published studies. *Clin Ophthalmol.* 2019;1439-1460.  
<https://doi.org/10.2147/OPTH.S208700>
21. Mirza Ertugrul G, Sarper Karaküçük, Erhan Temel. Comparison of the effects of 0.5% timolol maleate, 2% carteolol hydrochloride, and 0.3% metipranolol on intraocular pressure and perimetry findings and evaluation of their ocular and systemic effects. *J Glaucoma.* 2000;9(1):45-50.  
<https://doi.org/10.1097/00061198-200002000-00009>
22. Denis D, et al. Treatment of common ocular allergic disorders; a comparison of lodoxamide and NAAGA. *Br J Ophthalmol.* 1998;82(10):1135-1138.  
<https://doi.org/10.1136/bjo.82.10.1135>
23. Pfuhl Gerit, et al. A preliminary study on the role of personal history of infectious and parasitic diseases on self-reported health across countries. *Public Health.* 2025;242:220-227.  
<https://doi.org/10.1016/j.puhe.2025.02.030>
24. Ebekozien, Andrew, et al. Affordable housing in Nigerias slums: combating infectious diseases and advancing SDG 3. *Property Management* 2025;43(4):600-620.  
<https://doi.org/10.1108/PM-08-2024-0092>
25. Sachs Horst, Michael Stiebitz, Robin J Wilson. An historical note: Euler's Königsberg letters. *J Graph Theory.* 1988;12(1):133-139.  
<https://doi.org/10.1002/jgt.3190120114>
26. Çolakoglu, Özge, Muhammad Kamran, and Ebenezer Bonyah. M-Polynomial and NM-Polynomial of Used Drugs against Monkeypox. *J Math.* 2022;2022(1):9971255.  
<https://doi.org/10.1155/2022/9971255>
27. Bougueroua Sana, et al. Exploiting graph theory in MD simulations for extracting chemical and physical properties of materials. *Phys Chem Chem Phys.* 2025;27(3):1298-1309.  
<https://doi.org/10.1039/D4CP02764G>
28. Adamopoulos Ioannis, et al. Utilizing graph theory algorithms for the modeling and analysis of COVID-19 infection dynamics. *J Artif Intell Healthc.* 2025;2025:1-11.  
<https://doi.org/10.58496/MJAIH/2025/001>
29. Chawla Apporva, et al. QSPR modeling. *Computational Drug Delivery.* 2024;183.  
<https://doi.org/10.1515/9783111208671-009>
30. Hosoya Haruo. Topological index. A newly proposed quantity characterizing the topological nature of structural isomers of saturated hydrocarbons. *Bull Chem Soc Jpn.* 197144(9):2332-2339.  
<https://doi.org/10.1246/bcsj.44.2332>
31. Wiener, Harry. Structural determination of paraffin boiling points. *J Am Chem Soc.* 1947;69(1):17-20.  
<https://doi.org/10.1021/ja01193a005>
32. Voit Brigitte I, Alben Lederer. Hyperbranched and Highly Branched Polymer Architectures? Synthetic Strategies and Major Characterization Aspects. *Chem Rev.* 2009;109(11):5924-5973.  
<https://doi.org/10.1021/cr900068q>
33. Kamran Muhammad, et al. Physico-Chemical Characterization of Amylose and Amylopectin Using Revan Topological Indices. Physico-Chemical Characterization of Amylose and Amylopectin Using Revan Topological Indices. *J Math.* 2022;2022(1):2840217.  
<https://doi.org/10.1155/2022/2840217>
34. Gutman Ivan, Nenad Trinajstić. Graph theory and molecular orbitals. *New concepts II. Berlin, Heidelberg: Springer Berlin Heidelberg.* 2005;49-93.  
[https://doi.org/10.1007/3-540-06399-4\\_5](https://doi.org/10.1007/3-540-06399-4_5)
35. Shirdel Gholam Hassan, Hassan Rezapour, AM Sayadi. The hyper-Zagreb index of graph operations. 2013:213-220.  
<https://doi.org/10.22052/IJMC.2013.5294>
36. Khadikar Padmakar V, et al. The Szeged index and an analogy with the Wiener index. *J Chem Inf Comput Sci.* 35.3 (1995): 547-550.  
<https://doi.org/10.1021/ci00025a024>
37. Uyanik Gülden Kaya, Nese Güler. A study on multiple linear regression analysis. *Procedia Soc Behav Sci.* 2013;106:234-240.  
<https://doi.org/10.1016/j.sbspro.2013.12.027>
38. Kamran Muhammad, et al. Enhancing decision-making in environmental management: a complex picture fuzzy soft set approach to sustainability assessment: M. Kamran et al. *Comp Appl Math.* 2025;44(5):258.  
<https://doi.org/10.1007/s40314-025-03221-y>
39. Saeid ABM. Comprehensive Examination of Case Studies: Applications of Multi-Criteria

- Decision-Making (MCDM) Approach Across Diverse Fields. *FROM THE EDITOR* 2025;3(1):57-74.  
<https://doi.org/10.5281/zenodo.16751556>
40. Pawar, Sumathi, et al. Survey and Analysis on Machine Learning Approaches for Exploring Aspects of Covid-19. International Conference on Machine Vision and Augmented Intelligence. Singapore: Springer Nature Singapore, 2023.  
[https://doi.org/10.1007/978-981-97-4359-9\\_25](https://doi.org/10.1007/978-981-97-4359-9_25)
  41. Iqbal, Shereen, et al. Evaluation of antiarrhythmia drug through QSPR modeling and multi criteria decision analysis. *Sci Rep.* 2025;15(1):29216.  
<https://doi.org/10.1038/s41598-025-14892-2>
  42. Zaidi Noushi, et al. Pharmaceuticals and personal care products as emerging contaminants: environmental fate, detection, and mitigation strategies. *J Environ Anal Chem.* 2025;105(20):8837-8865.  
<https://doi.org/10.1080/03067319.2025.2484456>
  43. Yu, Dejian, and Tong Ye. Uncovering the academic evolution of VIKOR method: A comprehensive main path analysis. *J Oper Res Soc.* 2025;76(2):393-409.  
<https://doi.org/10.1080/01605682.2024.2359726>
  44. Zeleny, Milan. Multiple criteria decision making (MCDM): From paradigm lost to paradigm regained? *Journal of Multi-Criteria Decision Analysis.* 2011;18(1-2):77-89.  
<https://doi.org/10.1002/mcda.473>
  45. Ballesteros Enrique, Carlos Romero. Multiple Criteria Decision Making: An Introduction. Multiple Criteria Decision Making and its Applications to Economic Problems. Boston, MA: Springer US, 1998;1-10.  
[https://doi.org/10.1007/978-1-4757-2827-9\\_1](https://doi.org/10.1007/978-1-4757-2827-9_1)
- Muhammad Kamran** holds multiple academic affiliations, serving as an adjunct faculty member at International Center for Interdisciplinary Research in Sciences, The University of Lahore, Lahore, Pakistan. His research is dedicated to fuzzy logics, graph-based networks, and intelligent systems.  
 <https://orcid.org/0009-0000-5467-0497>
- Shereen Iqbal** is a Ph.D. scholar in the Department of Mathematics and Statistics at The University of Lahore, Pakistan. Her current research interests lie at the intersection of chemical graph theory, QSPR analysis, and the application of machine learning techniques. She holds an M.Phil. degree, during which her work focused on mathematical inequalities.  
 <https://orcid.org/0009-0006-0734-1184>
- Hifza Iqbal** completed her Ph.D. at The University of Lahore, Pakistan, in 2021. She is currently serving as an Assistant Professor in the Department of Mathematics and Statistics at the same institution. Her research interests are primarily in graph theory and computational wave propagation.  
 <https://orcid.org/0009-0009-1614-9110>
- Muhammad Akhtar Tarar** is Associate Professor (Civil Engineering) at The University of Lahore, Lahore, Pakistan.  
 <https://orcid.org/0009-0008-6788-0870>
- Muhammad Farman** did Ph.D. Mathematics from the University of Lahore in 2019, Pakistan, and a Postdoc fellowship from Universitas Airlangga Indonesia, Lebanese American University, and Near East University, North Cyprus, Turkey. Adjunct Professor (Research), at Universitas Airlangga Indonesia and University of Lahore, Pakistan. Currently, he is working as an Associate Professor Dr. at Near East University. He has more than eight years of teaching and research experience. The research field is Mathematical biology, Control theory, Numerical Analysis, and Application of fractional calculus on Epidemic models. He published more than three hundred research papers in well-reputed SCI and Scopus journals and presented his work at several national and international conferences. He is the winner of several awards, such as the Research Excellent Awards (Best Researcher of the University), in 2019 and 2020 at the University of Lahore. Stanford University's Top 2% Scientists of 203, 2024 in Applied Mathematics. Best Academic Performance Award 2024, Publication Honorary Award 2024, Most Impactful Researcher Award 2024, Near East University, 2024.  
 <https://orcid.org/0000-0001-7616-0500>
- Evren Hincal** is a Professor at the Mathematics Research Center, Department of Mathematics, Near East University, Mersin, Turkey. His research focuses on computational mathematics.  
 <https://orcid.org/0000-0001-6175-1455>
- Mohamad Hafez** is a Professor at the Faculty of Engineering and Quantity Surveying, INTI International University Colleges, Nilai, Negeri Sembilan, Malaysia. His research focuses on sustainability and various other applications of mathematics.

