

REVIEW ARTICLE

Nanomaterials in organoid research: Advancements and applications

Shuo Zhang^{1,2†}, Xu Chu^{3†}, Zihang Zhao⁴, Guilai Zuo^{4*}, Hongbo Zhang^{1,2,5*} , and Wenguo Cui^{1*} 

¹Department of Orthopaedics, Shanghai Key Laboratory for Prevention and Treatment of Bone and Joint Diseases, Shanghai Institute of Traumatology and Orthopaedics, Ruijin Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai, China

²Pharmaceutical Sciences Laboratory, Faculty of Science and Engineering, Åbo Akademi University, Turku, Southwest Finland, Finland

³Department of Thoracic Surgery, The First Affiliated Hospital of Bengbu Medical University, Bengbu, Anhui, China

⁴Department of Orthopedic Oncology, The Affiliated Hospital of Qingdao University, Qingdao, Shandong, China

⁵Turku Bioscience Centre, University of Turku and Åbo Akademi University, Turku, Southwest Finland, Finland

*Corresponding authors: Guilai Zuo (zuoguilai@qduhospital.cn); Hongbo Zhang (hongbo.zhang@abo.fi); Wenguo Cui (wgcai@sjtu.edu.cn)

[†]These authors contributed equally to this work.

Citation: Zhang S, Chu X, Zhao Z, Zuo G, Zhang H, Cui W. Nanomaterials in organoid research: Advancements and applications. *Organoid Res.* 2026;2(3):026090015. doi: 10.36922/OR026090015

Received: March 1, 2026

Revised: April 14, 2026

Accepted: April 15, 2026

Published online: May 12, 2026

Copyright: © 2026 Author(s). This is an Open-Access article distributed under the terms of the Creative Commons Attribution License, permitting distribution, and reproduction in any medium, which provided that the original work is properly cited.

Publisher's Note: AccScience Publishing remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Abstract

In recent years, organoids have demonstrated immense potential as three-dimensional in vitro models of human tissue, capable of simulating physiological and pathological environments. Nanomaterials, owing to their unique physicochemical properties, have become pivotal tools in advancing organoid research and applications. By integrating nanomaterials, scientists can more effectively explore disease mechanisms, optimize drug delivery systems, and develop novel therapeutic strategies. For example, nanoparticle-based metabolic reprogramming techniques offer novel avenues for regulating organ function, while advances in gene editing technologies have markedly enhanced the precision and complexity of organoid models. Furthermore, organoids demonstrate broad application prospects in fields such as virology research, cardiac tissue engineering, and cancer therapy. These advances not only propel fundamental scientific inquiry but also provide crucial support for clinical translation. The future holds promise for achieving more precise, personalized medical solutions through the integration of nanotechnology and organoid models.

Keywords: Nanomaterials; Organoid models; Drug delivery; Disease modeling; Personalized medicine

1. Introduction

1.1. Background and importance of organoids

Over the past few decades, traditional cell lines have been the most commonly used experimental models in cell biology and medical research, playing a crucial role in many major medical discoveries.¹ However, as research questions become increasingly complex, traditional cell lines have begun to reveal significant limitations, making them inadequate for modern biomedical research. This is especially true in areas such as drug pharmacology

assessment for complex diseases and the study of the mechanisms of organogenesis and development. The applicability of classical cell lines is constrained by their limited differentiation ability, homogeneous cell sources, and inability to replicate the multicellular composition and dynamic changes typical of in vivo tissues.²

Thus, the development of new research models that more closely mimic real physiological states has become a critical issue in modern biomedical research. In recent years, as stem cell technologies have progressed, organoid technology has matured, offering a new approach in this

field.³ Organoids are three-dimensional cell aggregates formed by stem cells in vitro and have a defined spatial structure.⁴ These structures have the ability to self-renew and self-organize, closely mimicking the structure and function of actual tissues.⁵ By cultivating stem cells derived from normal or diseased tissues, it is possible to recreate key features of tissues or tumors in vitro, providing an effective model for studying their development mechanisms.^{6,7}

Compared with traditional two-dimensional cell culture systems, organoid models have significant advantages in mimicking the tissue microenvironment. By enhancing interactions between cells, they more accurately replicate the spatial configuration and functional characteristics of in vivo tissues.⁸ As such, organoids bridge the gap between two-dimensional cell models and animal experiments, offering a more physiologically relevant experimental platform for drug screening, toxicity assessments, and disease mechanism studies.⁹

1.2. Role of nanomaterials in organoid engineering

Nanomaterials are materials that have at least one dimension in the range of nanometers (1–100 nm), positioned between atomic, molecular, and macroscopic structures. These materials are composed of nanoparticles and possess unique physicochemical properties, making them widely applied in various biomedical fields. Studies have shown that nanomaterials offer greater targeting precision, therapeutic accuracy, and safety in cancer treatment.¹⁰ Additionally, owing to the size effects of nanotechnology, nanomaterials play a significant role in drug delivery, medical imaging, and immunotherapy.^{11–13} Their optical, magnetic, and electrical properties have made them a focal point in biotechnology and bioengineering research, further promoting the development of organoid-related studies.¹⁴

In organoid research, nanomaterials have been extensively applied in various functional modulations. For example, in liver organoid models, iron oxide nanoparticles are used for non-invasive dynamic tracking of hepatocyte differentiation.¹⁵ Moreover, nanomaterials responsive to physical stimuli, such as light and electrical signals, have been developed to directly modulate stem cell surface receptor activity, providing new technological means for exploring the complex signaling networks involved in organoid construction and differentiation. However, the incorporation of nanomaterials into organoid research does not simply confer technical advantages but also introduces additional challenges associated with system complexity. On the one hand, nanomaterials provide new strategies for addressing key limitations that are difficult to overcome using conventional approaches, such as precise delivery to deep tissue regions, real-time non-invasive monitoring of the organoid microenvironment, and spatiotemporal

regulation of stem cell fate. On the other hand, their integration may increase the physicochemical complexity of the system, alter local microenvironmental parameters, and potentially disturb the intrinsic physiological homeostasis of organoids.¹⁶ Therefore, the integration of nanotechnology into organoid systems should be guided by clearly defined mechanistic objectives or practical therapeutic needs, rather than by simple technical accumulation. Overall, such interdisciplinary integration is of substantial mechanistic and translational value only when the biological insights gained or therapeutic benefits achieved clearly outweigh the added complexity introduced by nanomaterials.¹⁷ However, systematic reviews focusing on the interaction between nanomaterials and organoids are still relatively scarce, particularly research reports centered on using nanoscience and nanotechnology to construct organoid microenvironments.^{18,19} There remains a lack of comprehensive reviews on how nanomaterials directly influence stem cell fate decisions and how they can provide a three-dimensional biomimetic microenvironment for organoids.²⁰

Therefore, the role of nanomaterials—both inorganic and carbon-based—in regulating organoid structure and function requires further investigation. This paper reviews the latest advancements in nanomaterials in organoid research, discussing aspects such as material design, functionality, and applications in biomedical fields (Figure 1).

2. Design of nanomaterials in organoid engineering

2.1. Types and properties of nanomaterials: Inorganic/organic/composite nanomaterials

2.1.1. Inorganic materials

Inorganic nanomaterials, due to their unique physicochemical properties, exhibit significant advantages in the diagnosis and treatment of diseases. For example, gold nanoparticles exhibit exceptional biocompatibility, a high surface-to-volume ratio amenable to functionalization, and superior optical and surface plasmon resonance characteristics, rendering them highly attractive for photothermal therapy and drug delivery applications.²¹ In photothermal therapy, gold nanoparticles exhibit the capacity to absorb light at specific wavelengths and convert it efficiently into thermal energy, enabling localized thermal ablation of tumor tissue with minimal collateral damage to adjacent healthy tissues.²²

Silica nanoparticles, especially mesoporous silica nanoparticles (MSNs), are increasingly used as multifunctional carrier platforms due to their regular porous structure, high surface area, and tunable pore sizes²³ (Figure 2). These features allow them to effectively load




Figure 1. Research and progress of nanomaterials in organoid engineering. Image created by the authors with Nano Banana Pro.

various therapeutic molecules, including small molecules and biological macromolecules. Furthermore, MSNs are often designed as theranostic systems that integrate both diagnostic and therapeutic functions, enabling simultaneous disease imaging and drug delivery.²⁴ Owing to their distinctive optical characteristics—including broad excitation spectra, narrow emission bands, and tunable emission wavelengths—quantum dots have emerged as pivotal tools in biomedical applications.²⁵ These advantages make them excellent for high-sensitivity fluorescence labeling, widely used for protein and nucleic acid detection and analysis. Quantum dots also serve as high-resolution imaging probes for live-cell single-molecule tracking and studying intracellular dynamic behaviors.²⁶

2.1.2. Organic materials

Lipid-based nanodelivery systems are mature and widely used in drug delivery platforms due to their excellent biocompatibility and efficient drug encapsulation capabilities.²⁷ Liposomes, as a typical representative, consist of phospholipid bilayer vesicles capable of carrying both hydrophilic and hydrophobic drugs.²⁸ Doxil exemplifies a liposomal formulation that exhibits preferential accumulation at tumor sites via the enhanced permeability and retention effect, thereby enhancing therapeutic efficacy.²⁹ Solid lipid nanoparticles are composed of biocompatible solid lipids and are suitable for delivering poorly soluble drugs. Their structure helps improve drug stability, reduce enzymatic degradation, and enable controlled release, improving the solubility of drugs.³⁰ Nanostructured lipid

carriers integrate the advantageous characteristics of both solid and liquid lipids, thereby improving drug-loading capacity and encapsulation efficiency, while enabling precise modulation of drug release kinetics^{31,32} (Figure 3).

2.1.3. Composite nanomaterials

Polymer nanoparticles constructed from biodegradable and biocompatible materials, such as poly(lactic-co-glycolic acid) (PLGA), polycaprolactone, and chitosan, serve as important platforms for drug delivery. They are commonly used for sustained and controlled drug release, extending treatment durations and reducing dosing frequency.³³ By adjusting the composition ratio and molecular weight of the polymer, the degradation behavior and drug release kinetics can be precisely controlled.³⁴ For example, PLGA nanoparticles are widely used for delivering anti-inflammatory and anti-tumor drugs, enhancing targeting efficiency, reducing systemic side effects, and improving therapeutic efficacy.³⁵ Chitosan-based nanoparticles, recognized for their outstanding biocompatibility and versatile chemical modifiability, hold significant promise for non-oral drug delivery applications.³⁶

2.2. Surface functionalization of nanomaterials

Through surface functionalization, nanodrug delivery systems can achieve active targeting capabilities, enhancing their selective accumulation in specific tissues or cells.³⁷ This is typically done by conjugating antibodies, peptides, aptamers, or small-molecule ligands to the surface of nanoparticles, enabling them to recognize and bind to




Figure 2. Mesoporous silica nanoparticle (MSN) synthesis via sol-gel method and evaporation-induced self-assembly. Reprinted from Dutta Gupta *et al.*²³

Abbreviation: CTAB: Cetyltrimethylammonium bromide.




Figure 3. Structural features of liposomes, solid lipid nanoparticles, and nanostructured lipid carriers. Reprinted with permission from Barroso *et al.*³¹

overexpressed receptors or specific biomarkers in disease cells. This molecular recognition strategy not only improves therapeutic efficiency but also reduces nonspecific toxicity to normal tissues.

Antibodies, known for their highly specific antigen-binding abilities, are widely used in the construction of targeted nanoparticle platforms. For example, cetuximab-conjugated polysaccharide-ester nanoparticles have demonstrated enhanced anti-tumor efficacy in murine models.³⁸ Hyaluronic acid-functionalized nanoparticles have also demonstrated enhanced targeted delivery by binding to receptors overexpressed on tumor cells.³⁹

Furthermore, polyethylene glycolation (PEGylation) diminishes the recognition and clearance of nanoparticles by the mononuclear phagocyte system, thereby extending their circulation half-life and improving their stability and bioavailability.⁴⁰ However, the role of PEGylation in in vitro models such as organoids should be re-evaluated in the context of the specific experimental setting. Although organoid systems cannot recapitulate systemic circulation or mononuclear phagocyte system-mediated clearance in vivo, PEG modification remains relevant under in vitro conditions. In particular, PEGylation can improve the colloidal stability of nanoparticles, reduce nonspecific

adsorption in the culture system, and minimize particle aggregation. In addition, the PEG layer may modulate nanoparticle interactions with the extracellular matrix (ECM), thereby indirectly influencing their distribution and penetration within organoids. Nevertheless, it should be noted that the well-recognized advantages of PEGylation in prolonging circulation time and reducing immune clearance cannot be fully assessed in static organoid systems. Therefore, when PEGylated nanoparticles are

investigated using organoid models, their physicochemical effects *in vitro* should be clearly distinguished from their biological functions in systemic *in vivo* delivery, in order to avoid overinterpretation of their overall performance.⁴¹ By combining targeting molecules with pharmacokinetic control strategies, rationally designing the surface structure of nanoparticles helps overcome biological barriers and achieve precise spatial control and efficient drug delivery (Figure 4).




Figure 4. Schematic diagram of the treatment of liver cancer through inducing cell apoptosis using Apt/PPII-NPs. (A) Synthesis process. (B) Anti-tumor effect of Apt/PPII-NPs. (C) Apoptosis mechanism induced by Apt/PPII-NPs. Reprinted from Huang *et al.*²⁷

Abbreviations: Apt: Aptamer AS1411; Bax: Bcl-2-associated X protein; Bcl-2: B-cell lymphoma 2; Cyt-c: Cytochrome c; FADD: Fas-associated protein with death domain; FasL: Fas ligand; FasR: Fas receptor; NP: Nanoparticle; PARP: Poly (ADP-ribose) polymerase; PDA: Polydopamine; PPII: Polyphyllin II.

2.3. Conceptual framework for nanomaterial selection

A practical design of nanoparticle–organoid systems requires an integrated conceptual framework for nanomaterial selection. Such a framework should consider the intended application type, the biological characteristics of the organoid model, the presence of biological barriers, payload compatibility, targeting requirements, and safety considerations. Specifically, material choice should be aligned with the research objective, such as therapy, imaging, sensing, or tissue engineering, while also accounting for tissue-specific and pathological features of the organoid model.⁴² In addition, barriers such as dense ECM, elevated interstitial pressure, and hypoxic cores may critically influence nanoparticle penetration and distribution. Carrier selection should also match the properties of the intended payload, whereas surface functionalization strategies should be tailored to the need for active targeting of specific cell populations. Equally important is the evaluation of biodegradability, long-term toxicity, and unintended effects on organoid development. By standardizing these considerations, this framework may support the development of more precise and application-oriented nanoengineered organoid platforms.⁴³

3. Distribution of nanomaterials in organoids

The spatial distribution of nanoparticles within the body is one of the key factors determining their therapeutic efficacy, as their localization in tissues directly influences the recognition efficiency of specific target cells and the subsequent activation of biological responses. In this process, the ECM plays a crucial role as a physical barrier. ECM is composed of proteins and polysaccharides that form a dense, complex network structure, significantly limiting the diffusion and penetration of nanoparticles within tissues. To address these limitations and enhance research consistency, engineered and chemically defined matrices are increasingly employed to standardize nanoparticle transport studies. Researchers are increasingly turning to engineered and chemically defined matrices. Compared with naturally derived matrices such as Matrigel, these synthetic or semi-synthetic matrices can improve standardization by allowing precise control over several key parameters.⁴⁴ These include matrix density and porosity, which determine the physical channels available for nanoparticle diffusion through modulation of polymer concentration and crosslinking density; stiffness, which enables the simulation of tissue-specific mechanical properties and thereby standardizes the influence of biomechanical cues on nanoparticle uptake; and biochemical composition, where defined ligand densities and spatial organization of biochemical factors help reduce nonspecific interactions and render transport

studies more controllable.

Nevertheless, it should be noted that full standardization remains challenging, because matrix effects interact in complex ways with the intrinsic variability of organoids themselves, including differences in cellular organization and secretory profiles.⁴⁵ In organoid systems, particularly tumor organoids, the ECM not only replicates the structural environment of solid tumors but also mimics typical mechanical features, such as increased tissue stiffness and elevated interstitial fluid pressure. These factors have been shown to reduce the penetration ability of nanoparticles, leading to uneven drug distribution.

The intricate three-dimensional architecture within organoids—encompassing stratified cellular organization, nutrient and oxygen gradients, and a hypoxic core—further modulates the spatial distribution and diffusion dynamics of nanomaterials (Figure 5). In tumor organoids specifically, hypoxic regions frequently establish significant additional barriers to drug delivery. It should also be noted that conventional tumor organoid models cannot fully recapitulate the complex vascular architecture *in vivo*. Most current organoid systems primarily reproduce tissue structural features and the associated diffusion barriers, but lack authentic vascular networks, limiting their ability to accurately model nanoparticle delivery *in vivo*.

In recent years, vascularized organoids and microfluidic perfusion systems have partially addressed this limitation.⁴⁶ By incorporating endothelial cells or dynamic fluidic environments, these models can simulate certain aspects of vascular-associated transport, thereby improving the accuracy of nanoparticle delivery studies. However, these advanced systems still fall short of fully reproducing the complex hemodynamic properties and vascular–tissue interactions found *in vivo*, and therefore remain limited in their ability to predict nanoparticle biodistribution and delivery efficiency. Accordingly, when organoid models are used to evaluate nanoparticle delivery performance, their limitations in vascular simulation should be explicitly recognized, and the results should ideally be interpreted in combination with *in vivo* or other physiologically relevant models to obtain more reliable conclusions.⁴⁷

To overcome limitations in nanoparticle distribution within complex tissues, advanced imaging technologies are widely used to evaluate and optimize nanoparticle localization and penetration behaviors. For example, multiphoton microscopy (MPM) and optical coherence tomography (OCT) enable high-resolution, three-dimensional imaging of nanoparticles within organoids. By employing nonlinear fluorescence excitation mechanisms and long-wavelength lasers, MPM overcomes the resolution and tissue penetration constraints inherent in conventional single-photon microscopy.⁴⁸ This technology

has demonstrated significant efficacy in studies of brain organoids in vivo, facilitating the observation of cell migration, neuronal network anomalies, differentiation impairments, and the progression of neurodegenerative diseases and tumor models.⁴⁹ Similarly, in retinal organoid research, MPM has been instrumental in elucidating human-specific neurodevelopmental mechanisms and related disease processes.⁵⁰

On the other hand, OCT technology, based on interference measurement principles, offers high spatial resolution and non-invasive imaging advantages, making it valuable for three-dimensional structural reconstruction and real-time dynamic monitoring in organoid research.⁵¹ For example, OCT has been used for real-time three-dimensional imaging of retinal organoids, providing detailed structural information on cell dynamics and tissue remodeling.⁵² Despite these advantages, both techniques have inherent limitations in the context of nanoparticle–organoid interaction analysis.

To more accurately evaluate the penetration capacity and spatial heterogeneity of nanoparticles within organoids, a variety of quantitative analytical approaches have recently been developed, thereby improving the rigor and reliability of related studies. First, three-dimensional fluorescence imaging-based methods, such as confocal microscopy and MPM, are widely used. Through three-dimensional reconstruction of organoids, the fluorescence intensity of nanoparticles can be quantified at different depths, providing an overall assessment of their penetration behavior.⁵³

In addition, radial penetration analysis is commonly applied by using the organoid center as a reference point and calculating the decay profile of fluorescence signals along the radial axis, thereby quantitatively describing the diffusion efficiency of nanoparticles from the periphery toward the core. Furthermore, core-to-periphery distribution metrics can be used to assess the uniformity of nanoparticle distribution within organoids. For example, comparing fluorescence intensity ratios between the core and peripheral regions can directly indicate whether nanoparticles effectively penetrate into deeper structures. At the single-cell level, organoids can be dissociated into individual cells and combined with flow cytometry or single-cell sequencing techniques to quantify nanoparticle uptake. Such approaches help reveal uptake heterogeneity among different cell types and thus provide deeper insight into nanoparticle distribution in complex multicellular systems. In addition, quantitative techniques such as inductively coupled plasma mass spectrometry (ICP-MS) can be employed to precisely measure the metal content of nanoparticles, thereby determining their overall or region-specific accumulation in organoids.⁵⁴ Owing to its high sensitivity and accuracy, this method often serves as a powerful complement to fluorescence imaging.

Collectively, these multiscale and multimodal quantitative strategies are essential for comprehensively evaluating nanoparticle penetration and distribution heterogeneity in organoids, while also providing a more robust basis for cross-study comparison.⁵⁵




Figure 5. Whole-mount staining of vascular spheres (VSs) and cardiac organoids (vcCOs) for endothelial cells (platelet endothelial cell adhesion molecule 1 [PECAM1], red), cardiomyocytes (α -actinin, green), and the nucleus (Hoechst 33342, blue) at indicated days (D10, D15, and D25). Scale bars: 50 μ m. The right panel shows the three-dimensional reconstruction of whole VSs (D10) and vcCOs (D25) using a light plate microscope. Scale bar: 100 μ m. Reprinted from Yang *et al.*⁵⁶

4. Nanomaterials regulating cell proliferation and differentiation in organoids

Organoid models have the capacity to reconstruct tissue-specific three-dimensional structures and microenvironments in vitro, providing an important platform to study the regulatory mechanisms of cell proliferation and differentiation. However, in complex three-dimensional systems such as organoids, ligand–receptor interactions do not always operate in a stable or uniform manner, and their effects are often highly dependent on the local microenvironment.

In organoid models, these interactions are regulated by multiple factors and exhibit marked context dependence.⁵⁷ First, receptor expression levels are often heterogeneous among different cell types and differentiation states, potentially leading to substantial differences in targeting efficiency. Second, the ECM can exert a matrix-shielding effect on nanoparticles, thereby limiting their effective contact with receptors on target cell surfaces. Third, dynamically changing microenvironmental factors within organoids, such as pH, oxygen tension, and metabolic status, may alter the affinity and stability of ligand–receptor binding. Fourth, nanoparticles may undergo surface modifications in biological environments, such as protein corona formation, further affecting their recognition capability and binding behavior. Therefore, ligand–receptor-mediated targeting in organoid systems should be considered a dynamic process that depends on specific experimental conditions, rather than a universally stable mechanism.

Recognizing this context dependence is important for optimizing nanoparticle design strategies and for improving the interpretation of experimental findings.⁵⁸ For example, in reconstituted testicular organoids, spermatogonial stem cells (SSCs) have the potential for proliferation and differentiation. To study the proliferation and differentiation abilities of SSCs in testicular organoids, researchers have conducted immunohistochemical staining.⁵⁹ Proliferating cell nuclear antigen (PCNA) and Ki67-positive SSCs indicate their proliferative potential (Figure 6). Zhang *et al.*⁶⁰ found that the expression patterns of Ki67 and PCNA in SSCs from testicular organoids resembled those in normal mouse testis tissues. C-kit, a marker of differentiating spermatogonia, especially A1–A4 type spermatogonia, was found in testicular organoids, suggesting that SSCs differentiate into A1–A4 spermatogonia within eight weeks. Therefore, testicular organoids can promote the proliferation and initial differentiation of SSCs within seminiferous tubules.

Further reverse transcription-polymerase chain reaction and Western blot analyses confirmed the

expression of multiple germ cell and somatic markers, such as *PLZF*, *GFRA1*, *THY1*, *SYCP3*, and *MLH1*.⁶¹ This indicates that organoids not only maintain the stem cell population but also support the developmental progression toward spermatocytes and more mature stages. It is important to note that cell seeding density and hormonal environments of recipient animals significantly influence the self-renewal and differentiation efficiency of stem cells within organoids. Moderate cell density (4×10^7 cells/mL) is conducive to the formation of seminiferous tubule-like structures, while excessively high or low cell densities may affect tissue reconstruction efficiency.⁶² Hormonal status also regulates SSC differentiation potential, highlighting the critical role of microenvironmental factors in stem cell fate determination.⁶³

5. Nanomaterials guiding organoid tissue development

Nanomaterials in organoid engineering are not intended to replace advanced ECM-based culture systems, but rather to serve as powerful complementary tools that endow these platforms with dynamic functions that cannot be readily achieved by ECM alone.⁶⁴ The integration of nanomaterials with ECM-based systems enables the construction of a microenvironment that is both more physiologically relevant and more controllable.⁶⁵ While ECM provides structural support and essential biochemical cues, nanomaterials introduce several unique engineering advantages. Owing to their precisely tunable physicochemical properties, including size, shape, and surface chemistry, nanomaterials can be optimized to enhance biocompatibility and regulate cell–material interactions in accordance with the specific requirements of different tissues.⁶⁶

In addition, stimulus-responsive nanomaterials, designed to respond to light, magnetic fields, or electrical signals, allow non-invasive and remote modulation of the local microenvironment within organoids or direct activation of cellular receptors. The incorporation of imaging probes, such as quantum dots and magnetic nanoparticles, also enables long-term and real-time monitoring of cell migration, cell differentiation, and neuronal network abnormalities without disrupting organoid architecture. Furthermore, through surface functionalization with antibodies or ligands, nanoparticles can actively target specific cell subpopulations within organoids,⁶⁷ thereby increasing the local concentration of therapeutic agents while reducing nonspecific effects.

Beyond these functions, nanoplateforms can integrate multiple therapeutic and imaging components within a single system, allowing seamless combination of disease modeling, drug delivery evaluation, and real-time therapeutic monitoring.⁶⁸ Looking forward, embedding nanocarriers with spatiotemporally controlled release




Figure 6. Morphological features of mouse testicular organoids. (A) The precise localization of transplantation (left panel) and a representative testicular organoid (right panel) at 8 weeks post-transplantation. Scale bar: 1 mm. (B) Hematoxylin and eosin staining of testicular organoids at 8, 10, 12, 14, and 16 weeks of cell transplantation. The red arrowhead indicates spermatogonial stem cells (SSCs), while the blue arrowhead denotes Sertoli cells; blue circles indicate spermatocytes, and green circles denote spermatids. Scale bars: 10 μm. (C) Double immunostaining of SOX9+ Sertoli cells and PLZF+ SSCs, as well as SOX9+ Sertoli cells and UCHL1+ SSCs, in the eighth testis organoids. White circles denote seminiferous tubules. The red arrows referred to a local zoomed-in image. Scale bars: 10 μm. Reprinted from Zhang *et al.*⁶⁰

Abbreviations: DAPI: 4',6-diamidino-2-phenylindole; PLZF: Promyelocytic leukemia zinc finger; SOX9: Sex-determining region Y-box 9; UCHL1: Ubiquitin C-terminal hydrolase L1.

properties into engineered ECM scaffolds may offer a more realistic means of recapitulating morphogen gradients and dynamic mechanical changes during *in vivo* development. Such deep integration of nanomaterials with ECM is advancing organoids from static culture models toward dynamic and predictive preclinical platforms.⁶⁹

5.1. Nanomaterials as precision delivery tools for organoids

Studies have shown that organoid systems can simulate the structure and signaling networks of *in vivo* tissues, supporting stem cell proliferation and lineage differentiation.⁷⁰ These systems provide an ideal platform for exploring the regulatory mechanisms of exogenous factors (such as nanomaterials, bioactive molecules, and mechanical signals) on cell fate. Under the context of nanomaterial applications, these systems can be used

to evaluate the effects of nanocarriers on stem cell self-renewal, differentiation direction, and microenvironment remodeling, thus laying the experimental foundation for developing nanostrategies to regulate tissue regeneration and functional reconstruction.⁷¹

5.2. Active regulatory mechanisms of nanomaterials on organoid fate

In organoid engineering, the role of nanomaterials has evolved from that of passive carriers to active components capable of modulating cell fate. This active regulation can be achieved through several interconnected mechanisms. At the receptor level, nanomaterials can directly interfere with or activate cell-surface signaling pathways through their specific physicochemical properties;⁷² for example, stimulus-responsive nanomaterials that react to light, magnetic fields, or electrical signals can non-invasively

activate specific receptors on stem cell surfaces, thereby influencing lineage commitment, while controlled nanoparticle arrangement on the cell membrane may further induce receptor clustering and trigger downstream signaling cascades. In addition, nanomaterials can regulate cell behavior through mechanotransduction by altering the local mechanical environment within organoids.⁷³

The incorporation of nanocomposite materials into ECM-based scaffolds can locally modulate stiffness and mimic dynamic mechanical changes during tissue development, whereas magnetic nanoparticles exposed to external magnetic fields can exert subtle mechanical forces on selected cell populations, thereby guiding tissue morphogenesis.⁷⁴ Nanomaterials may also influence organoid fate through redox and metabolic regulation, as certain nanoparticles have been shown to induce metabolic reprogramming in stem cells or diseased cells, while redox-active nanomaterials, such as specific metal oxides, can regulate local reactive oxygen species levels and thereby affect cell survival, proliferation, and differentiation.⁷⁵

Furthermore, unlike conventional passive diffusion, nanosystems enable spatiotemporal control of bioactive signal delivery. By responding to local cues within organoids, including pH gradients, enzyme activity, and hypoxic conditions, nanomaterials can achieve on-demand release of signaling molecules at defined developmental stages or pathological regions. At the same time, the non-uniform distribution of nanoparticles within three-dimensional structures may be exploited to mimic *in vivo* morphogen gradients, thereby promoting the establishment of more complex spatial patterning and polarity. Collectively, these mechanisms indicate that nanomaterials are not merely auxiliary delivery tools but functional regulators capable of reshaping the biochemical, biophysical, and spatiotemporal signaling environment of organoids, thus providing new opportunities for directing organoid development and enhancing the precision of disease modeling and regenerative strategies.¹⁷

6. Impact of nanomaterials on the organoid tissue microenvironment

To systematically evaluate the physicochemical properties and biological effects of nanodrugs, it is essential to construct high-fidelity models that accurately replicate the complex disease microenvironment.⁷⁶ This is because microenvironmental factors directly determine the *in vivo* behavior, therapeutic response, and safety of nanoparticles. Traditional two-dimensional monolayer cell cultures, due to their simplicity, lower cost, and suitability for high-throughput screening, have been widely used in nanomedicine research for many years (Figure 7). However, this system is primarily limited to preliminary analyses of

nanoparticle–cell interactions and cytotoxicity, failing to accurately predict their clinical efficacy.⁷⁷ The fundamental issue is that two-dimensional models lack a three-dimensional ECM, which plays a key role in regulating nanoparticle tissue penetration, diffusion kinetics, and intracellular uptake.⁷⁸

In contrast, three-dimensional organoid models, by reconstructing the natural tissue's spatial structure, cell diversity, and functional characteristics,⁷⁹ provide a more physiologically relevant experimental platform for nanodrug research. Organoids, which are typically derived from stem cells or patient tissues, can self-organize to form multicellular systems with organ-specific structures and functions, providing a microecological environment similar to that *in vivo* for preclinical studies.^{80,81} Disease-specific organoids—including models for cancer, neurodegenerative disorders, and infectious diseases—provide a pathologically relevant platform for investigating nanodrug behavior.⁸² Automated bioreactors can markedly improve the size uniformity and production throughput of organoids by precisely controlling parameters such as hydrodynamics and oxygen tension. However, standardization of the external culture environment does not necessarily translate into uniformity of nanoparticle distribution within organoids. Physical barriers persist, as the dense cellular organization and ECM inside organoids continue to restrict nanoparticle penetration even in scaled-up systems. Therefore, in large-scale organoid platforms, the penetration depth and intratissue distribution of nanoparticles should still be directly validated using three-dimensional imaging or quantitative analytical techniques to ensure the reliability of efficacy assessment.⁸³

7. Applications of nanomaterials in organoid engineering

In organoid engineering, the application of nanomaterials has become an important area of research due to their unique physicochemical properties, making them highly applicable in tumor models, tissue repair, drug screening, and multicellular organoids. Nanomaterials, with their tunable size, shape, and surface chemistry, play a significant role in organoid engineering, enhancing biocompatibility and regulating growth, development, and functionality. This section provides a detailed exploration of the application of nanomaterials in tumor organoids, tissue repair, drug screening, and their integrated role in multicellular organoids (Figure 8).

7.1. Applications in tumor organoids

The core goal of cancer nanomedicine is to construct nanocarriers that can accurately deliver drugs to the tumor site, thereby enhancing anti-tumor efficacy while




Figure 7. Organoid model-driven nanomedicine screening *in vitro*. The study provides a comparative evaluation of two-dimensional (2D) and three-dimensional (3D) organoid models for nanomedicine screening, underscoring distinctions in structural organization, cellular crosstalk, drug penetration efficiency, and predictive accuracy. The 3D organoid systems demonstrate superior physiological relevance and more reliable predictions of drug efficacy and toxicity, advantages that are further enhanced by automated screening platforms and high-throughput evaluation methodologies. Reprinted with permission from Wang *et al.*⁸⁴




Figure 8. Applications of organoid-driven platforms in nanomedicine. The application of organoid platforms in personalized nanomedicine has expanded across diverse fields, including cancer treatment, neurological diseases, infectious disorders, and tissue regeneration. Reprinted with permission from Wang *et al.*⁸⁴

Abbreviations: BBB: Blood-brain barrier; CNS: Central nervous system; PDO: Patient-derived organoid.

minimizing damage to normal tissues.⁸⁵ Tumor organoids, derived from patient biopsy specimens or tumor cell lines, serve as physiologically relevant three-dimensional platforms for investigating nanoparticle behavior within the complex and dynamic tumor microenvironment. By maintaining critical features of the parental tumor, including cellular heterogeneity, epithelial–mesenchymal interactions, and tumor-specific ECM components, these models offer an advantageous platform for evaluating the delivery performance, tissue distribution, and therapeutic efficacy of nanomedicines.⁸⁶

Beyond the well-established pathway of apoptosis, emerging modalities of programmed cell death, including ferroptosis and cuproptosis, have recently attracted considerable interest within cancer research and nanomedicine.⁸⁷ Nanomaterials are capable of initiating these cell death pathways through the modulation of iron and copper homeostasis, the induction of lipid peroxidation, and the instigation of mitochondrial proteotoxic stress, offering a promising strategy to circumvent tumor resistance and augment anti-tumor efficacy.⁵⁴ Tumor organoid models, which retain three-dimensional tissue architecture, cellular heterogeneity, and key elements of the tumor microenvironment, represent a physiologically relevant *in vitro* platform for assessing these nanomaterial-induced cell death mechanisms, thereby refining and expanding our comprehension of nanomaterial-mediated anti-tumor actions.⁸⁸

By reconstructing the complexity of tumor tissue in terms of spatial structure and biochemical signals, tumor organoids provide crucial tools for studying the interactions between nanoparticles, cancer cells, and ECM components, helping deepen understanding of drug uptake mechanisms, intracellular transport pathways, and local therapeutic responses.⁸⁹ In solid tumors, the hypoxic core is of particular importance because it is marked by low oxygen levels, dense ECM deposition, and poor vascular perfusion, all of which pose substantial obstacles to drug penetration.⁹⁰ Tumor organoids can faithfully recapitulate oxygen and nutrient gradients, providing a reliable model for assessing the capacity of nanoparticles to penetrate and exert therapeutic effects within hypoxic regions.⁹¹

Although patient-derived organoids hold considerable promise for optimizing nanoformulations in a personalized manner, their broader application remains constrained by several practical factors.⁹² First, resource and time limitations are major challenges, as access to primary tumor tissues is often uncertain, and the establishment and expansion of organoids can be time-consuming, which may not be compatible with the urgent therapeutic window of some patients with advanced cancer. Second, the cost and scalability of individualized screening remain substantial barriers, as this approach requires intensive culture

and analytical resources, while standardized automated workflows for processing large numbers of patient samples are still lacking. Finally, successful clinical translation requires the establishment of standardized procedures covering tissue processing, nanoparticle administration protocols, and efficacy evaluation criteria, in order to ensure the reproducibility of predictive results and their clinical relevance.⁹³

7.2. Applications in tissue repair

In regenerative medicine research, nanoparticles have gradually become an important tool for promoting tissue repair and functional reconstruction, primarily by delivering growth factors, regulating local microenvironments, and targeting stem cell populations to enhance tissue regeneration.⁹⁴ Organoid models can reconstruct the spatial structure, cell composition, and some functional characteristics of natural tissues, providing a highly relevant three-dimensional experimental platform for evaluating the biological effects of nanomaterials in the regeneration process.⁹⁵

Organoids derived from tissues such as the liver, kidney, and skin provide suitable platforms for functionally validating nanocarriers intended for tissue repair applications.⁹⁶ These models can simulate tissue microenvironments under damaged or pathological conditions. On this basis, nanoparticles loaded with growth factors—such as vascular endothelial growth factor (VEGF) and epidermal growth factor (EGF)—can be used to assess their ability to promote cell proliferation, induce angiogenesis, and drive tissue remodeling in organoid systems, particularly in the context of fibrosis or chronic injury.⁹⁷

7.3. Applications in drug screening

In drug screening, nanomaterials serve not only as therapeutic carriers but also as essential tools for constructing efficient screening systems. By loading antiviral, anti-malaria, or immune-modulating molecules onto nanocarriers, their delivery efficiency, cellular uptake capacity, and therapeutic potential can be systematically evaluated *in vitro*, enabling more targeted drug screening strategies.⁹⁸ However, the clinical translation of organoid-based nanodrug screening remains incompletely validated. Although organoid-based studies of drug response have shown some correlation with patient treatment outcomes, direct clinical validation specifically for nanodrug screening remains limited. Most existing evidence has been derived from studies of conventional small molecules or chemotherapeutic agents, whereas nanocarrier-based systems, owing to their more complex delivery mechanisms and *in vivo* behaviors, have not yet been sufficiently validated in terms of how organoid responses correspond to clinical outcomes.⁹⁹ In contrast to

traditional two-dimensional cell culture models, organoids derived from tissues such as the lung and intestine provide physiologically relevant three-dimensional platforms for nanodrug screening and more accurately recapitulate host–pathogen interactions in native tissues.¹⁰⁰

Organoids can reconstruct tissue-specific structures and functions, including epithelial barrier integrity, mucus secretion, and immune response patterns, thus forming infection models that closely mimic the *in vivo* environment.¹⁰¹ For example, lung organoids have been widely used to evaluate nanodrug candidates targeting respiratory pathogens, such as the influenza virus, respiratory syncytial virus, and SARS-CoV-2.¹⁰² In these systems, nanoparticles loaded with small-molecule drugs, peptides, or small interfering RNA (siRNA) can be used to address targeting efficiency, viral replication inhibition, and immune-modulating effects, thereby optimizing nanomedicine design.

Moreover, organoid models can reproduce key microenvironmental factors in infected tissues, such as pathogen storage regions, hypoxic conditions, and immune cell involvement, thereby aligning nanomaterial screening more closely with clinical scenarios.¹⁰³ For example, nanoparticles with surface-modified specific ligands can be validated in organoids to determine their selective binding to infected cells or pathogens, thereby selecting candidates with superior targeting ability.¹⁰⁴ Additionally, the multicellular dynamic environment helps analyze immune responses induced by nanoparticles, such as changes in cytokine secretion and macrophage activation levels, providing a basis for comprehensive evaluation of drug safety and efficacy.¹⁰⁵

At present, organoid-based nanotoxicology research is primarily focused on acute toxicity, while the evaluation of long-term effects is still in its early stages. To improve clinical translational value, future studies should prioritize the following: chronic and repeated dosing designs, including the establishment of long-term exposure models that simulate clinical treatment cycles to assess the cumulative toxicity of nanoparticles within organoids; functional and genomic stability, with attention extending beyond cell viability to the effects of prolonged exposure on organoid physiological functions, such as barrier integrity, as well as on genomic stability; and dynamic monitoring technologies, such as organ-on-a-chip platforms, to enable real-time and dynamic monitoring of long-term interactions between nanoparticles and organoids.¹⁰⁶

Organoid-driven nanomedicine platforms hold enormous promise; however, most current organoid systems still lack complex immune components, greatly limiting the comprehensive evaluation of nanoparticle immunogenicity and immune-mediated therapeutic

effects.¹⁰⁷ To address this limitation, research is evolving toward a higher level of physiological complexity. One key challenge is the limitation in immunogenicity evaluation: the absence of endogenous immune cells makes it difficult for researchers to observe critical biological responses induced by nanoparticles, such as cytokine storms, macrophage polarization, and complement activation. Immune cell co-culture strategies offer one potential solution: by introducing peripheral blood mononuclear cells, T cells, or macrophages into organoid systems, it becomes possible to simulate immune infiltration within the tumor microenvironment and thereby more accurately assess the cytotoxic efficacy of immuno-nanomedicines.¹⁰⁸ Another important approach is the organoid-on-a-chip strategy: microfluidic-based organoid chip systems can simulate dynamic blood circulation and immune cell recruitment, providing a valuable tool for investigating the real-time interactions between nanoparticles and the immune system. Through the integration of these strategies, organoid models will more faithfully reflect the immune response mechanisms of nanomedicines *in vivo*, thereby significantly enhancing the translational value of preclinical research.

Based on three-dimensional organoid systems, the nanodrug screening platform can more accurately reflect the *in vivo* microenvironment compared to traditional two-dimensional cultures, improving the predictive accuracy of nanomedicine performance and reducing the gap between *in vitro* screening and clinical application.¹⁰⁹ Therefore, the combination of nanomaterials and organoid technology is becoming a key strategy for precise drug screening and optimization in anti-infective drug development.¹¹⁰

7.4. Applications in multi-cell line organoids

Three-dimensional multicellular organoids, as highly biomimetic *in vitro* models, can reconstruct the spatial structure, cell heterogeneity, and microenvironment characteristics of tissues or tumors. Compared to traditional two-dimensional cell culture systems, they offer greater physiological relevance and clinical predictive value in nanomaterial research. Multicellular organoids typically include cancer cells, fibroblasts, immune cells, and ECM components, enabling the simulation of cell–cell and cell–matrix interactions that occur in real tissues.¹¹¹

In multicellular organoids, different cell types exhibit differential nanoparticle uptake behaviors. For example, in tumor organoids, cancer cells, stromal fibroblasts, and infiltrating immune cells show significant differences in their uptake patterns.¹¹² This variation provides important insights for the precise design of targeted nanodrugs. Organoids reflect the complex tissue microenvironment, including the ECM, oxygen gradient, nutrient gradient, and spatial structural barriers, all of which influence

nanoparticle distribution and penetration.¹¹³ Therefore, organoids can more accurately reflect the transport behavior of nanodrugs *in vivo*. Nanoparticle size, surface charge, and functional modifications significantly affect their diffusion, penetration, and retention within organoids.¹¹⁴ Research has shown that negatively charged nanoparticles generally exhibit better tissue penetration in organoids, and multivalent ligand or targeting molecule modifications can enhance their endocytosis efficiency in specific cell populations.¹¹⁴

Overall, multicellular organoids, by reconstructing complex tissue microenvironments and cellular heterogeneity, provide a highly biomimetic research platform for evaluating nanoparticle uptake, distribution, and therapeutic efficacy in biological systems.⁷⁷ However, it is important to clarify that current organoid models only partially reproduce the intricate stromal and vascular interactions found *in vivo*. Although the integration of multicellular organoids, vascularized platforms, and microfluidic technologies represents a major advance in bioengineering and has substantially improved the physiological relevance of these models,¹¹⁵ it must be acknowledged that current organoid systems can only partially reproduce the complex stromal and vascular interactions present *in vivo*. First, with regard to stromal complexity, although co-culture with fibroblasts and immune cells has enhanced cellular heterogeneity, these systems still struggle to fully recapitulate the highly dynamic and hierarchically organized ECM architecture of native tissues.¹¹⁶ Second, in terms of vascularization, most current vascularized organoids rely on the integration of endothelial cells or on artificial scaffolds. While these approaches can improve nutrient and oxygen delivery to some extent, they remain insufficient compared with native vascular networks in terms of branching density, hemodynamic shear stress, and the fine microstructure of capillary beds. Third, microfluidic organ-on-a-chip platforms are capable of simulating certain hydrodynamic conditions and partially compensating for the limitations of traditional static culture systems; however, they still face significant technical challenges in maintaining long-term vascular stability and in reconstructing complex intertissue crosstalk. Overall, despite the substantial progress enabled by these advanced platforms, a considerable gap remains between current models and the full complexity of native stromal and vascular systems, and bridging this gap will be a critical priority for future translational research.¹¹⁷ Their applications in mechanism studies, immunotherapy collaboration research, high-throughput screening, and precision medicine significantly enhance the clinical translation potential of nanomedicines, making the intersection of nanomedicine and organoids a key area of research development.¹¹⁸

However, this increase in complexity also poses serious challenges for experimental standardization, and the establishment of unified criteria covering the following parameters is a prerequisite for clinical translation. To improve interlaboratory reproducibility and enhance the translational potential of organoid-based studies, the standardization of experimental parameters is essential.¹¹⁹ The following summarizes the core elements that should be standardized in studies of nanoparticle–organoid interactions: organoid characterization, including detailed documentation of organoid origin (such as patient biopsies and stem cell lines), passage number, culture duration, and quantitative size distribution to ensure baseline consistency of the biological model; microenvironmental control, including clear definition of the chemical composition, stiffness, and matrix components of the hydrogel or extracellular scaffold, as these physical properties strongly influence nanoparticle penetration kinetics and distribution heterogeneity; nanoparticle physicochemical properties, including precise characterization of particle size, surface charge, morphology, and surface modifications such as ligand density, since these features determine tissue diffusion and cellular uptake patterns; exposure protocols, including standardized dosing regimens based on mass concentration or surface area concentration, exposure duration, and the specific administration mode, such as static incubation or dynamic microfluidic perfusion; and analytical endpoints, including standardized quantitative metrics such as radial penetration depth, core-to-periphery distribution ratio, single-cell uptake rate, and absolute accumulation measured by ICP-MS.¹²⁰

8. Challenges and future directions

Despite the promising potential of organoid-driven nanomedicine platforms, there are still many challenges in their biomedical research and therapeutic applications that require systematic optimization.^{121–122} The foremost issue is the lack of a unified standard system.¹²³ Beyond insufficient standardization, batch-to-batch variability commonly observed in organoid systems is another major factor affecting the consistency of experimental outcomes.¹⁰⁶ Even under identical culture conditions, substantial differences may still arise between batches in organoid size, cellular composition, differentiation state, and ECM composition. In addition, the intratumoral-like microenvironment within organoids is highly heterogeneous, with pronounced spatial gradients in oxygen, nutrients, and metabolic byproducts. These factors can markedly influence nanoparticle uptake efficiency, penetration capacity, and subsequent biological responses.¹²⁴ For example, larger organoids usually exhibit a more pronounced diffusion barrier effect, thereby limiting nanoparticle access to the core region, whereas cells at different differentiation states may display

distinct nanoparticle uptake profiles, further increasing result variability. Accordingly, batch-to-batch variability has been recognized as one of the major sources of poor reproducibility in nanoparticle–organoid studies. At the current stage, effectively reducing inter-batch variability and establishing standardized evaluation frameworks remain critical challenges for the field. Future work should therefore place greater emphasis on the standardization of organoid preparation procedures, harmonization of experimental parameters, and incorporation of quantitative assessment methods to improve the reliability and comparability of experimental results.

In addition, although nanomaterials have shown considerable potential in regulating metabolic reprogramming in organoid systems, the reproducibility of these effects across different organoid models has not yet been sufficiently validated. Most existing studies have focused on specific models or single experimental settings, and no unified or broadly generalizable conclusions have yet been established. Therefore, this area remains largely exploratory, and the robustness and reproducibility of current findings still require systematic evaluation.

Available evidence suggests that nanomaterial-induced metabolic reprogramming is strongly influenced by multiple factors.¹²⁵ First, the origin of the organoids, such as whether they are patient-derived or stem cell-derived, as well as their differentiation status, can directly shape their metabolic features and responsiveness to external stimuli. Second, the physicochemical properties of the nanomaterials, including composition, particle size, surface charge, and surface modification, can alter their interactions with cells and thereby affect their metabolic regulatory capacity.¹²⁶ Third, experimental conditions, such as dosage and exposure duration, also play important roles in determining the magnitude and direction of metabolic reprogramming. Consequently, at present, the consistency of nanomaterial-mediated metabolic modulation across different organoid systems remains highly uncertain.

This issue is now regarded as a major limitation of the field. Future studies will need to improve reproducibility and comparability by standardizing organoid construction workflows, unifying nanomaterial characterization parameters, and optimizing experimental design, thereby supporting more reliable mechanistic investigations and facilitating clinical translation.¹²⁷ Currently, there are no general guidelines for organoid culture, maintenance, and functional characterization, leading to significant differences in size, cell composition, and functional characteristics across different laboratories.^{128,129} This inter-laboratory variability weakens the reproducibility of models and affects their reliability as preclinical evaluation tools. Therefore, establishing standardized culture medium formulations, quality control systems, and functional

verification standards is essential for enhancing model stability and comparability.¹³⁰

Another major limitation is the insufficient level of clinical validation. Most current studies remain at the stage of preliminary toxicity and efficacy assessment, and the lack of large-scale clinical data makes it difficult to accurately predict the actual *in vivo* performance of these systems in patients. To validate the predictive accuracy of organoid models for patient clinical responses, researchers should adopt a systematic validation framework.¹³¹ First, matched testing should be performed using paired patient-derived organoids to evaluate both free drugs and their corresponding nanoformulations, thereby assessing the therapeutic gain conferred by nanomedicine strategies.¹³² Second, clinical correlation should be established by directly comparing *in vitro* organoid screening results with patients' actual clinical outcomes, either retrospectively or prospectively, in order to determine the predictive sensitivity and specificity of the model. Third, multi-parameter readouts should be integrated to encompass diverse indicators, including cell viability, morphological changes, gene expression profiles, and the tissue penetration depth of nanoparticles, to construct a comprehensive evaluation system. Finally, the clinical predictive value of organoid-based nanomedicine platforms should be confirmed through longitudinal patient follow-up and multicenter validation across laboratories.

Another major limitation is the lack of scalable production capacity.¹³³ At present, the organoid construction process requires high precision, involving directed differentiation of stem cells and fine-tuning of three-dimensional culture systems. The overall process is time-consuming and labor-intensive.¹³⁴ Although organoid models overcome some of the limitations of conventional two-dimensional cell models, their ability to bridge the gap between preclinical findings and clinical outcomes remains limited. The major advantage of organoid systems lies in their capacity to preserve three-dimensional tissue architecture, cellular heterogeneity, and disease-relevant microenvironmental features, thereby providing a more physiologically relevant platform for the *in vitro* evaluation of nanomedicines.¹³⁵ However, these models cannot fully recapitulate the complex systemic processes that occur *in vivo*, including pharmacokinetics, immune regulation, and inter-organ interactions. Therefore, while organoids may help narrow the gap between preclinical success and clinical failure to a certain extent, they cannot yet replace *in vivo* models and have significant limitations in predicting the clinical efficacy of nanomedicines.¹³⁶ Small fluctuations in oxygen tension, nutrient gradients, or fluid dynamics can induce heterogeneity among batches, making it difficult to ensure batch consistency.¹³⁷ This limitation restricts its application in high-throughput drug screening, where large

numbers of standardized and stable biological models are needed.¹³⁸

To enhance scalability, automated bioreactors and microfluidic systems are considered key technological pathways. Studies have shown that automated stirred-tank bioreactors, by precisely controlling fluid dynamics, stirring rate, and oxygen tension, can stably produce billions of human-induced pluripotent stem cell aggregates of controllable size while maintaining their pluripotent state.¹³⁹ Additionally, modular organ-on-a-chip platforms show great promise. For example, Carvalho *et al.*⁹⁶ developed a LEGO-like microfluidic system that combines continuous perfusion with real-time monitoring of DO₂ and pH, enabling the cultivation of thyroid organoids for up to 10 days under strictly controlled conditions. These technologies combine automated scaling with precise environmental control, laying the foundation for the industrial-scale, high-quality, reproducible production of organoids.

9. Conclusion

Organoid-driven nanomedicine platforms, by addressing key issues such as insufficient standardization, limited scalability, and incomplete physiological complexity, have the potential to transition from experimental research tools to predictive preclinical models. These technological advancements will aid in the realization of personalized, precise, and highly safe therapeutic strategies, promoting the translation of preclinical research findings into clinical applications.

Organoid technology provides a more in vivo-like experimental platform for biomedical research, showing immense potential in drug screening, disease model establishment, and therapeutic method exploration. The application of nanomaterials in organoids will propel the field to new heights, offering new strategies for drug targeting and delivery, while also providing a solid foundation for the study of disease mechanisms and the development of precision medicine.⁵⁶ However, despite significant progress in both organoid and nanotechnology fields, challenges such as model standardization, scalability, and physiological complexity remain. Addressing these issues will drive deeper integration of organoid and nanomaterial research and promote their widespread application in biomedical fields.

In the future, as technology continues to advance, the combination of organoids and nanomaterials will enable breakthroughs in multiple directions. Firstly, the vascularization of organoids will become a key focus of research, particularly through the integration of endothelial cell co-culture systems, artificial vascular

scaffolds, and microfluidic perfusion systems to support nanoparticle penetration and therapeutic efficacy in highly vascularized tissues. Secondly, nanodrug design will focus on microenvironment-responsive features, such as developing multifunctional nanoparticles that respond to hypoxia or pH changes to enhance targeting precision and reduce nonspecific side effects. Within the three-dimensional architecture of organoids, particularly in tumor organoids, pronounced hypoxic and redox gradients are commonly present. It should be clarified that nanoparticles do not biologically “adapt” to these changes on their own; rather, they can be molecularly engineered to exhibit physicochemical responsiveness to specific microenvironmental cues. In terms of hypoxia responsiveness, nanomaterials can be designed with hypoxia-sensitive chemical bonds or functional groups, enabling precise drug release or structural transformation within the hypoxic core regions of organoids and thereby helping to overcome the penetration barriers imposed by dense tissues. In addition, redox-sensitive designs can take advantage of variations in intracellular glutathione levels and other redox-related indicators within organoids, allowing engineered nanocarriers to undergo spatiotemporally controlled disassembly and thus improving the local bioavailability of therapeutic agents. Therefore, the key concept is not active adaptation, but intelligent sensing through environment-triggered mechanisms. This strategy provides a precise means of regulation in organoid research and enables the design of more predictive and pathology-oriented drug-delivery systems for complex microenvironments. Moreover, the construction of multi-organ models will provide more comprehensive data for systematic efficacy and safety studies, especially when combined with immune components and vascularized structures, significantly improving clinical translation potential.

In summary, with technological advancements and increasing interdisciplinary collaboration, organoid-driven nanomedicine platforms are expected to provide new solutions for personalized medicine and precision treatments, accelerating the development and clinical application of innovative drugs and ultimately benefiting human health.

Acknowledgments

None.

Funding

None.

Conflict of interest

The authors declare they have no competing interests.

Author contributions

Conceptualization: Wenguo Cui, Hongbo Zhang

Visualization: Zihang Zhao

Writing—original draft: Shuo Zhang, Xu Chu

Writing—review & editing: Guilai Zuo

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data

Not applicable.

References

- Schutgens F, Clevers H. Human organoids: Tools for understanding biology and treating diseases. *Annu Rev Pathol.* 2020;15:211-234.
doi: 10.1146/annurev-pathmechdis-012419-032611
- Lancaster MA, Knoblich JA. Organogenesis in a dish: modeling development and disease using organoid technologies. *Science.* 2014;345(6194):1247125.
doi: 10.1126/science.1247125
- Clevers H. Modeling development and disease with organoids. *Cell.* 2016;165(7):1586-1597.
doi: 10.1016/j.cell.2016.05.082
- Kapałczyńska M, Kolenda T, Przybyła W, et al. 2D and 3D cell cultures - a comparison of different types of cancer cell cultures. *Arch Med Sci.* 2018;14(4):910-919.
doi: 10.5114/aoms.2016.63743
- Zhou LB, Lei YT, Han X. Charting bioethical frontiers: China's human organoid guidelines in a global context. *Mil Med Res.* 2025;12(1):64.
doi: 10.1186/s40779-025-00651-x
- Abdel Fattah AR, Ranga A. Nanoparticles as versatile tools for mechanotransduction in tissues and organoids. *Front Bioeng Biotechnol.* 2020;8:240.
doi: 10.3389/fbioe.2020.00240
- Fei K, Zhang J, Yuan J, Xiao P. Present application and perspectives of organoid imaging technology. *Bioengineering (Basel).* 2022;9(3):121.
doi: 10.3390/bioengineering9030121
- Pomerantseva E, Bonaccorso F, Feng X, Cui Y, Gogotsi Y. Energy storage: The future enabled by nanomaterials. *Science.* 2019;366(6468):eaan8285.
doi: 10.1126/science.aan8285
- Shen C, Zhang ZJ, Li XX, et al. Intersection of nanomaterials and organoids technology in biomedicine. *Front Immunol.* 2023;14:1172262.
doi: 10.3389/fimmu.2023.1172262
- Mahapatra C, Lee R, Paul MK. Emerging role and promise of nanomaterials in organoid research. *Drug Discov Today.* 2022;27(3):890-899.
doi: 10.1016/j.drudis.2021.11.007
- Pampaloni F, Reynaud EG, Stelzer EH. The third dimension bridges the gap between cell culture and live tissue. *Nat Rev Mol Cell Biol.* 2007;8(10):839-845.
doi: 10.1038/nrm2236
- Badir A, Refki S, Sekkat Z. Utilizing gold nanoparticles in plasmonic photothermal therapy for cancer treatment. *Heliyon.* 2025;11(4):e42738.
doi: 10.1016/j.heliyon.2025.e42738
- Narayan R, Nayak UY, Raichur AM, Garg S. Mesoporous silica nanoparticles: A comprehensive review on synthesis and recent advances. *Pharmaceutics.* 2018;10(3):118.
doi: 10.3390/pharmaceutics10030118
- Vallet-Regí M, Schüth F, Lozano D, Colilla M, Manzano M. Engineering mesoporous silica nanoparticles for drug delivery: Where are we after two decades? *Chem Soc Rev.* 2022;51(13):5365-5451.
doi: 10.1039/d1cs00659b
- Miyamoto Y, Koshidaka Y, Murase K, Kanno S, Noguchi H, Miyado K, et al. Functional evaluation of 3D liver models labeled with polysaccharide functionalized magnetic nanoparticles. *Materials.* 2022;15(21):7823.
doi: 10.3390/ma15217823
- Choudhery MS, Umezawa A. Stem cells as biological drugs for incurable diseases. *Stem Cell Res Ther.* 2025;16(1):482.
doi: 10.1186/s13287-025-04632-8
- Shi Y, Han X, Zou S, Liu G. Nanomaterials in organoids: From interactions to personalized medicine. *ACS Nano.* 2024;18(49):33276-33292.
doi: 10.1021/acsnano.4c13330
- Pérez-Piñero J, Sánchez-Cea F, Arce MP, et al. Stability study of graphene oxide-bovine serum albumin dispersions. *J Xenobiot.* 2023;13(1):90-101.
doi: 10.3390/jox13010008
- Shah S, Dhawan V, Holm R, Nagarsenker MS, Perrie Y. Liposomes: Advancements and innovation in the manufacturing process. *Adv Drug Deliv Rev.* 2020;154-155:102-122.
doi: 10.1016/j.addr.2020.07.002
- Lin L, Song X, Dong X, Li B. Nano-photosensitizers for enhanced photodynamic therapy. *Photodiagnosis Photodyn Ther.* 2021;36:102597.

- doi: 10.1016/j.pdpdt.2021.102597
21. Wang H, Chen Y, Wei R, *et al.* Synergistic chemioimmunotherapy augmentation via sequential nanocomposite hydrogel-mediated reprogramming of cancer-associated fibroblasts in osteosarcoma. *Adv Mater.* 2024;36(15):e2309591.
doi: 10.1002/adma.202309591
 22. El Moukhtari SH, Garbayo E, Amundarain A, *et al.* Lipid nanoparticles for siRNA delivery in cancer treatment. *J Control Release.* 2023;361:130-146.
doi: 10.1016/j.jconrel.2023.07.054
 23. Dutta Gupta Y, Mackeyev Y, Krishnan S, Bhandary S. Mesoporous silica nanotechnology: promising advances in augmenting cancer theranostics. *Cancer Nanotechnol.* 2024;15(1):9.
doi: 10.1186/s12645-024-00250-w
 24. Viegas C, Patrício AB, Prata J, *et al.* Advances in pancreatic cancer treatment by nano-based drug delivery systems. *Pharmaceutics.* 2023;15(9):2363.
doi: 10.3390/pharmaceutics15092363
 25. Chiu JZ, Tucker IG, McLeod BJ, McDowell A. Arginine-tagging of polymeric nanoparticles via histidine to improve cellular uptake. *Eur J Pharm Biopharm.* 2015;89:48-55.
doi: 10.1016/j.ejpb.2014.11.014
 26. Dilliard SA, Siegwart DJ. Passive, active and endogenous organ-targeted lipid and polymer nanoparticles for delivery of genetic drugs. *Nat Rev Mater.* 2023;8(4):282-300.
doi: 10.1038/s41578-022-00529-7
 27. Huang H, Chang A, Peng H, *et al.* Preparation and anti-tumor effect in hepatocellular carcinoma treatment of AS1411 aptamer-targeted polyphyllin II-loaded PLGA nanoparticles. *J Sci Adv Mater Devices.* 2024;9(3):100755.
doi: 10.1016/j.jsamd.2024.100755
 28. Shaw I, Aryee AA, Ali YS, *et al.* Harmonizing tradition and technology: Liposomal nanocarriers unlocking the power of natural herbs in Traditional Chinese Medicine. *Chin J Nat Med.* 2025;23(6):700-713.
doi: 10.1016/s1875-5364(25)60889-2
 29. Mitchell MJ, Billingsley MM, Haley RM, Wechsler ME, Peppas NA, Langer R. Engineering precision nanoparticles for drug delivery. *Nat Rev Drug Discov.* 2021;20(2):101-124.
doi: 10.1038/s41573-020-0090-8
 30. Omlor AJ, Nguyen J, Bals R, Dinh QT. Nanotechnology in respiratory medicine. *Respir Res.* 2015;16(1):64.
doi: 10.1186/s12931-015-0223-5
 31. Barroso L, Viegas C, Vieira J, Ferreira-Pêgo C, Costa J, Fonte P. Lipid-based carriers for food ingredients delivery. *J Food Eng.* 2021;295:110451.
doi: 10.1016/j.jfoodeng.2020.110451
 32. Bag J, Banerjee S, De A, *et al.* Nanoengineered approaches to improve the efficacy of targeted drug delivery for the treatment of malignancy: a comprehensive review. *Futur J Pharm Sci.* 2023;9(1):88.
doi: 10.1186/s43094-023-00541-w
 33. López-Estevez AM, Gref R, Alonso MJ. A journey through the history of PEGylated drug delivery nanocarriers. *Drug Deliv Transl Res.* 2024;14(8):2026-2031.
doi: 10.1007/s13346-024-01608-8
 34. Vakhshiteh F, Bagheri Z, Soleimani M, *et al.* Heterotypic tumor spheroids: a platform for nanomedicine evaluation. *J Nanobiotechnology.* 2023;21(1):249.
doi: 10.1186/s12951-023-02021-y
 35. Lacin ME, Yildirim M. Applications of multiphoton microscopy in imaging cerebral and retinal organoids. *Front Neurosci.* 2024;18:1360482.
doi: 10.3389/fnins.2024.1360482
 36. Zhao WP, Wang HW, Liu J, *et al.* Positive PCNA and Ki-67 expression in the testis correlates with spermatogenesis dysfunction in fluoride-treated rats. *Biol Trace Elem Res.* 2018;186(2):489-497.
doi: 10.1007/s12011-018-1338-6
 37. Mohammed MA, Syeda JTM, Wasan KM, Wasan EK. An overview of chitosan nanoparticles and its application in non-parenteral drug delivery. *Pharmaceutics.* 2017;9(4):53.
doi: 10.3390/pharmaceutics9040053
 38. Alves-Lopes JP, Söder O, Stukenborg JB. Use of a three-layer gradient system of cells for rat testicular organoid generation. *Nat Protoc.* 2018;13(2):248-259.
doi: 10.1038/nprot.2017.140
 39. Kita K, Watanabe T, Ohsaka K, *et al.* Production of functional spermatids from mouse germline stem cells in ectopically reconstituted seminiferous tubules. *Biol Reprod.* 2007;76(2):211-217.
doi: 10.1095/biolreprod.106.056895
 40. Dobrinski I. De novo morphogenesis of functional testis tissue after ectopic transplantation of isolated cells. *Organogenesis.* 2007;3(2):79-82.
doi: 10.4161/org.3.2.4944
 41. Sato T, Clevers H. Growing self-organizing mini-guts from a single intestinal stem cell: mechanism and applications. *Science.* 2013;340(6137):1190-1194.
doi: 10.1126/science.1234852
 42. Proceedings of the 17th Italian convention of FFC investigators in cystic fibrosis. *BMC Pulm Med.* 2020;20(1):100.
doi: 10.1186/s12890-020-1104-3
 43. Yang T, Wang YY, Ma F, Xu BH, Qian HL. Build the virtual cell with artificial intelligence: a perspective for cancer research. *Mil Med Res.* 2025;12(1):4.

doi: 10.1186/s40779-025-00591-6

44. Kaur SD, Kapoor DN. Cancer in translation: cyanine carborane salts: a multifunctional theranostic approach to cancer treatment. *EJNMMI Radiopharm Chem.* 2025;10(1):36.
doi: 10.1186/s41181-025-00360-9
45. van Ravensteijn BGP, Vermonden T. Dynamic, tunable, biocompatible: Functional polymers advancing biomedical innovation. *Biomacromolecules.* 2026;27(1):1-2.
doi: 10.1021/acs.biomac.5c02557
46. Turpin I, Modrego A, Sarrias AM, *et al.* Deep learning applied to live imaging for classifying brain organoids derived from neurodevelopmental disorders considering the intrinsic variability of the in vitro model. *IBRO Neurosci Rep.* 2023;15:S209.
doi: 10.1016/j.ibneur.2023.08.331
47. Singh A, Lutolf M. Breaking the barriers in engineering organoids and tissues with advanced materials. *Adv Funct Mater.* 2020;30(48):2008531.
doi: 10.1002/adfm.202008531
48. Robinson M, Sparanese S, Witherspoon L, Flannigan R. Human in vitro spermatogenesis as a regenerative therapy - where do we stand? *Nat Rev Urol.* 2023;20(8):461-479.
doi: 10.1038/s41585-023-00723-4
49. Shankaran A, Prasad K, Chaudhari S, Brand A, Satyamoorthy K. Advances in development and application of human organoids. *3 Biotech.* 2021;11(6):257.
doi: 10.1007/s13205-021-02815-7
50. Takasato M, Er PX, Chiu HS, *et al.* Kidney organoids from human iPS cells contain multiple lineages and model human nephrogenesis. *Nature.* 2015;526(7574):564-568.
doi: 10.1038/nature15695
51. Karkampouna S, La Manna F, Benjak A, *et al.* Patient-derived xenografts and organoids model therapy response in prostate cancer. *Nat Commun.* 2021;12(1):1117.
doi: 10.1038/s41467-021-21300-6
52. Scholler J, Groux K, Goureau O, *et al.* Dynamic full-field optical coherence tomography: 3D live-imaging of retinal organoids. *Light Sci Appl.* 2020;9:140.
doi: 10.1038/s41377-020-00375-8
53. Bruun J, Kryeziu K, Eide PW, *et al.* Patient-derived organoids from multiple colorectal cancer liver metastases reveal moderate intra-patient pharmacotranscriptomic heterogeneity. *Clin Cancer Res.* 2020;26(15):4107-4119.
doi: 10.1158/1078-0432.CCR-19-3637
54. Dorbala S, Ando Y, Bokhari S, *et al.* ASNC/AHA/ASE/EANM/HFSA/ISA/SCMR/SNMMI Expert consensus recommendations for multimodality imaging in cardiac amyloidosis: Part 1 of 2-Evidence base and standardized methods of imaging. *Circ Cardiovasc Imaging.* 2021;14(7):e000029.
doi: 10.1161/HCI.0000000000000029
55. Xiang X, Pang H, Ma T, *et al.* Ultrasound targeted microbubble destruction combined with Fe-MOF based bio-/enzyme-mimics nanoparticles for treating of cancer. *J Nanobiotechnology.* 2021;19(1):92.
doi: 10.1186/s12951-021-00835-2
56. Achilefu S. Introduction to concepts and strategies for molecular imaging. *Chem Rev.* 2010;110(5):2575-2578.
doi: 10.1021/cr1001113
57. Yang J, Lei W, Xiao Y, *et al.* Generation of human vascularized and chambered cardiac organoids for cardiac disease modelling and drug evaluation. *Cell Prolif.* 2024;57(8):e13631.
doi: 10.1111/cpr.13631
58. Shi R, Li Y, Liu H, Chen J. Lung microbiota: Unexploited treasure hidden in the immune microenvironment of lung cancer. *Thorac Cancer.* 2021;12(21):2964-2966.
doi: 10.1111/1759-7714.14159
59. Taylor LS, Bergström CAS, Lavasanifar A, Qian F, Suryanarayanan R, Thurecht KJ. Celebrating women in the pharmaceutical sciences. *Mol Pharm.* 2021;18(4):1487-1490.
doi: 10.1021/acs.molpharmaceut.0c01197
60. Chen N, Brachmann C, Liu X, *et al.* Albumin-bound nanoparticle (nab) paclitaxel exhibits enhanced paclitaxel tissue distribution and tumor penetration. *Cancer Chemother Pharmacol.* 2015;76(4):699-712.
doi: 10.1007/s00280-015-2833-5
61. Zhang D, Jin W, Cui Y, He Z. Establishment and characterization of testis organoids with proliferation and differentiation of spermatogonial stem cells into spermatocytes and spermatids. *Cells.* 2024;13(19):1642.
doi: 10.3390/cells13191642
62. Metselaar JM, Lammers T. Challenges in nanomedicine clinical translation. *Drug Deliv Transl Res.* 2020;10(3):721-725.
doi: 10.1007/s13346-020-00740-5
63. Miller MA, Gadde S, Pfirschke C, *et al.* Predicting therapeutic nanomedicine efficacy using a companion magnetic resonance imaging nanoparticle. *Sci Transl Med.* 2015;7(314):314ra183.
doi: 10.1126/scitranslmed.aac6522
64. Pan B, Wei X, Xu X. Patient-derived xenograft models in hepatopancreatobiliary cancer. *Cancer Cell Int.* 2022;22(1):41.
doi: 10.1186/s12935-022-02454-9
65. Heo JH, Kang D, Seo SJ, Jin Y. Engineering the extracellular matrix for organoid culture. *Int J Stem Cells.* 2022;15:60-69.
doi: 10.15283/ijsc21190

66. Chen X. Biomedical poly(amino acid) materials. *Nano Lett.* 2026;26(11):3645-3648.
doi: 10.1021/acs.nanolett.6c00156
67. Chen X. From quantum dots to cancer nanotheranostics. *Nano Lett.* 2025;25(38):13967-13969.
doi: 10.1021/acs.nanolett.5c03619
68. Mao Y, Li C, Fan C, Schanze KS, Fernandez L. Forum on materials for brain research. *ACS Appl Mater Interfaces.* 2023;15(1):1-2.
doi: 10.1021/acsami.2c21906
69. Singh LS, Singh CS, Premata T. Magnetically controlled nanocarriers for site-specific drug delivery and theranostics in cancer precision medicine. *Mol Imaging Biol.*
doi: 10.1007/s11307-026-02097-6
70. Becker ML, Burdick JA. Introduction: Polymeric biomaterials. *Chem Rev.* 2021;121(18):10789-10791.
doi: 10.1021/acs.chemrev.1c00354
71. Sridhar SS, Blais N, Tran B, *et al.* Efficacy and safety of nab-paclitaxel vs paclitaxel on survival in patients with platinum-refractory metastatic urothelial cancer: The Canadian Cancer Trials Group BL.12 Randomized Clinical Trial. *JAMA Oncol.* 2020;6(11):1751-1758.
doi: 10.1001/jamaoncol.2020.3927
72. Takebe T, Wells JM. Organoids by design. *Science.* 2019;364(6444):956-959.
doi: 10.1126/science.aaw7567
73. Hammond PT. Shooting for the moon: Nanoscale approaches to cancer. *ACS Nano.* 2016;10(2):1711-1713.
doi: 10.1021/acs.nano.6b01095
74. Choi Y, Cho BK, Seok SH, Kim C, Ryu JH, Kwon IC. Controlled spatial characteristics of ligands on nanoparticles: Determinant of cellular functions. *J Control Release.* 2023;360:672-686.
doi: 10.1016/j.jconrel.2023.07.020
75. Jookan S, Deschaume O, Bartic C. Nanocomposite hydrogels as functional extracellular matrices. *Gels.* 2023;9(2):153.
doi: 10.3390/gels9020153
76. Yang Z, Jiang H. Control of hematopoietic stem cell function through epigenetic regulation of energy metabolism and genome integrity. *Blood.* 2017;130:772.
doi: 10.1182/blood.v130.suppl_1.772.772
77. Teli MK, Mutalik S, Rajanikant GK. Nanotechnology and nanomedicine: going small means aiming big. *Curr Pharm Des.* 2010;16(16):1882-1892.
doi: 10.2174/138161210791208992
78. Wang W, Cai J, Wen J, *et al.* Boosting ferroptosis via abplatin(iv) for treatment of platinum-resistant recurrent ovarian cancer. *Nano Today.* 2022;44:101459.
doi: 10.1016/j.nantod.2022.101459
79. van der Meel R, Sulheim E, Shi Y, Kiessling F, Mulder WJM, Lammers T. Smart cancer nanomedicine. *Nat Nanotechnol.* 2019;14(11):1007-1017.
doi: 10.1038/s41565-019-0567-y
80. Yoshida GJ. Applications of patient-derived tumor xenograft models and tumor organoids. *J Hematol Oncol.* 2020;13(1):4.
doi: 10.1186/s13045-019-0829-z
81. Zazo H, Colino CI, Lanao JM. Current applications of nanoparticles in infectious diseases. *J Control Release.* 2016;224:86-102.
doi: 10.1016/j.jconrel.2016.01.008
82. Hill DR, Spence JR. Gastrointestinal organoids: Understanding the molecular basis of the host-microbe interface. *Cell Mol Gastroenterol Hepatol.* 2016;3(2):138-149.
doi: 10.1016/j.jcmgh.2016.11.007
83. Schlaermann P, Toelle B, Berger H, *et al.* A novel human gastric primary cell culture system for modelling Helicobacter pylori infection in vitro. *Gut.* 2016;65(2):202-213.
doi: 10.1136/gutjnl-2014-307949
84. Stanković A, Mirković M, Radović M, *et al.* Abstracts from the 20th European symposium on radiopharmacy and radiopharmaceuticals. *EJNMMI Radiopharm Chem.* 2023;8(1):11.
doi: 10.1186/s41181-023-00193-4
85. Wang J, Wang F, Jiang Y, Bai L, Chen X, Su J. Organoid-driven nanomedicine platform development. *Biomaterials.* 2026;325:123611.
doi: 10.1016/j.biomaterials.2025.123611
86. Giobbe GG, Bonfante F, Jones BC, *et al.* SARS-CoV-2 infection and replication in human gastric organoids. *Nat Commun.* 2021;12(1):6610.
doi: 10.1038/s41467-021-26762-2
87. Chen YW, Huang SX, de Carvalho ALRT, *et al.* A three-dimensional model of human lung development and disease from pluripotent stem cells. *Nat Cell Biol.* 2017;19(5):542-549.
doi: 10.1038/ncb3510
88. Li Y, Guo Y, Zhang K, *et al.* Cell death pathway regulation by functional nanomedicines for robust antitumor immunity. *Adv Sci (Weinh).* 2024;11(3):e2306580.
doi: 10.1002/advs.202306580
89. Allenspach K. I32 | 3-Dimensional organoid application in veterinary medicine – focus on oncology. *J Vet Pharmacol Ther.* 2023;46(S1):21-23.
doi: 10.1111/jvp.13162
90. Sun M, Lee J, Chen Y, Hoshino K. Studies of nanoparticle delivery with in vitro bio-engineered microtissues. *Bioact*

- Mater.* 2020;5(4):924-937.
doi: 10.1016/j.bioactmat.2020.06.016
91. Wang Y, Jeon H. 3D cell cultures toward quantitative high-throughput drug screening. *Trends Pharmacol Sci.* 2022;43(7):569-581.
doi: 10.1016/j.tips.2022.03.014
 92. Shapiro RL, DeLong K, Zulfiqar F, Carter D, Better M, Ensign LM. In vitro and ex vivo models for evaluating vaginal drug delivery systems. *Adv Drug Deliv Rev.* 2022;191:114543.
doi: 10.1016/j.addr.2022.114543
 93. Palma M, Qi B. Advancing phage therapy: A comprehensive review of the safety, efficacy, and future prospects for the targeted treatment of bacterial infections. *Infect Dis Rep.* 2024;16(6):1127-1181.
doi: 10.3390/idr16060092
 94. Hu T, Brinker CJ, Chan WCW, *et al.* Publishing translational research of nanomedicine in ACS Nano. *ACS Nano.* 2022;16(11):17479-17481.
doi: 10.1021/acsnano.2c10967
 95. Liang W, Zhou C, Liu X, *et al.* Current status of nano-embedded growth factors and stem cells delivery to bone for targeted repair and regeneration. *J Orthop Translat.* 2025;50:257-273.
doi: 10.1016/j.jot.2024.12.006
 96. Abbasalizadeh S, Larijani MR, Samadian A, Baharvand H. Bioprocess development for mass production of size-controlled human pluripotent stem cell aggregates in stirred suspension bioreactor. *Tissue Eng Part C Methods.* 2012;18(11):831-851.
doi: 10.1089/ten.TEC.2012.0161
 97. Carvalho DJ, Kip AM, Tegel A, *et al.* A modular microfluidic organoid platform using LEGO-like bricks. *Adv Healthc Mater.* 2024;13(13):e2303444.
doi: 10.1002/adhm.202303444
 98. Sakamoto JH, van de Ven AL, Godin B, *et al.* Enabling individualized therapy through nanotechnology. *Pharmacol Res.* 2010;62(2):57-89.
doi: 10.1016/j.phrs.2009.12.011
 99. Yang W, Liu H, Liu X, *et al.* Biomimetic fucoidan nanoparticles with regulation of macrophage polarization for targeted therapy of acute lung injury. *Carbohydr Polym.* 2025;351:122967.
doi: 10.1016/j.carbpol.2024.122967
 100. Fancher IS, Rubinstein I, Levitan I. Potential strategies to reduce blood pressure in treatment-resistant hypertension using Food and Drug Administration-approved nanodrug delivery platforms. *Hypertension.* 2019;73(2):250-257.
doi: 10.1161/HYPERTENSIONAHA.118.12005
 101. Ju S, Mu J, Dokland T, *et al.* Grape exosome-like nanoparticles induce intestinal stem cells and protect mice from DSS-induced colitis. *Mol Ther.* 2013;21(7):1345-1357.
doi: 10.1038/mt.2013.64
 102. Smith TT, Stephan SB, Moffett HF, *et al.* In situ programming of leukaemia-specific T cells using synthetic DNA nanocarriers. *Nat Nanotechnol.* 2017;12(8):813-820.
doi: 10.1038/nnano.2017.57
 103. Abdel Fattah AR, Kolaitis N, Van Daele K, Daza B, Rustandi AG, Ranga A. Targeted mechanical stimulation via magnetic nanoparticles guides in vitro tissue development. *Nat Commun.* 2023;14(1):5281.
doi: 10.1038/s41467-023-41037-8
 104. Ye L, Jiang S, Hu J, *et al.* Induction of metabolic reprogramming in kidney by singlet diradical nanoparticles. *Adv Mater.* 2023;35(36):e2301338.
doi: 10.1002/adma.202301338
 105. Tang XY, Wu S, Wang D, *et al.* Human organoids in basic research and clinical applications. *Signal Transduct Target Ther.* 2022;7(1):168.
doi: 10.1038/s41392-022-01024-9
 106. Moradi S, Mahdizadeh H, Šarić T, *et al.* Research and therapy with induced pluripotent stem cells (iPSCs): social, legal, and ethical considerations. *Stem Cell Res Ther.* 2019;10(1):341.
doi: 10.1186/s13287-019-1455-y
 107. Sean G, Baner AJ, Gangaraju R. Organoids and tissue/organ chips. *Stem Cell Res Ther.* 2024;15(1):241.
doi: 10.1186/s13287-024-03859-1
 108. Hong Y, Gao D, Zhao B, Ma J, Yang Z, Guo H. Thermal immuno-nanomedicine: A new strategy for cancer treatment. *Clin Transl Med.* 2023;13(5):e1256.
doi: 10.1002/ctm2.1256
 109. Patton J, Smith E, Smith Z, *et al.* Characterization of cytotoxic macrophages in a pre-clinical model of Epstein-Barr Virus (EBV)-driven lymphoproliferative disease. *Blood.* 2021;138:3502.
doi: 10.1182/blood-2021-154161
 110. Jin H, Xue Z, Liu J, Ma B, Yang J, Lei L. Advancing organoid engineering for tissue regeneration and biofunctional reconstruction. *Biomater Res.* 2024;28:0016.
doi: 10.34133/bmr.0016
 111. Di X, Gao X, Peng L, *et al.* Cellular mechanotransduction in health and diseases: from molecular mechanism to therapeutic targets. *Signal Transduct Target Ther.* 2023;8(1):282.
doi: 10.1038/s41392-023-01501-9
 112. Chhabra A, Song HG, Grzelak KA, *et al.* A vascularized model of the human liver mimics regenerative responses. *Proc Natl Acad Sci USA.* 2022;119(28):e2115867119.

- doi: 10.1073/pnas.2115867119
113. Chooi WH, Ng CY, Ow V, *et al.* Defined alginate hydrogels support spinal cord organoid derivation, maturation, and modeling of spinal cord diseases. *Adv Healthc Mater.* 2023;12(9):e2202342.
doi: 10.1002/adhm.202202342
 114. Ohnsorg ML, Hushka EA, Anseth KS. Photoresponsive chemistries for user-directed hydrogel network modulation to investigate cell-matrix interactions. *Acc Chem Res.* 2025;58(1):47-60.
doi: 10.1021/acs.accounts.4c00548
 115. Thang NH, Chien TB, Cuong DX. Polymer-based hydrogels applied in drug delivery: An overview. *Gels.* 2023;9(7):523.
doi: 10.3390/gels9070523
 116. Singh R, Mori N, Kida YS. Methodological innovations in perfusable vascular networks: Advancing high-density tissue models. *Regen Ther.* 2026;32:101109.
doi: 10.1016/j.reth.2026.101109
 117. Kaur N, Yang D, Ali J. Abstract 1292: Innovative 3D ECM platforms for tumoroid development, drug screening, and personalized treatment. *Cancer Res.* 2025;85(5_Suppl):1292.
doi: 10.1158/1538-7445.am2025-1292
 118. Zhou R, Brislinger D, Fuchs J, *et al.* Vascularised organoids: Recent advances and applications in cancer research. *Clin Transl Med.* 2025;15(3):e70258.
doi: 10.1002/ctm2.70258
 119. Li G, Zhou D, Sheng S, *et al.* Hydrogel for bone microenvironment: Strategy and application. *Chem Eng J.* 2024;499:156554.
doi: 10.1016/j.cej.2024.156554
 120. Luce E, Duclos-Vallee JC. Stem cells and organoids: A paradigm shift in preclinical models toward personalized medicine. *Pharmaceuticals (Basel).* 2025;18(7):992.
doi: 10.3390/ph18070992
 121. Andr s IE, Cho HJ, Toborek M. Abstracts from the International Symposium "Signal Transduction at the Blood-Brain Barriers" 2022. *Fluids Barriers CNS.* 2023;20(1):26.
doi: 10.1186/s12987-023-00417-4
 122. Ma W, Dong Z, Zheng Z, Bai L, Zhang X, Su J. Multiscale construction, evaluation, and application of organoids. *Adv Sci (Weinh).* 2025;12(47):e08534.
doi: 10.1002/advs.202508534
 123. Vallmajo-Martin Q, Broguiere N, Millan C, Zenobi-Wong M, Ehrbar M. PEG/HA hybrid hydrogels for biologically and mechanically tailorable bone marrow organoids. *Adv Funct Mater.* 2020;30(48):1910282.
doi: 10.1002/adfm.201910282
 124. Joddar B, Natividad-Diaz SL, Padilla AE, *et al.* Engineering approaches for cardiac organoid formation and their characterization. *Transl Res.* 2022;250:46-67.
doi: 10.1016/j.trsl.2022.08.009
 125. Shaw CA, Mortimer GM, Deng ZJ, *et al.* Protein corona formation in bronchoalveolar fluid enhances diesel exhaust nanoparticle uptake and pro-inflammatory responses in macrophages. *Nanotoxicology.* 2016;10(7):981-991.
doi: 10.3109/17435390.2016.1155672
 126. Wang C, Ma X. The role of acetylation and deacetylation in cancer metabolism. *Clin Transl Med.* 2025;15(1):e70145.
doi: 10.1002/ctm2.70145
 127. Bilardo R, Traldi F, Vdovchenko A, Resmini M. Influence of surface chemistry and morphology of nanoparticles on protein corona formation. *Wiley Interdiscip Rev Nanomed Nanobiotechnol.* 2022;14(4):e1788.
doi: 10.1002/wnan.1788
 128. Chen C, Leong D, Lynch I. Rethinking nanosafety Part II: Leveraging progress to pioneer new approaches and solutions. *Small.* 2020;16(36):2004934.
doi: 10.1002/sml.202004934
 129. Ghosh S, Pati F. Decellularized extracellular matrix and silk fibroin-based hybrid biomaterials: A comprehensive review on fabrication techniques and tissue-specific applications. *Int J Biol Macromol.* 2023;253(Pt 8):127410.
doi: 10.1016/j.ijbiomac.2023.127410
 130. Khodayari H, Khodayari S, Rezaee M, *et al.* Promotion of cardiac microtissue assembly within G-CSF-enriched collagen I-cardiogel hybrid hydrogel. *Regen Biomater.* 2024;11:rbae072.
doi: 10.1093/rb/rbae072
 131. Khetani SR, Berger DR, Ballinger KR, Davidson MD, Lin C, Ware BR. Microengineered liver tissues for drug testing. *SLAS Technol.* 2015;20(3):216-250.
doi: 10.1177/2211068214566939
 132. Zhao R, Feng Q, Xia Y, Liao L, Xie S. Iteration of tumor organoids in drug development: Simplification and integration. *Pharmaceuticals.* 2025;18(10):1540.
doi: 10.3390/ph18101540
 133. Jun HR, Chang SY, Kim H, *et al.* Abstract 5199: Enhancing precision medicine in advanced lung, gastric and rectal cancer: Exploring the reliability of Patient-derived organoids. *Cancer Res.* 2025;85(5_Suppl):5199.
doi: 10.1158/1538-7445.am2025-5199
 134. Vives J, Batlle-Morera L. The challenge of developing human 3D organoids into medicines. *Stem Cell Res Ther.* 2020;11(1):72.
doi: 10.1186/s13287-020-1586-1
 135. Sato T, Vries RG, Snippert HJ, *et al.* Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal

- niche. *Nature*. 2009;459(7244):262-265.
doi: 10.1038/nature07935
136. Franco C, Rezzani R. Methods and models for studying mycobacterium tuberculosis in respiratory infections. *Int J Mol Sci*. 2025;26(1):18.
doi: 10.3390/ijms26010018
137. Lin YJ, Komperda I. Biomimetic targeting and autonomous optimization in nanomedicine. *ChemRxiv*. Preprint posted online 2025.
doi: 10.26434/chemrxiv-2025-fxnmb
138. Lancaster MA, Renner M, Martin CA, *et al*. Cerebral organoids model human brain development and microcephaly. *Nature*. 2013;501(7467):373-379.
doi: 10.1038/nature12517
139. Huch M, Gehart H, van Boxtel R, *et al*. Long-term culture of genome-stable bipotent stem cells from adult human liver. *Cell*. 2015;160(1):299-312.
doi: 10.1016/j.cell.2014.11.050